

Universality in p -spin glasses with correlated disorder

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

J. Stat. Mech. (2013) L02003

(<http://iopscience.iop.org/1742-5468/2013/02/L02003>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 194.94.224.254

The article was downloaded on 25/03/2013 at 10:41

Please note that [terms and conditions apply](#).

LETTER

Universality in p -spin glasses with correlated disorder

Valentin Bonzom¹, Razvan Gurau¹ and Matteo Smerlak²

¹ Perimeter Institute for Theoretical Physics, 31 Caroline Street N,
ON N2L 2Y5, Waterloo, Canada

² Max-Planck-Institut für Gravitationsphysik, Am Mühlenberg 1,
D-14476 Golm, Germany

E-mail: vbonzom@perimeterinstitute.ca, rgurau@perimeterinstitute.ca
and smerlak@aei.mpg.de

Received 21 December 2012

Accepted 2 February 2013

Published 27 February 2013

Online at stacks.iop.org/JSTAT/2013/L02003

[doi:10.1088/1742-5468/2013/02/L02003](https://doi.org/10.1088/1742-5468/2013/02/L02003)

Abstract. We introduce a new method, based on the recently developed *random tensor theory*, for studying the p -spin glass model with non-Gaussian, correlated disorder. Using a suitable generalization of Gurau's theorem on the universality of the large N limit of the p -unitary ensemble of random tensors, we exhibit an infinite family of such non-Gaussian distributions which leads to the *same* low temperature phase as the Gaussian distribution. While this result is easy to show (and well known) for uncorrelated disorder, its robustness with respect to strong quenched correlations is surprising. We show in detail how the critical temperature is renormalized by these correlations. We close with speculation on possible applications of random tensor theory to finite-range spin glass models.

Keywords: disordered systems (theory), spin glasses (theory), random matrix theory and extensions

ArXiv ePrint: [1206.5539](https://arxiv.org/abs/1206.5539)

Contents

1. Introduction	2
1.1. p -spin glass models	3
1.2. Large random tensors	4
1.3. Universality in the couplings	6
2. Conclusion and outlook	7
Acknowledgments	8
References	8

1. Introduction

It is well known that the phase diagram of the p -spin glass model [1, 2] does not depend on the details of the disorder distributions, in the following sense: if $J_{i_1 \dots i_p}$ denotes a set of independent and identically distributed p -valent couplings between sites $i_1 \dots i_p$, a non-quadratic potential $V(J_{i_1 \dots i_p})$ in the coupling distribution

$$\prod_{i_1 \dots i_p} dJ_{i_1 \dots i_p} e^{-J_{i_1 \dots i_p}^2 / \sigma^2 + V(J_{i_1 \dots i_p})} \quad (1)$$

is irrelevant in the thermodynamic limit. That a similar result would hold for *correlated* disorder distributions, with terms such as

$$\sum_{\{i_l, j_l\}} J_{i_1 i_2 i_3} J_{i_1 j_2 j_3} J_{j_1 j_2 j_3} J_{j_1 i_2 i_3}, \quad (2)$$

in the potential, is much less obvious. In fact, to our knowledge, no analytic framework for dealing with such correlated, non-Gaussian disorder has been reported so far. Since disorder correlations are to be expected in actual physical systems, understanding their effect is an important problem.

In this work we exhibit an infinite class of non-Gaussian terms of the kind (2) such that (i) the thermodynamic limit $N \rightarrow \infty$ is exactly soluble, and (ii) the spin glass phase has the same structure as for uncorrelated disorder, except for a renormalization of the critical temperature. This provides the first general result on spin glasses with strongly correlated disorder.

Our approach is based on new results in *random tensor theory*. A natural generalization of random matrices—random tensors—have recently been shown to possess a large N limit [3] dominated by a few, well-identified ‘melonic’ graphs (the tensor equivalent of ’t Hooft’s planar graphs in matrix theory [4]). Furthermore, the melonic family can actually be resummed exactly, and turns out to exhibit interesting critical and multicritical behavior [5]. These results have not been applied to spin glass problems previously, and our hope in this work is to convey that random tensors are potentially powerful tools for spin glass theory, just like random matrices [6].

From the perspective of random tensor theory, the quenched couplings of spin glasses with p -spin interactions are non-Gaussian rank- p random tensors. The behavior of such tensors in the large N limit has been investigated in [7, 8], with a striking conclusion: in a suitable ensemble with p -unitary symmetry (more details are given in the text), this limit is *universally Gaussian*. This means that, in this ensemble, in the large N limit the sole effect of the self-interactions of large tensors is to dress the propagator. Here, we show how this result can be generalized to include interactions between tensors and spin variables, and thus obtain the aforementioned universality result.

This work is organized as follows. We first recall the Hamiltonian for p -spin models and insist on the need for a correlated disorder. Then, we recall the relevant properties of large random tensors in the p -unitary ensemble. This enables us to show how non-Gaussian, correlated, quenched variables can be integrated exactly in the large N (thermodynamic) limit, yielding our universality theorem. We conclude with a few words on the possible relevance of tensor techniques for finite-range p -spin glasses.

1.1. p -spin glass models

We consider a p -spin Hamiltonian [1, 2]

$$H_J(S) = - \sum_{1 \leq i_1 \dots i_p \leq N} J_{i_1 \dots i_p} S_{i_1} \dots S_{i_p} + c.c. \quad (3)$$

where $J_{i_1 \dots i_p}$ is a complex³ tensor describing the couplings and $S = (S_i)_{1 \leq i \leq N}$ is a set of real spins⁴ with lattice index i , weighted by a (normalized) probability measure $d\Omega(S)$ such that

$$\int d\Omega(S) \sum_{i=1}^N S_i^2 = \mathcal{O}(N). \quad (4)$$

This includes in particular Ising [1] and spherical [2] spins.

When the couplings are Gaussianly distributed, it is well known that such p -spin glass models exhibit replica symmetry breaking in the low temperature phase [2, 11, 12] and have a dynamical transition at a higher temperature where a large number of metastable states (growing exponentially with N) dominates the free energy landscape [13, 14]; their relevance is conjectured to extend to structural glasses [15]. These results extend easily to the case of independent and identically distributed (i.i.d.) couplings: all terms of order higher than 2 in $J_{i_1 \dots i_p}$ and $\bar{J}_{i_1 \dots i_p}$ in the measure on (J, \bar{J}) are irrelevant in the thermodynamic limit $N \rightarrow \infty$.

In this work we aim to study a family of *correlated* non-Gaussian measures on the disorder. Physically, randomness of the couplings comes from randomness of the positions of the spins, and in general we should not expect the couplings between different sets of p spins to be independent (for instance due to the geometric relations between the positions of the spins). One should therefore perturb the Gaussian distribution (with covariance σ^2)

³ The use of complex rather than real tensors is motivated by purely technical convenience and does not change the physics in any way.

⁴ It is also possible to include Potts or vector spins coupled according to some fixed multi-linear map.

on the couplings with a polynomial $V(J, \bar{J})$. The quenched free energy is given by

$$[F(J, \bar{J})] = \frac{\int dJ d\bar{J} e^{-N^{p-1}(J \cdot \bar{J}/\sigma^2 + V(J, \bar{J}))} F(J, \bar{J})}{\int dJ d\bar{J} e^{-N^{p-1}(J \cdot \bar{J}/\sigma^2 + V(J, \bar{J}))}}, \quad (5)$$

with

$$-\beta F(J, \bar{J}) = \ln \int d\Omega(S) e^{-\beta H_J(S)}, \quad (6)$$

and $J \cdot \bar{J}$ is shorthand for $\sum_{j_1 \dots j_p} J_{j_1 \dots j_p} \bar{J}_{j_1 \dots j_p}$.

The evaluation of $[F]$ for a generic potential V is of course a completely open problem. However, the theory of large random tensors provides an exact calculation for an infinite family of potentials satisfying a particular kind of invariance.

1.2. Large random tensors

We now review the relevant properties of large random tensors discovered in [3, 5, 7]. The first obvious observation is that, unlike symmetric/Hermitian matrices, tensors cannot be diagonalized. Hence, a key concept in random matrix theory, the eigenvalue distribution, does not carry over to the case of higher rank. It turns out however that this fact does not preclude the development of random tensor theory, which in fact relies on the identification of an *ensemble* with suitable symmetry properties.

One such ensemble of tensors—indeed the only one identified so far—is the p -unitary ensemble, defined as follows. Consider a rank- p complex tensor in N dimensions J , with components $J_{i_1 \dots i_p}$ in a fixed basis, and for each set of p unitary matrices $U^{(1)}$ to $U^{(p)}$, define

$$J'_{i_1 \dots i_p} = \sum_{j_1 \dots j_p} U_{i_1 j_1}^{(1)} \cdots U_{i_p j_p}^{(p)} J_{j_1 \dots j_p}. \quad (7)$$

Then we say that a function $V(J, \bar{J})$ of J and its complex conjugate \bar{J} is a p -unitary invariant if⁵

$$V(J', \bar{J}') = V(J, \bar{J}). \quad (8)$$

The set of p -unitary invariants is conveniently parameterized by p -bubbles B , that is p -valent bipartite connected graphs with edges colored by numbers between 1 and p , such that each ‘color’ is incident exactly once to each vertex; see figure 1. A bubble represents an invariant denoted as $\text{tr}_B(J, \bar{J})$, by associating a tensor J with each ‘white’ vertex of B and a conjugate \bar{J} to each ‘black’ vertex, and contracting their k th indices along the edges colored with k . By the fundamental theorem of classical invariants of $U(N)$ (see for instance [9]), a general p -unitary invariant can be expanded as

$$V(J, \bar{J}) = \sum_B t_B \text{tr}_B(J, \bar{J}), \quad (9)$$

where the t_B are coupling constants.

⁵ The corresponding symmetry group is known as the external tensor product of p copies of $U(N)$.




Figure 1. Some p -bubbles at $p = 3$. Up to color relabeling, there is a single bubble with four vertices (on the left), whose invariant is $\sum_{\{i_l, j_l\}} J_{i_1 i_2 i_3} \bar{J}_{i_1 j_2 j_3} J_{j_1 j_2 j_3} \bar{J}_{j_1 i_2 i_3}$. But there exist different bubbles with six vertices (the three on the right).

For a given invariant potential $V(J, \bar{J})$, we define the average of $f(J, \bar{J})$ over J by

$$\left[f(J, \bar{J}) \right] = \frac{\int dJ d\bar{J} e^{-N^{p-1}(J \cdot \bar{J} / \sigma^2 + V(J, \bar{J}))} f(J, \bar{J})}{\int dJ d\bar{J} e^{-N^{p-1}(J \cdot \bar{J} / \sigma^2 + V(J, \bar{J}))}}. \quad (10)$$

The Feynman diagrammatic expansion of these quantities involves $(p+1)$ -colored bipartite graphs, made of p -bubbles connected together via extra lines with color ‘0’ incident on each vertex and corresponding to the propagator σ^2 in (10).

The following results concerning the large N limit of (10) have been proved:

- The Feynman expansion is dominated in the large N limit by a simple class of graphs, called *melonic graphs*, which generalize ’t Hooft’s planar graphs [3]. Intuitively, a $(p+1)$ -colored graph is melonic if it can be built from recursive insertions on any line of two vertices connected together by p lines, as in figure 2.
- The large N limit is *Gaussian*, in the sense that up to subleading corrections in $1/N$,

$$\left[\text{tr}_B(J, \bar{J}) \right] = N G_2^{|B|/2}, \quad (11)$$

where $|B|$ is the number of vertices of the bubble B and $G_2 = [J \cdot \bar{J}] / N$ is the *dressed* propagator depending on the potential V [7].

- The following Schwinger–Dyson equation holds in the $N \rightarrow \infty$ limit [8]:

$$\frac{[J \cdot \bar{J}]}{\sigma^2 N} + \sum_B t_B \frac{|B|}{2} \frac{[\text{tr}_B(J, \bar{J})]}{N} = 1. \quad (12)$$

The first result implies that all non-melonic bubbles B in the potential drop out in the large N limit, and therefore we can restrict the sum in (9) to melonic bubbles (and hence hereafter B will always denote a melonic bubble). In figure 1, all bubbles are melonic except the non-planar one on the right.

The second result has been coined the *universality* property of the p -unitary ensemble of random tensors, and can be seen as a non-trivial generalization of the central limit theorem. Its origin is that there is only one way to dress a melonic bubble B with propagators in a melonic way, which happens to correspond to Gaussian contractions. This feature is specific to tensors and does not hold for random matrices. In a way, this work can be read as the physics counterpart of this surprising mathematical result. We refer the reader to the review [10] and to the original papers for more details on random tensor theory.




Figure 2. A patch of a melonic graph, recursively built by inserting on any line a pair of vertices, a black one and a white one, connected together via p lines (here $p = 3$).

1.3. Universality in the couplings

Let us now come back to spin glasses. Following the standard recipe for computing quenched quantities [17], we consider the averaged replicated partition function

$$[Z^n] = \int \prod_{a=1}^n d\Omega(S^a) e^{-\beta H_{\text{eff}}(\{S^a\})}, \quad (13)$$

where a is the replica index and the effective Hamiltonian is defined by

$$e^{-\beta H_{\text{eff}}(\{S^a\})} = \left[e^{-\beta \sum_{a=1}^n H_J(S^a)} \right]. \quad (14)$$

In diagrammatic language, H_{eff} is given by the sum over all connected $(p+1)$ -colored bipartite graphs (henceforth ‘graphs’) with spins S_i^a on the external legs. Denoting as k the order of the effective coupling between replicas $a_1 \cdots a_k$, this can be pictured as

$$-\beta H_{\text{eff}}(\{S^a\}) = \sum_k \beta^k \sum_{a_1 \dots a_k} \sum_{a_k} \text{Diagram} \quad (15)$$

Here the solid line is the J -propagator (tensor lines with color 0), and the p dashed line emerging from each external leg represents the external spin variables $S_{i_l}^{a_k}$. The blob G_k is the large N tensor connected k -point function, i.e. the sum over all connected melonic graphs with k external (solid) legs. For each graph contributing to the blob amplitude, the site indices i_l of the spins are contracted along ‘broken faces’, i.e. connected paths with alternating color $1 \leq c \leq p$ and 0, from one external dashed leg to another, through the graph.

Let us now show that the $k = 2$ term dominates in the large N limit. Observe that powers of N in $H_{\text{eff}}(\{S^a\})$ have three sources: the tensor propagators, the bubble interactions $\text{tr}_B(J, \bar{J})$, and the sums over site indices i . The first two contributions are those of a melonic graph with k cut lines of color 0 whose scaling has been found in the appendix of [7] to be $p - (p-1)k - \rho$, where ρ is a positive number independent of k . As for the spin contribution, from (4) we see that it gives at most a factor of N per broken face, and there are at most $pk/2$ of them. This gives for the scaling degree $\omega(k)$ in N of the order- k term of (15)

$$\omega(k) \leq p - \left(\frac{p}{2} - 1\right) k. \quad (16)$$

models are judged (at least partially) unphysical. We expect however that random tensor techniques should be applicable to finite-range models too. Indeed, from the random tensor perspective, a finite-range spin glass model is one for which the J -propagator is non-trivial, and in particular depends on the tensor indices of J . A typical example of interest here would be

$$\sigma_{i_1 \dots b_i p; j_1 \dots b_j p}^2 = \frac{\prod_{k=1}^p \delta_{i_k j_k}}{\sum_{1 \leq l < k \leq p} (i_l - j_k)^2 + 1}, \quad (22)$$

which goes to zero when the lattice sites are far away. Such tensor models have already been considered in the context of quantum gravity [18], where they have been called *tensor field theories* (TFT). The key difference between TFT and the simple tensor models considered in this work is the appearance of a renormalization flow. The first renormalizable TFT has been identified in [19], and developments are happening quickly in this area. We expect that these new techniques will prove useful in the difficult field of finite-range spin glass theory.

Acknowledgments

The authors thank Romain Mari for his careful reading of the manuscript and Francesco Zamponi for useful comments. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research and Innovation. The research leading to these results received funding from the European Union Seventh Framework Programme FP7-People-2010-IRSES under grant agreement 269217.

References

- [1] Derrida B, *Random-energy model: an exactly solvable model of disordered systems*, 1981 *Phys. Rev. B* **24** 2613
- [2] Crisanti A and Sommers H J, *The spherical p -spin interaction spin glass model: the statics*, 1992 *Z. Phys. B* **87** 341
- [3] Gurau R, *The $1/N$ expansion of colored tensor models*, 2011 *Ann. Henri Poincaré* **12** 829 [arXiv:1011.2726]
Gurau R and Rivasseau V, *The $1/N$ expansion of colored tensor models in arbitrary dimension*, 2011 *Europhys. Lett.* **95** 50004 [arXiv:1101.4182]
Gurau R, *The complete $1/N$ expansion of colored tensor models in arbitrary dimension*, 2012 *Annales Henri Poincaré* **13** 399 [arXiv:1102.5759]
- [4] 't Hooft G, *A planar diagram theory for strong interactions*, 1974 *Nucl. Phys. B* **72** 461
- [5] Bonzom V, Gurau R, Riello A and Rivasseau V, *Critical behavior of colored tensor models in the large N limit*, 2011 *Nucl. Phys. B* **853** 174 [arXiv:1105.3122]
- [6] Kosterlitz J, Thouless D and Jones R, *Spherical model of a spin-glass*, 1976 *Phys. Rev. Lett.* **36** 1217
- [7] Gurau R, 2011 *Universality for random tensors* arXiv:1111.0519
- [8] Bonzom V, Gurau R and Rivasseau V, *Random tensor models in the large N limit: uncoloring the colored tensor models*, 2012 *Phys. Rev. D* **85** 084037 [arXiv:1202.3637]
- [9] Collins B, *Moments and cumulants of polynomial random variables on unitary groups, the Itzykson–Zuber integral, and free probability*, 2003 *Int. Math. Res. Not.* **17** 953 [arXiv:math-ph/0205010]
- [10] Gurau R and Ryan J, *Colored tensor models—a review*, 2012 *SIGMA* **8** 020 [arXiv:1109.4812]
- [11] Gross D and Mezard M, *The simplest spin glass*, 1984 *Nucl. Phys. B* **240** 431
- [12] Gardner E, *Spin glasses with p -spin interactions*, 1985 *Nucl. Phys. B* **257** 747
- [13] Crisanti A and Sommers H J, *Thouless–Anderson–Palmer approach to the spherical p -spin spin glass model*, 1995 *J. Physique I* **5** 805

- [14] Crisanti A, Leuzzi L and Rizzo T, *Complexity in mean-field spin-glass models: Ising p -spin*, 2005 *Phys. Rev. B* **71** 094202 [arXiv:cond-mat/0406649]
- [15] Kirkpatrick T R and Thirumalai D, *Dynamics of the structural glass transition and the p -spin-interaction spin-glass model*, 1987 *Phys. Rev. Lett.* **58** 2091
- [16] Castellani T and Cavagna A, *Spin-glass theory for pedestrians*, 2005 *J. Stat. Mech.* **P05012** 2005 [arXiv:cond-mat/0505032]
- [17] Mezard M, Parisi G and Virasoro M, 1986 *Spin Glass Theory and Beyond* (Singapore: World Scientific)
- [18] Rivasseau V, *Quantum gravity and renormalization: the tensor track*, 2011 arXiv:1112.5104
- [19] Geloun J B and Rivasseau V, *A renormalizable 4-dimensional tensor field theory*, arXiv:1111.4997