

The very energetic, broad-lined Type Ic supernova 2010ah (PTF10bzf) in the context of GRB/SNe

Paolo A. Mazzali,^{1,2,3*} Emma S. Walker,^{4,5} Elena Pian,^{5,6} Masaomi Tanaka,⁷
Alessandra Corsi,⁸ Takashi Hattori⁹ and Avishay Gal-Yam¹⁰

¹INAF–Osservatorio Astronomico, vicolo dell’Osservatorio, 5, I-35122 Padova, Italy

²Astrophysics Research Institute, Liverpool John Moores University, CH41 1LD Liverpool, UK

³Max-Planck-Institut für Astrophysik, Karl-Schwarzschildstr. 1, D-85748 Garching, Germany

⁴Yale University, New Haven, CT, USA

⁵Scuola Normale Superiore, 7, I-56126 Pisa, Italy

⁶INAF–IASF Bologna, via Gobetti 101, I-40129 Bologna, Italy

⁷National Astronomical Observatory, 2-21-1 Osawa, Mitaka, Tokyo, Japan

⁸George Washington University, Washington, D.C. 20052, USA

⁹Subaru Telescope, National Astronomical Observatory of Japan, Hilo, HI 96720, USA

¹⁰Weizmann Institute of Science, Rehovot 76100, Israel

Accepted 2013 April 9. Received 2013 April 6; in original form 2013 February 24

ABSTRACT

SN 2010ah, a very broad-lined Type Ic supernova (SN) discovered by the Palomar Transient Factory, was interesting because of its relatively high luminosity and the high velocity of the absorption lines, which was comparable to that of gamma-ray burst (GRB)/SNe, suggesting a high explosion kinetic energy. However, no GRB was detected in association with the SN. Here, the properties of SN 2010ah are determined with higher accuracy than previous studies through modelling. New *Subaru* telescope photometry is presented. A bolometric light curve is constructed taking advantage of the spectral similarity with SN 1998bw. Radiation transport tools are used to reproduce the spectra and the light curve. The results thus obtained regarding ejecta mass, composition and kinetic energy are then used to compute a synthetic light curve. This is in reasonable agreement with the early bolometric light curve of SN 2010ah, but a high abundance of ^{56}Ni at high velocity is required to reproduce the early rise, while a dense inner core must be used to reproduce the slow decline at late phases. The high-velocity ^{56}Ni cannot have been located on our line of sight, which may be indirect evidence for an off-axis, aspherical explosion. The main properties of SN 2010ah are: ejected mass $M_{\text{ej}} \approx 3 M_{\odot}$; kinetic energy $E_{\text{kin}} \approx 10^{52}$ erg, $M(^{56}\text{Ni}) \approx 0.25 M_{\odot}$. The mass located at $v \gtrsim 0.1 c$ is $\sim 0.2 M_{\odot}$. Although these values, in particular the E_{kin} , are quite large for a Type Ic SN, they are all smaller (especially M_{ej}) than those typical of GRB/SNe. This confirms the tendency for these quantities to correlate, and suggests that there are minimum requirements for a GRB/SN, which SN 2010ah may not meet although it comes quite close. Depending on whether a neutron star or a black hole was formed following core collapse, SN 2010ah was the explosion of a CO core of ~ 5 to $6 M_{\odot}$, pointing to a progenitor mass of ~ 24 – $28 M_{\odot}$.

Key words: radiation mechanisms: thermal – supernovae: general – supernovae: individual: SN 2010ah.

1 INTRODUCTION

Type Ib/c supernovae (SNe Ib/c) are perhaps the most diverse sub-type of SNe. Their spectroscopic definition [no H (Type Ib), no He (Type Ic), weak Si, see Filippenko (1997)] means that this is the class in which almost all ‘strange’ SNe are confined. SNe Ib/c include

all explosions of massive stars that have lost their outer envelopes. The diversity of possible progenitors and evolutionary histories is reflected in the huge variety of properties of SNe Ib/c, which range from low (‘normal’?) luminosity and kinetic energy (e.g. SN 1994I; Iwamoto et al. 1994; Richmond et al. 1996; Sauer et al. 2006) to broad-lined SNe Ic (e.g. SN 2002ap; Gal-Yam, Ofek & Shemmer 2002; Mazzali et al. 2002; Foley et al. 2003), to X-ray Flash (XRF)-connected SNe Ib (e.g. SN 2008D; Mazzali et al. 2008; Modjaz et al. 2008; Soderberg et al. 2008) and SNe Ic (e.g. SN 2006aj; Pian et al.

*E-mail: mazzali@mpa-garching.mpg.de

2006), to luminous and energetic gamma-ray burst (GRB)/SNe Ic [SN 1998bw, Galama et al. (1998); SN 2003dh, Stanek et al. (2003); Hjorth et al. (2003); Matheson et al. (2003); SN 2003lw, Maesani et al. (2004)]. The SN I Ib subtype, where a small fraction of the H envelope survives, showing up in the spectrum but leaving practically no imprint on the light curve (LC; Hachinger et al. 2012), also belongs to the same physical group. SNe I Ib are also known to range from low (e.g. SN 1993J; Filippenko, Matheson & Ho 1993) to high energy events (e.g. SN 2003bg; Hamuy et al. 2009). Binarities are suspected to play a major role in the evolution of their progenitors (Nomoto, Iwamoto & Suzuki 1995; Maund et al. 2004, 2011; Crockett et al. 2008; Arcavi et al. 2011; Van Dyk et al. 2011). Some very bright SNe Ic are possibly linked to the pair instability phenomenon in very massive stars (e.g. the SN Ic 2007bi; Gal-Yam et al. 2009). Among the members of this class are also very luminous SNe whose origin is still debated (e.g. SN 2005ap, Quimby et al. 2007, 2011; Gal-Yam 2012, and references therein or PS1-10afx, Chornock et al. 2013). Finally, at the opposite end of the distribution, very dim SNe Ib such as 2005E and other ‘Ca-rich SNe’ (Perets et al. 2010; Kasliwal et al. 2012) may be the result of the explosive ejection of a shell from the surface of a He-accreting white dwarf. All these SNe are also formally members of the SN Ib/c class, but their physical origin can be very different.

Still, the connection between SNe Ib/c, massive stars and GRBs is one of the most important reasons to study these stellar explosions. While extremely massive stars ($M \geq 80 M_{\odot}$) are believed to explode via the pair instability mechanism, which leaves no remnant, stars more massive than $\sim 25 M_{\odot}$ should collapse to form black holes. This mechanism is thought to be responsible for the emission of long-soft GRBs (Woosley 1993). Although the details of the energy extraction are not yet understood, there is sufficient energy to power a relativistic jet (MacFadyen & Woosley 1999; Bromberg et al. 2011). In fact, all well-observed GRB/SNe are broad-lined SNe Ic, and all have been inferred to have large ejected mass and kinetic energy (Iwamoto et al. 1998; Mazzali et al. 2003, 2006b). These quantities are derived from modelling the LCs and spectra of these SNe, and the results consistently yield ejected masses of $\sim 10 M_{\odot}$, kinetic energies of several 10^{52} erg, ^{56}Ni masses of $\sim 0.5 M_{\odot}$, which have been used to infer progenitor star masses of 35–50 M_{\odot} .

On the other hand, X-ray Flash/SNe like SN 2006aj seem to come from stars of lower mass ($\sim 20 M_{\odot}$), suggesting that energy injection from a magnetar is the mechanism which originated the XRF and energized the explosion (Mazzali et al. 2006a).

There are various cases of broad-lined SNe Ic whose properties are intermediate between those of GRB/SNe and those of XRF/SNe and are apparently not accompanied by a relativistic outflow [e.g. SN 1997ef, Mazzali, Iwamoto & Nomoto (2000); SN 1997dq, Mazzali et al. (2004); SN 2002ap, Mazzali et al. (2002)]. This may suggest that some of the properties of these SNe were not sufficient for the generation of a GRB. Alternatively, they may all have off-axis GRBs, but this seems to be ruled out by radio observations (Soderberg et al. 2006). Studies of the emission line profiles in the late, nebular phase (Mazzali et al. 2004, 2007, 2010) do not suggest the off-axis option for most of these SNe, although they clearly do so for some of them (Mazzali et al. 2005; Tanaka et al. 2008; van den Horst et al. 2011). Spectropolarimetry, which is available only in very few cases, also suggests that SNe Ic are aspherical (e.g. Kawabata et al. 2002, 2003; Gorosabel et al. 2006, 2010), and that more energetic SNe are more aspherical than lower energy ones (Tanaka et al. 2012).

Clearly, in order to improve our understanding of these explosions, we need to extend the observed sample and cover a

broader range of SN properties. A particularly interesting domain is SNe Ib/c with properties straddling those of GRB- and non-GRB/SNe. The presence of a high-energy transient is certainly one of the dimensions that determine SN Ib/c properties, although not the only one. One of the few SNe that falls in this category is SN 2010ah/PTF10bzf, a bright, broad-lined SN Ic without a GRB, for which extremely high velocities were inferred (Corsi et al. 2011). In this paper, we model the spectra and LC of SN 2010ah with our Monte Carlo radiation transport codes and determine its properties. Our work is based on the data presented by Corsi et al. (2011), but we add very accurate photometry obtained with the *Subaru* telescope.

The paper is organized as follows. In Section 2, we recap the properties of SN 2010ah. In Section 3, we present the new *Subaru* photometry. In Section 4, we discuss how we use the spectral similarity to SN 1998bw to build a bolometric LC for SN 2010ah given the lack of multicolour photometry. The LC of SN 2010ah is then compared to those of other hypernovae (HNe). In Section 5, we briefly recap our modelling method and present spectral models for SN 2010ah. In Section 6, the synthetic LCs computed based on the spectral modelling results are presented and compared to the bolometric LC of SN 2010ah. Finally, Section 7 contains a discussion and conclusions.

2 PTF10BZF = SN 2010AH: OBSERVATIONAL PROPERTIES

The bright, broad-lined SN Ic SN 2010ah was discovered by the Palomar Transient Factory (PTF; Law et al. 2009; Rau et al. 2009) on 2010 February 23 in a galaxy at a redshift $z = 0.0498$ (Corsi et al. 2011). It reached $r = 18.3$ mag on 2010 March 2. At a distance modulus of 36.7 mag, this corresponds to an absolute magnitude $M_R \approx -18.4$ mag. This makes SN 2010ah a luminous SN, but not a record breaker. Data from the PTF follow-up were published in Corsi et al. (2011). The SN was classified as a Type Ic from spectra obtained at Gemini-N and Keck Obs, which show very broad lines. Velocities in excess of $0.1c$ were measured in Ca II and Si II lines (Corsi et al. 2011, fig. 5). In fact, absorption lines in SN 2010ah are among the broadest ever, rivalling with GRB/SNe such as 1998bw. Corsi et al. (2011) suggested that the SN is intermediate in spectral properties between the GRB/SN 1998bw (Patat et al. 2001) and the non-GRB SN 1997ef (Mazzali et al. 2000). Actually, SN 1998bw is the closest match, as can be seen from figs 2 and 3 of Corsi et al. (2011). However, no X-ray counterpart was detected for SN 2010ah.

The striking spectral similarity to SN 1998bw and the lack of an associated γ - or X-ray counterpart make SN 2010ah an interesting event. The *R*-band luminosity was comparable to that of other broad-lined SNe Ic without a GRB, like SN 2003jd or SN 2009bb, for both of which an off-axis GRB has been suggested (Mazzali et al. 2005; Soderberg et al. 2010, respectively). However, SNe 2003jd and 2009bb resemble one another spectroscopically and look much more like the XRF/SN 2006aj (Pignata et al. 2011, figs 11 and 12) than SN 2010ah or SN 1998bw.

In Fig. 1, we compare the two available spectra of SN 2010ah with the spectra of SNe 1998bw and 2002ap that look most similar based on an eye estimate of the available ones. The overall similarity is apparent: all three SNe show extreme line blending. In particular, Ca II IR and O I 7774 Å are blended, which requires sufficient ejecta with $v \gtrsim 0.1c$. In blue, most pseudo-emission peaks (which actually correspond to regions of low line opacity where photons can escape) are obliterated because of line blocking by high-velocity material (Fe, Cr and Ti). This effect is more apparent in SN 1998bw than




Figure 1. Comparison of the spectra of SNe 2010ah, 1998bw (Patat et al. 2001) and 2002ap (William Herschel Telescope service programmes) at similar epochs relative to the respective LC evolution. Ions responsible for the strongest spectral features are marked.

in either SN 2002ap or 2010ah, indicating a higher metal content at high velocity. On the other hand, SN 2010ah seems to have the highest line velocity of the three SNe (Corsi et al. 2011). This is shown in particular by the blueshift of the Si II 6355 Å absorption in the March 7 spectrum, which has $v \approx 18\,000$ km s⁻¹. Considering that the epoch of the spectrum is at least 12 d after explosion (Corsi et al. 2011), this velocity is exceptionally high, and similar only to GRB/SNe (Pian et al. 2006; Bufano et al. 2012).

The spectra of SN 2010ah and SN 2002ap used for the comparison are closer to one another in time than those of SN 1998bw. Based on the epochs of the spectra it can be roughly estimated that the time evolution of SN 2010ah is faster than that of SN 1998bw by about a factor of 1.3. In Section 4, we show how this is in fact confirmed by the evolution of the LCs.

3 NEW *Subaru* PHOTOMETRY AND THE LIGHT CURVES OF SN 2010AH

The PTF follow-up campaign collected a good *r*-band LC of SN 2010ah, but other bands were poorly sampled (Corsi et al. 2011). New *Subaru* data presented here improve the coverage in the post-maximum phase and make the calculation of a bolometric LC more reliable.

Imaging observations of SN 2010ah in the *B*, *V* and *R* bands were performed under non-photometric conditions with the *Subaru* telescope equipped with the Faint Object Camera and Spectrograph (FOCAS; Kashikawa et al. 2002) on 2010 March 19.25, 20.25, 21.25 and 22.24 UT (MJD = 552 74.25, 552 75.25, 552 76.25 and 552 77.24, respectively). The exposure time was 2 s for each image. The images were bias subtracted and flat fielded. Photometry was performed relative to five stars in the field for the *R*- and *V*-band images and to four stars for the *B*-band images. We used an aperture of 1.5 times the full width at half-maximum (FWHM) of the point spread function of each image, estimated as the mean of the FWHM




Figure 2. The UBVRI LCs of SN 2010ah. The LCs, dereddened with $E(B - V) = 0.012$, include the data presented in Corsi et al. (2011) and the new *Subaru* photometry (larger dots). The lines show the LCs of SN 1998bw, dereddened with $E(B - V) = 0.016$, dimmed by 0.5 mag and stretched in time by a factor of 1.3^{-1} . The LCs of SN 2010ah approximately match those of SN 1998bw.

measured over the reference stars and SN 2010ah. The magnitudes of the reference stars, originally in the Sloan Digital Sky Survey system, were calibrated to *BVR* using the conversions described in Jordi, Grebel & Ammon (2006). Colour corrections were applied to each image assuming that the measurement using the filter at the adjacent longer wavelength contains the information required to perform the correction. Zero-point and colour-term coefficients were derived performing a linear fit on the reference stars for each image, and then used to derive the SN photometry for the corresponding image. The $B - V$, $V - R$ and $R - I$ colours of SN 2010ah were derived from *WISE* photometry (Corsi et al. 2011) at epochs close to those of the *Subaru* images. Since only two reference stars could be identified in the *B*-band image of 19.25 March, the zero-point and colour term could not be determined, and a magnitude from this image could not be derived.

Fig. 2 shows the multiband photometry of SN 2010ah from Corsi et al. (2011) and the new *Subaru* data, highlighted as large circles. In order to build this figure, the photometric data originally acquired with filters different from the Johnson Cousins system in the PTF campaign have been reduced to this system as follows. For the P48 data, $R = R_{\text{Mould}} - 0.1$ was assumed. The P60 *griz* magnitudes were transformed to *BRI* using colours from simultaneous observations and normalizations as in Fukugita et al. (1996). Lacking spectroscopic information for SN 2010ah after maximum and based on the spectral similarity with SN 1998bw (see Fig. 1), the *g*, *r*, *i*, *z* and Mould *R* magnitudes of SN 2010ah have been converted to the *BVRI* system and *k*-corrected following Hamuy et al. (1993) and Oke & Sandage (1968), using as templates the spectra of SN 1998bw at phases comparable to those of SN 2010ah, after application of a time-folding factor of 1.3. The Ultraviolet/optical Telescope *U*-band filters were transformed to *U* magnitudes using the zero-points of Poole et al. (2008) and a λ^3 power law for the ultraviolet spectrum. Finally, synthetic magnitudes were extracted

from the Gemini and Keck spectra. As these were both taken with the slit positioned near the parallactic angle, we assumed that in both cases flux calibration is reasonable across the whole spectrum. When we measured the synthetic photometry we conservatively assigned an uncertainty of ± 0.3 mag to both points, as in Corsi et al. (2011). The error on the first two R -band points has been increased to 0.2 mag because the k -correction for those points is based on extrapolation from an SN 1998bw spectrum whose epoch is 3.5 d later in the time frame of SN 2010ah. At early epochs the spectral shape may change significantly over this time interval.

A fiducial time of explosion was selected. Since the explosion date is constrained by an upper limit on 2010 February 19, 4 d before the first PTF detection (Corsi et al. 2011, fig. 2), we assumed that the explosion occurred 2 d before first detection.

In Fig. 2, all magnitudes have been dereddened with $E(B - V) = 0.012$ (Schlegel, Finkbeiner & Davis 1998), adopting $R_V = 3.1$ and the Galactic extinction curve of Cardelli, Clayton & Mathis (1989). The comparison of the rest-frame LCs of SN 2010ah with the much better sampled ones of SN 1998bw in the same filters, and especially in the R band, where SN 2010ah has been most intensively observed, confirms that the time evolution of SN 2010ah is faster than that of SN 1998bw by a factor of ~ 1.3 . If the rest-frame monochromatic LCs of SN 1998bw are divided by this factor they match those of SN 2010ah reasonably well (Fig. 2), although bluer bands in SN 2010ah seem to evolve slightly faster than redder bands with respect to SN 1998bw. In the U band, the two points available for SN 2010ah are below the SN 1998bw template, suggesting a more rapid decay in this band, and perhaps some modest local absorption.

4 SN 2010AH: THE BOLOMETRIC LIGHT CURVE

In order to build a successful model for SN 2010ah it is necessary to construct a bolometric LC. This is essential for the estimate of the mass of ^{56}Ni synthesized in the explosion and as an input to our spectrum synthesis code. Since PTF photometry was obtained almost exclusively in the r band, it is difficult to construct a well-sampled bolometric LC. At the same time it would be dangerous to use r as a proxy for bolometric because the bolometric correction from r may be different from SN to SN.

In order to compute the bolometric LC of SN 2010ah we made use of the spectroscopic similarity between SN 2010ah and SN 1998bw and of the similar temporal evolution after rescaling in time by a factor of 1.3^{-1} (Fig. 2, see also Deng et al. 2005). We therefore adopted for SN 2010ah the bolometric correction from r obtained for SN 1998bw (Patat et al. 2001) at corresponding epochs (stretched in time by 1.3^{-1}). The resulting bolometric LC is shown in Fig. 3. The epoch and brightness of the peak are not very well determined, but the good coverage of the decline phase allows a model to be developed.

As a verification of this method, for the two epochs when spectra are available we integrated the flux under the spectra, added 20 per cent to account for flux outside the optical range taking SN 1998bw as an example (e.g. Melandri et al. 2012), and allowed for a conservatively large error to account for uncertain flux calibration of the spectra.

Also, again as a verification, a bolometric magnitude was computed directly from the monochromatic data when sufficient coverage of the spectrum was available. This includes the epochs of the *Subaru* photometry, when quasi-simultaneous *WISE* I -band data are also available (Corsi et al. 2011), and an epoch near 50 d, when b , r and i photometry is available. A correction of 20 per cent was




Figure 3. The PTF r -band LC of SN 2010ah (red) and the bolometric LC built from the photometry and correction with respect to SN 1998bw (black). The larger black points indicate estimates obtained directly from the spectra (near maximum) and from multicolour photometry.

again added for flux outside the optical bands. These points are highlighted in Figs 3 and 4.

4.1 SN properties derived from the light curve

A useful first step towards establishing a model for SN 2010ah is to derive basic information from the LC and the spectra. In the spirit of Arnett (1982) this requires measuring the width of the LC and a characteristic ejecta velocity from the spectra. These in turn can be transformed into first guesses for the ejected mass and kinetic energy. In Fig. 4, we compare the bolometric LC of SNe 2010ah, 1998bw and 2002ap, all of which look similar spectroscopically. The LC of SN 2010ah is very similar in shape to that of SN 2002ap, showing a fast rise to peak, which is reached within ~ 10 d. The LC of SN 2010ah is ~ 5 per cent broader than that of SN 2002ap, but significantly narrower (~ 25 per cent) than that of SN 1998bw. The characteristic velocity (at maximum) of SN 2010ah can be derived from fig. 5 of Corsi et al. (2011).

In order to obtain a rough estimate of the properties of SN 2010ah we can apply the well-known relations between ejected mass and velocity on the one hand and kinetic energy and LC shape on the other (Arnett 1982):

$$E_{\text{kin}} \propto M_{\text{ej}} v^2 \quad (1)$$

$$\tau_{\text{LC}} \propto \frac{\kappa^{1/2} M_{\text{ej}}^{3/4}}{E_{\text{kin}}^{1/4}}, \quad (2)$$

where E_{kin} is the kinetic energy of the ejecta, M_{ej} the ejected mass, κ the optical opacity, τ_{LC} the characteristic width of the LC. Adopting the commonly made assumption that κ is constant in all SNe Ib/c, simple scaling relations can be obtained:

$$\frac{M_1}{M_2} \propto \left(\frac{\tau_1}{\tau_2} \right)^2 \frac{v_1}{v_2} \quad (3)$$




Figure 4. Comparison of bolometric LCs of different SNe Ic: the filled dots show the bolometric LC of SN 2010ah, 1998bw and 2002ap. The empty green circles show the LC of SN 2010ah shifted upwards by 0.9 mag. This is a good match to SN 1998bw in the declining phase. The green crosses show the LC of SN 2010ah shifted upwards by 0.5 mag and stretched in time by a factor of 1.3. This is a good match to SN 1998bw except the earliest epochs. The empty blue dots show the LC of SN 2002ap shifted upwards by 1.1 mag. This is a reasonable match to SN 2010ah.

$$\frac{E_1}{E_2} \propto \left(\frac{\tau_1}{\tau_2}\right)^2 \left(\frac{v_1}{v_2}\right)^3. \quad (4)$$

The results that are obtained from these relations depend sensitively on the choice of v . This is a characteristic ejecta velocity, and in itself it is not a very well defined quantity. Measuring line velocity and using that value has a number of shortcomings: (1) the velocity changes (decreases) with time; (2) the velocity at which lines form can be different (higher) than the momentary photosphere; (3) different lines may form at different velocities because of ionization or optical depth effects.

Adopting a velocity used for modelling [as for example was done for the SNe shown in fig. 5 of Corsi et al. (2011), with the exception of SN 2010ah itself] eliminates at least some of these problems. Still, a single velocity does not represent the actual velocity distribution of the ejected matter. The velocity changes rapidly with time, and LCs have different shapes.

Very different results can be obtained if velocities at the same SN physical age (i.e. days from explosion, which is also an uncertain quantity except for GRB/SNe or for cases where shock breakout is detected) or at a particular *phase* of the LC (e.g. maximum) are used, since different SNe peak at different physical times from explosion. Let us show some examples of this.

Since the epoch of maximum of the LC of SN 2010ah is not well defined, different epochs can be chosen. Additionally, only two spectra are available. We use the velocity derived from the second spectrum (2010 March 7), 18 000 km s⁻¹. Corsi et al. (2011) assigned to this spectrum an epoch of 11 d, based on the first detection. Corresponding velocities at this epoch can be obtained for both SN 1998bw (via interpolation) and SN 2002ap (from direct modelling). All values are shown in Table 1. The value of E_{kin}

Table 1. Derived properties of SN 2010ah (M_{ej} , E_{kin}) with different assumptions about the epoch and using different reference SNe, whose M_{ej} and E_{kin} are given as are their velocities at different reference epochs.

SN	τ (d)	Vel (11 d) (km s ⁻¹)	M_{ej} (M_{\odot})	E_{kin} (10^{51} erg)
1998bw	17	22 500	10	50
2002ap	12	12 000	2.5	4
2010ah	13	18 000	5	15 (using 98bw)
	13	18 000	4	15 (using 02ap)
SN	τ (d)	vel (13 d) (km s ⁻¹)	M_{ej} (M_{\odot})	E_{kin} (10^{51} erg)
1998bw	17	19 000	10	50
2002ap	12	10 000	2.5	4
2010ah	13	18 000	6	25 (using 98bw)
	13	18 000	5	25 (using 02ap)

changes somewhat if we start from SN 1998bw or from SN 2002ap, reflecting the uncertainty in LC shape and time of maximum for SN 2010ah. Alternatively, we can consider the non-detection limit 4 d earlier, and assign to the 7 March spectrum a fiducial epoch of 13 d, as we do in the spectroscopic analysis below. The velocities of SNe 1998bw and 2002ap change significantly, despite the small change in epoch, and so do the results obtained for M_{ej} and E_{kin} of SN 2010ah. The value of E_{kin} , in particular, is now much higher. Indicative results for SN 2010ah are $M_{\text{ej}} \sim 4\text{--}6 M_{\odot}$, $E_{\text{kin}} \sim (1.5\text{--}3) 10^{52}$ erg, with large uncertainties. Similarly, uncertain results would be obtained if we used the earlier spectrum as a reference for velocity. This shows that the rescaling method should be applied with caution, especially if the data coverage is not excellent. Furthermore, it is imperative that SNe are used that have very similar spectral properties in order to avoid carrying incorrect information (especially about E_{kin} as derived from velocities) from the SN used as template to the one which is being studied.

The other important quantity that can be derived by interpolation is the mass of ^{56}Ni . This can be done using as reference the peak of the LC, or alternatively the late phase. The use of different and independent methods makes the estimate of the $M(^{56}\text{Ni})$ more reliable than those of M_{ej} or E_{kin} . Fig. 4 shows the two different approaches. The temporal evolution of the bolometric LC of SN 2010ah looks rather similar to that of SN 2002ap. If the bolometric LC of SN 2002ap is shifted up by 1.1 mag it tracks that of SN 2010ah very well, except for the first point, which suggests that SN 2010ah had a broader LC. The early rise of the LC of SN 2010ah may be due to mixing out of some ^{56}Ni , possibly in an aspherical explosion like that of SN 1998bw (Maeda et al. 2002; Maeda, Mazzali & Nomoto 2006). A shift of 1.1 mag corresponds to a factor of 2.75 in luminosity. The mass of ^{56}Ni for SN 2002ap was determined from modelling as $0.07 M_{\odot}$ (Mazzali et al. 2002). This yields for SN 2010ah $M(^{56}\text{Ni}) = 0.19 M_{\odot}$, consistent with the value derived by Corsi et al. (2011).

If we use as reference SN 1998bw, on the other hand, we can follow two different approaches. The post-maximum LC of SN 2010ah can be superposed on that of SN 1998bw if it is scaled up by 0.9 mag. This is a factor of 2.3 in luminosity. Adopting for SN 1998bw a ^{56}Ni mass of $0.41 M_{\odot}$ (Nakamura et al. 2001; Mazzali et al. 2006b), a ^{56}Ni mass of $0.19 M_{\odot}$ is obtained for SN 2010ah, consistent with the value obtained when rescaling from SN 2002ap. In fact, differences in the relative times are negligible at advanced epochs. This is therefore the safest approach.

Alternatively, one can rescale the bolometric LC of SN 2010ah in flux and stretch it by a factor of 1.3 to generate a LC that looks like that of SN 1998bw. In this case, the rescaling factor is only 1.6 (0.5 mag), and the ^{56}Ni mass for SN 2010ah is $0.26 M_{\odot}$. This result is clearly inconsistent. The different time evolution of the two LCs must be taken into account: at early times the contribution of ^{56}Co can be ignored, and shifting the peak from 17 d (SN 1998bw) to 13 d (SN 2010ah) leads to a ^{56}Ni mass for SN 2010ah smaller by a factor of 0.63 with respect to the initial estimate. Hence, we recover $M(^{56}\text{Ni}) = 0.16 M_{\odot}$, which is reasonably consistent with the estimate based on the LC tail. We conclude that for SN 2010ah $M(^{56}\text{Ni}) = 0.18 \pm 0.02 M_{\odot}$. A departure from this value can be introduced by (a) any ^{56}Ni which is mixed to high velocities and is only visible in the rising phase; (b) some deeply hidden ^{56}Ni which at peak does not yet contribute to the luminosity and only affects the LC at very late times. This central ^{56}Ni is known to be present in a high density inner region in probably all HNe (Maeda et al. 2003). A signature of this deep ^{56}Ni may also be seen in SN 2010ah after about day 40, when the LC begins to flatten, although data unfortunately stop soon thereafter. Therefore, $M(^{56}\text{Ni}) = 0.20 M_{\odot}$ is probably a lower limit for the mass of ^{56}Ni in SN 2010ah.

5 SPECTRAL MODELLING

Given the description above, only accurate spectral (and LC) modelling can yield reliable information on the properties of SNe Ib/c. For SN 2010ah only two spectra are available, so a very accurate result should not be expected. However, since the two spectra are separated by several days, modelling may yield significant information.

We used our SN Monte Carlo spectrum synthesis code, which was first presented by Mazzali & Lucy (1993). The code computes radiation transport in an SN ejecta starting from an assumed luminosity emitted at a lower boundary as a blackbody. SNe ejecta are assumed to be in free expansion, so that $v = R/t$. Bound-free processes are ignored, but electron scattering and line absorption are considered. The latter is treated including the process of fluorescence, which is essential for the formation of Type I SN spectra (Lucy 1999; Mazzali 2000). Radiative equilibrium is enforced in the SN envelope by conserving photon packets. Level populations are computed adopting a modified nebular approximation, which successfully accounts for the deviation from Local Thermodynamic Equilibrium. The code has been successfully applied to a number of different SNe Ib/c (e.g. Mazzali et al. 2002), and it has proved a very useful tool to go beyond line identification: thanks to its physical treatment of opacities it allows different density and abundance profiles to be tested, so that various SN properties can be determined quantitatively.

Since our code uses a physical luminosity, flux-calibrated spectra are necessary for a meaningful comparison. The spectra are assumed to be reasonably well flux calibrated based on the observational techniques used to obtain them. We corrected them for a Milky Way extinction of $E(B - V) = 0.012$ (Schlegel et al. 1998) and measured R -band magnitudes using the `PYSYNPHOT`¹ package. In order to account for calibration errors we applied a large error of 0.3 mag for both measurements. The corrected observed magnitudes are $R = 17.95$ mag for the Gemini spectrum and $R = 18.15$ mag for the Keck spectrum.

Our code requires as input a density distribution. This is essential in order to define the properties of the SN. In particular, SNe Ic with progressively broader lines are characterized by increasing values of the parameter $E_{\text{kin}}/M_{\text{ej}}$. Low-energy, ‘normal’ SNe like SN 1994I have $E_{\text{kin}}/M_{\text{ej}} \sim 1$ (Sauer et al. 2006), while GRB/SNe like SN 1998bw, which show very broad lines, have $E_{\text{kin}}/M_{\text{ej}} > 2$ (Nakamura et al. 2001). Given the spectral similarity between SN 2010ah, SN 1998bw and SN 2002ap, we adopt the explosion model that was used for SN 1998bw, and rescaling it to match the parameters of SN 2010ah as estimated above.

We use a distance of 216 Mpc (i.e. a distance modulus $\mu = 36.67$ mag) assuming $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and standard cosmology, and a reddening $E(B - V) = 0.012$ (Corsi et al. 2011).

Given the estimated mass and energy of SN 2010ah, we started by rescaling the model for SN 1998bw (model CO138, $M_{\text{ej}} = 10 M_{\odot}$, $E_{\text{kin}} = 310^{52}$ erg; Iwamoto et al. 1998) by a factor between 0.3 and 0.5 in mass, which obviously implies a reduction in energy of the same magnitude.

Since both spectra are relatively early, our models can only probe the outer part of the ejecta. Therefore, while they can yield a reasonable estimate of the one-dimensional energy, which is mostly associated with the high-velocity outer ejecta, they cannot probe the bulk of the mass, which is located at lower velocities and deeper layers than the early spectra can reach. A more accurate estimate of the mass in this case requires modelling the LC. Nebular spectra would be very useful for this purpose (e.g. Mazzali et al. 2010), but none are available for SN 2010ah.

A fundamental element is the choice of rise time. The longer the rise time, the larger the ejected mass has to be in order to obtain a reasonable spectrum. SN 2010ah was first detected on 2010 February 23.5. The first spectrum was obtained on 2010 March 2, i.e. 7 d later. A deep non-detection limit 4 d prior to discovery means that the upper limit for the age of the first spectrum is 11 d (10.5 rest-frame days). Averaging these values, we assume $t = 9$ d as the epoch of the first spectrum. Considering the redshift ($z = 0.0498$), we adopted for the March 2 spectrum $t = 8.5$ d.

Our code computes a spectrum starting from the SN physical parameters L , t and v , in order to define a consistent temperature structure in the ejecta. The photospheric velocity v_{ph} can be inferred from the spectra. Corsi et al. (2011) measured an Si II 6355 Å absorption velocity $v = 29\,000 \text{ km s}^{-1}$ in the March 2 spectrum. We adopt for the model a slightly lower velocity, $25\,000 \text{ km s}^{-1}$, and obtain a reasonable solution for the spectrum.

The luminosity necessary to reproduce the flux in the March 2 spectrum is $5 \times 10^{42} \text{ erg s}^{-1}$, i.e. $\log(L/L_{\odot}) = 9.12$. Given the rather small epoch, we find that the best results are obtained if model CO138 is rescaled in mass by a factor of 0.3. This gives $M_{\text{ej}} = 3.2 M_{\odot}$; $E_{\text{kin}} = 10^{52}$ erg, but we should remark once more that the mass derived from the spectral fits alone can be highly uncertain since the spectra do not cover a large range of epochs. It appears however that a much smaller mass is located at high velocities than in the case of SN 1998bw, so we need to reduce the mass at $v > 30\,000 \text{ km s}^{-1}$. We did this by steepening the density slope, using a power law $\rho(r) \propto r^n$ with index $n \approx -7$ above $20\,000 \text{ km s}^{-1}$. We find $\approx 0.2 M_{\odot}$ at $v > 0.1c$, compared to $\approx 1 M_{\odot}$ for SN 1998bw. There is however practically no material at $v > 10^5 \text{ km s}^{-1}$, as was also the case for SN 1998bw. On 2010 March 2, only $\sim 0.5 M_{\odot}$ of material was found above v_{ph} . The equivalent model for SN 1998bw had $\sim 1.2 M_{\odot}$ located above $v_{\text{ph}} = 31\,600 \text{ km s}^{-1}$ on 1998 May 3 (Iwamoto et al. 1998), 8 d after explosion. Therefore, the velocity of SN 2010ah is actually slightly smaller than that of SN 1998bw.

¹ <http://stsdas.stsci.edu/pysynphot/>




Figure 5. The spectrum of SN 2010ah obtained on 2010 March 2 (black) and the corresponding synthetic model (blue).

The synthetic spectrum is compared to the observed one in Fig. 5. Main features are Fe II absorption at $\sim 4000, 4600$ Å, Si II near 6000 Å, the O I–Ca II blend near 7000 Å. The observed spectrum is reproduced adopting a composition above the photosphere dominated by oxygen (78 per cent by number), followed by Ne (20 per cent), intermediate-mass elements (5 per cent). Ca and Fe are only present at the level of ~ 0.5 per cent.

The next spectrum is only 5 d later, 2010 March 7. This corresponds to 14 d, (13.3 d in the rest frame). The velocity in this spectrum has dropped significantly (Corsi et al. 2011, measured $18\,000$ km s $^{-1}$ in the Si II line), which is compatible with the drop observed in both GRB/HNe and non-GRB/HNe. We can thus probe deeper layers of the ejecta. The luminosity required to match the flux is $L = 4 \times 10^{42}$ erg s $^{-1}$, i.e. $\log(L/L_{\odot}) = 9.01$. A velocity of $v_{\text{ph}} = 13\,500$ km s $^{-1}$ gives the best fit to the spectrum using the same density distribution as at the earlier epoch rescaled to the later epoch (Fig. 6). The photospheric velocity drops by much more than the observed Si II velocity, suggesting that the observed feature is affected by other contributions. On the other hand, our synthetic Si II line has a somewhat lower velocity than the observed one. At this epoch, the mass above the photosphere is $2.25 M_{\odot}$. The composition is similar to that used for the previous epoch, with a little less oxygen (60 per cent), replaced by neon (30 per cent). The spectrum is still formed inside the oxygen layer. Abundances do not indicate enrichment of intermediate-mass elements or mixing out of Fe. The synthetic O I line is too deep, but it is blended with Ca II IR, which actually dominates in strength near 7500 Å. A lower density may lead to a better O I line, but the Si II line would look worse. Since we only have these two spectra, we refrain from performing a more detailed but uncertain analysis. We rather turn to the LC in order to determine parameters such as M_{ej} , ^{56}Ni mass and distribution.

6 LIGHT CURVE MODELS

It is well known that LC models alone do not have the ability to determine uniquely all SN parameters. In particular, while the mass




Figure 6. The spectrum of SN 2010ah obtained on 2010 March 7 (black) and the corresponding synthetic model (blue).

of ^{56}Ni can be reasonably well established from the LC peak if the time of explosion is known to within a good accuracy, or from the late, exponential decline phase of the LC if this is observed, the two parameters that determine the shape of the LC, M_{ej} and E_{kin} are degenerate (equation 2; Arnett 1982). Only the simultaneous use of LC and spectra can break this degeneracy (Mazzali et al. 2000). Having established the most likely values of M_{ej} and E_{kin} through spectral modelling, we can test whether they can reproduce the LC (Mazzali et al. 2002). Fitting the LC yields additional information about the mass and distribution of ^{56}Ni , in particular at velocities lower than the photosphere sampled by the spectra. This is particularly useful for SN 2010ah, since for this SN we do not have nebular spectra, which would make it possible to obtain an accurate description of the inner ejecta (e.g. Mazzali et al. 2010).

The properties of the LC of an SN I (i.e. H-free) are determined by the mass of ^{56}Ni , which determines the overall luminosity, and by the mass, density and composition of the ejecta. Mass and energy have a direct impact on the density of the ejecta as a function of time, and hence on the diffusion time of the photons. The abundances in the ejecta are important in that they determine the opacity (in a Type I SN this is dominated by line opacity; Pauldrach et al. 1996). If both early and late-time spectra are available, all these quantities can be independently determined through spectral modelling. Typically, only small modifications to the mass and distribution of ^{56}Ni are required to reproduce the LC and validate the results (Mazzali et al. 2009). This is however not the case for SN 2010ah, since no nebular spectra are available. Here, we must rely on LC modelling alone to establish the mass and density of the inner layers. Still, the properties of the outer layers are known from the early-time spectra, and so E_{kin} is reasonably well established.

We use the one-dimensional, monochromatic LC code described in Cappellaro et al. (1997) and later modifications. The code is based on the Monte Carlo method: given a density structure of the ejecta and a ^{56}Ni distribution, the emission and deposition of gamma-rays and positrons produced in the radioactive decay of ^{56}Ni to ^{56}Co and hence to ^{56}Fe are computed. Deposition is assumed to be




Figure 7. The *uvoir* LC of SN 2010ah and two models, one obtained using the rescaled explosion model used to fit the spectra (blue) and one where corrections have been applied for mixed out ^{56}Ni and a dense inner core (black). The bigger blue points are those obtained from direct integration of the observed magnitudes from the spectra (near peak), from *Subaru* photometry (~ 25 d) and at one late epoch (2010 April 14, 49 rest-frame days) when multiple photometry was available.

instantaneous, and to be followed by the emission of optical photons. Photon diffusion, which determines the LC, is also followed with a Monte Carlo scheme. Optical opacity is treated assuming that line opacity dominates: accordingly, it is assumed to depend on composition, as in Mazzali & Podsiadlowski (2006), reflecting the different number of active spectral lines in different atomic species.

Initially, we use as input the density/abundance profiles derived from early-time spectral modelling. In particular, the model we used has $M_{\text{ej}} = 3.0 M_{\odot}$, $E_{\text{kin}} = 1.15 \times 10^{52}$ erg. The result is the synthetic LC shown in blue in Fig. 7. This reproduces the maximum using a ^{56}Ni mass of $0.19 M_{\odot}$, but the rise is delayed relative to the observations. The decline is also well reproduced, but only until day 30 or so. Neither of these problems is new in the context of SN Ic LC modelling: the slow rise of the synthetic LC is typically the result of insufficient mixing out of ^{56}Ni , while the slow observed late-time decline is actually common to all SNe Ic, and a solution was proposed that involves a dense low-velocity core, possibly a signature of an aspherical explosion (Maeda et al. 2003).

We therefore modified both the inner density and the outer ^{56}Ni abundance, and obtained the LC shown in black in Fig. 7. This modified model has a power-law index $n \approx -1$ below $20\,000 \text{ km s}^{-1}$. It also has a slightly larger mass: $3.3 M_{\odot}$. The additional $0.2 M_{\odot}$ is located at $v < 10\,000 \text{ km s}^{-1}$. The ^{56}Ni distribution is also enhanced: outer layers ($v > 20\,000 \text{ km s}^{-1}$) contain a ^{56}Ni mass fraction of ~ 10 per cent. This is much larger than the spectral modelling results, which indicate ~ 1 per cent. The ^{56}Ni mass is now $0.25 M_{\odot}$ (instead of $0.19 M_{\odot}$), but because of the location of the additional ^{56}Ni there is no effect on the peak of the LC, only on the rising phase. The highest ^{56}Ni abundance is at the lowest velocities, $v < 3000 \text{ km s}^{-1}$, where a mass fraction of more than 50 per cent is reached. However, most of the ^{56}Ni is located at intermediate velocities,

with a mass fraction of ~ 15 per cent at $5000\text{--}10\,000 \text{ km s}^{-1}$ and ~ 1 per cent at $10\,000\text{--}20\,000 \text{ km s}^{-1}$. The kinetic energy, on the other hand, is only marginally increased (now 1.19×10^{52} erg instead of 1.15×10^{52} erg). The modified model gives a better match to the overall LC of SN 2010ah. Only the last phase ($t > 60$ d) shows a discrepancy. A further increase in mass at very low velocities may be a solution, but we refrain from attempting such models because of the lack of later points to confirm the slower decline. Additionally, although galaxy subtraction has been applied homogeneously to all images (Corsi et al. 2011), since the galaxy brightness in *r* is comparable to that of SN 2010ah at about 30 d after explosion, the presence of some residual at epochs $\gtrsim 60$ d cannot be excluded. This may explain the excess seen in Fig. 7.

In the modified models, ^{56}Ni is added mostly at high velocity ($> 20\,000 \text{ km s}^{-1}$). This requires reaching a ^{56}Ni mass fraction of ~ 0.1 in the outer ejecta. Although the ^{56}Ni distribution we have used yields a reasonably good match to the peak of the LC of SN 2010ah, it is inconsistent with the abundance of ^{56}Ni and decay products required for the spectra. This discrepancy is potentially interesting. It may indicate that some ^{56}Ni was ejected at high velocity but not along our line of sight, so that it caused the rapid rise of the LC (Maeda et al. 2006) without leaving an imprint on the spectra in the form of absorption lines.

Although this suggestion depends mostly on the first LC point, this interpretation is appealing in view of the properties of highly energetic SNe Ic. We know from nebular spectroscopy that the distribution of elements is not spherically symmetric, in the sense that some ^{56}Ni and Fe are found at high velocities, probably in a funnel-like distribution which may track the position of the GRB jet in some cases (Mazzali et al. 2001), while oxygen and other unburned elements are ejected more in a plane, likely the equator of the progenitor. Two-dimensional models suggest this (Maeda et al. 2002). There is further evidence for asphericity in SNe Ib/c. Nebular studies suggest that even lower energy SNe can be aspherical, albeit to a lesser degree (Maeda et al. 2008; Modjaz et al. 2008; Taubenberger et al. 2009; Maurer et al. 2010). It is also particularly interesting that in an SN Ib linked to an X-ray Flash (SN 2008D; Soderberg et al. 2008) spectral analysis suggests a high kinetic energy and the presence of a jet which was quenched by the helium shell (Mazzali et al. 2008), and later data confirmed asphericity in the oxygen distribution (Tanaka et al. 2008).

The kinetic energy derived for SN 2010ah is larger than that of SN 2008D, although smaller than in GRB/SNe, and the ejected mass is compatible with that of SN 2008D if the He shell in the latter is not considered. An off-axis, jet-driven SN was proposed for SN 2008D (Mazzali et al. 2008). More energetic SNe Ic are all linked to GRBs and aspherical (e.g. Mazzali et al. 2001). Given the peculiarly high first point in the LC of SN 2010ah, an aspherical distribution of elements may also be speculated for SN 2010ah. The lack of nebular or spectropolarimetric data prevents us from testing this hypothesis.

7 DISCUSSION

The results of the analysis we performed can be used to estimate the properties of the progenitor of SN 2010ah. This is interesting, because of the suggested relation between E_{kin} , $M(^{56}\text{Ni})$ and M_{ej} , which should be a good proxy for the mass of the progenitor star (Mazzali et al. 2009). In the case of SN 2010ah, we found $M_{\text{ej}} \sim 3.3 M_{\odot}$, $E_{\text{kin}} \sim 1.2 \times 10^{52}$ erg (in spherical symmetry) and $M(^{56}\text{Ni}) \sim 0.25 M_{\odot}$.




Figure 8. The mass of ^{56}Ni and the E_{kin} as derived from spectral/LC modelling of a number of SNe Ib/c. Colour coding is by ejected mass.




Figure 9. The ejecta mass and the E_{kin} derived from spectral/LC modelling of a number of SNe Ib/c and some SNe IIP. Colour coding is by spectral type.

If we plot the properties of SN 2010ah and those of other well-studied SNe (Figs 8 and 9), we see a trend between M_{ej} , $M(^{56}\text{Ni})$ and E_{kin} . Fig. 8 shows $M(^{56}\text{Ni})$ versus E_{kin} . The trend is clear, although SN 2006aj deviates somewhat. In this case, the birth of a magnetar was suggested (Mazzali et al. 2006a), and the energy released may have led to an increased synthesis of ^{56}Ni . Slightly more massive progenitors probably collapsed to a black hole, thus, reducing the amount of material available for nucleosynthesis. At larger masses (and larger energies) the efficiency of production and ejection of ^{56}Ni increases again. Fig. 9 shows $M(^{56}\text{Ni})$ versus E_{kin} . Here, the




Figure 10. The mass of ^{56}Ni derived from spectral/LC modelling of a number of SNe Ib/c plotted against the inferred zero-age main sequence (ZAMS) mass of the progenitor star using single-star evolutionary models. Colour coding is by spectral type, and it is the same as in Fig. 9.




Figure 11. The E_{kin} derived from spectral/LC modelling of a number of SNe Ib/c plotted against the inferred ZAMS mass of the progenitor star using single-star evolutionary models. Colour coding is by spectral type, and it is the same as in Fig. 9.

relation is less well defined, but a trend is still visible. SNe Ib and Iib have a massive He envelope and lie slightly higher in the plot, suggesting that the E_{kin} is not influenced by the properties of the outer layers, as is reasonable to expect.

We can try to go from the estimated M_{ej} to a progenitor mass. Using the Nomoto & Hashimoto (1988) models, and assuming that the mass ejected by SN 2010ah corresponds to the CO core of a massive star minus the remnant mass, we end up with a black hole

remnant, a CO core of $\sim 5\text{--}6 M_{\odot}$ and a $M_{ZAMS} \sim 24\text{--}28 M_{\odot}$. In the more traditional plots of $M(^{56}\text{Ni})$ or E_{kin} versus M_{ZAMS} (Figs 10 and 11), SN 2010ah falls nicely on the relation that is taking shape (e.g. Mazzali et al. 2009). This relation is derived for single-star evolution, and using a single, albeit old set of stellar models. The process of stripping the outer H and He layers may require interaction with a companion in a binary system (Fryer et al. 2007) and may not be easy to parametrize. Also, the uncertainty on the nature of the remnant (a neutron star or a black hole?) may introduce errors in the mass estimate.

Still, in the case of SN 2010ah, it looks as if all properties we derived (M_{ej} , E_{kin} , $M(^{56}\text{Ni})$) are consistently smaller than those of GRB/SNe, although they are large compared to other non-GRB/HNe. We may be testing the limits for stars to produce GRBs. Another recent non-GRB/HN, SN 2010ay (Sanders et al. 2012) may come even closer to this limit and it will be an interesting case study.

A question remains concerning SN 2010ah: did it produce a GRB, or was it at least driven by a jet which failed to escape from the star, as was suggested for SN 2008D (Mazzali et al. 2008)? No detection of a GRB has been reported. Off-axis GRBs may be detected in the radio, but this is not the case for SN 2010ah (Corsi et al. 2011). The anomalously rapid rise of the LC may suggest the presence of highly processed material in a direction not along our line of sight, but this cannot be regarded as a strong argument in the absence of other supporting evidence. Spectropolarimetry in the early phase or late-time spectroscopy would be useful in order to reveal asphericities. These may in turn point to the presence of a jet which, as in the case of SN 2008D (Tanaka et al. 2008), may have aided exploding the star even though it may not have emerged from it. Unfortunately, no such data are available for SN 2010ah, so we cannot determine the degree of asphericity or the presence of any asymmetry. A suggestion for asphericity drawn from the rapid rise of the LC may be a point in favour of this. It would be important that exceptional SNe like SN 2010ah are studied in more detail in the future.

ACKNOWLEDGEMENTS

We acknowledge financial support from grants: INAF PRIN 2009 and 2011, ASI I/016/07/0, ASI I/088/06/0.

REFERENCES

- Arcavi I. et al., 2011, *ApJ*, 742, L18
 Arnett W. D., 1982, *ApJ*, 253, 785
 Bromberg O., Nakar E., Piran T., Sari R., 2011, *ApJ*, 740, 100
 Bufano M. et al., 2012, *ApJ*, 753, 67
 Cappellaro E., Mazzali P. A., Benetti S., Danziger I. J., Turatto M., della Valle M., Patat F., 1997, *A&A*, 328, 203
 Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
 Chornock R. et al., 2013, preprint (arXiv:1302.0009)
 Corsi A. et al., 2011, *ApJ*, 741, 76
 Crockett R. M. et al., 2008, *MNRAS*, 391, L5
 Deng J., Tominaga N., Mazzali P. A., Maeda K., Nomoto K., 2005, *ApJ*, 624, 898
 Filippenko A. V., 1997, *ARA&A*, 35, 309
 Filippenko A. V., Matheson T., Ho L. C., 1993, *ApJ*, 415, L103
 Foley R. J. et al., 2003, *PASP*, 115, 1220
 Fryer C. L. et al., 2007, *PASP*, 119, 1211
 Fukugita M., Ichikawa T., Gunn J. E., Doi M., Shimasaku K., Schneider D. P., 1996, *AJ*, 111, 1748
 Gal-Yam A., 2012, *Sci*, 337, 927
 Gal-Yam A., Ofek E. O., Shemmer O., 2002, *MNRAS*, 332, L73
 Gal-Yam A. et al., 2009, *Nat*, 462, 624
 Galama T. J. et al., 1998, *Nat*, 395, 670
 Gorosabel J. et al., 2006, *A&A*, 459, L33
 Gorosabel J. et al., 2010, *A&A*, 522, A14
 Hachinger S., Mazzali P. A., Taubenberger S., Hillebrandt W., Nomoto K., Sauer D. N., 2012, *MNRAS*, 422, 70
 Hamuy M., Phillips M. M., Wells L. A., Maza J., 1993, *PASP*, 105, 787
 Hamuy M. et al., 2009, *ApJ*, 703, 1612
 Hjorth J. et al., 2003, *Nat*, 423, 847
 Iwamoto K., Nomoto K., Hoflich P., Yamaoka H., Kumagai S., Shigeyama T., 1994, *ApJ*, 437, L115
 Iwamoto K. et al., 1998, *Nat*, 395, 672
 Jordi K., Grebel E. K., Ammon K., 2006, *A&A*, 460, 339
 Kashikawa N. et al., 2002, *PASJ*, 54, 819
 Kasliwal M. et al., 2012, *ApJ*, 755, 161
 Kawabata K. S. et al., 2002, *ApJ*, 580, L39
 Kawabata K. S. et al., 2003, *ApJ*, 593, L19
 Law N. M. et al., 2009, *PASP*, 121, 1395
 Lucy L. B., 1999, *A&A*, 345, 211
 MacFadyen A. I., Woosley S. E., 1999, *ApJ*, 524, 262
 Maeda K., Nakamura T., Nomoto K., Mazzali P. A., Patat F., Hachisu I., 2002, *ApJ*, 565, 405
 Maeda K., Mazzali P. A., Deng J., Nomoto K., Yoshii Y., Tomita H., Kobayashi Y., 2003, *ApJ*, 593, 931
 Maeda K., Mazzali P. A., Nomoto K., 2006, *ApJ*, 645, 1331
 Maeda K. et al., 2008, *Sci*, 319, 1220
 Malesani D. et al., 2004, *ApJ*, 609, L5
 Matheson T. et al., 2003, *ApJ*, 599, 394
 Maund J. R., Smartt S. J., Kudritzki R. P., Podsiadlowski P., Gilmore G. F., 2004, *Nat*, 427, 129
 Maund J. et al., 2011, *ApJ*, 739, L37
 Maurer J. I. et al., 2010, *MNRAS*, 402, 161
 Mazzali P. A., 2000, *A&A*, 363, 705
 Mazzali P. A., Lucy L. B., 1993, *A&A*, 279, 447
 Mazzali P. A., Podsiadlowski P., 2006, *MNRAS*, 369, L19
 Mazzali P. A., Iwamoto K., Nomoto K., 2000, *ApJ*, 545, 407
 Mazzali P. A., Nomoto K., Patat F., Maeda K., 2001, *ApJ*, 547, 988
 Mazzali P. A. et al., 2002, *ApJ*, 572, L61
 Mazzali P. A. et al., 2003, *ApJ*, 599, L95
 Mazzali P. A., Deng J., Maeda K., Nomoto K., Filippenko A. V., Matheson T., 2004, *ApJ*, 614, 858
 Mazzali P. A. et al., 2005, *Sci*, 308, 1284
 Mazzali P. A. et al., 2006a, *Nat*, 442, 1018
 Mazzali P. A. et al., 2006b, *ApJ*, 645, 1323
 Mazzali P. A. et al., 2007, *ApJ*, 670, 592
 Mazzali P. A. et al., 2008, *Sci*, 321, 1185
 Mazzali P. A., Deng J., Hamuy M., Nomoto K., 2009, *ApJ*, 703, 1624
 Mazzali P. A., Maurer I., Valenti S., Kotak R., Hunter D., 2010, *MNRAS*, 408, 87
 Melandri A., Pian et al., 2012, *A&A*, 547, A82
 Modjaz M., Kirshner R. P., Blondin S., Challis P., Matheson T., 2008, *ApJ*, 687, L9
 Nakamura T., Mazzali P. A., Nomoto K., Iwamoto K., 2001, *ApJ*, 550, 991
 Nomoto K., Hashimoto M., 1988, *Phys. Rep.*, 163, 13
 Nomoto K., Iwamoto K., Suzuki T., 1995, *Phys. Rep.*, 256, 173
 Oke J. B., Sandage A., 1968, *ApJ*, 154, 21
 Patat F. et al., 2001, *ApJ*, 555, 900
 Pauldrach A. W. A., Duschinger M., Mazzali P. A., Puls J., Lennon M., Miller D. L., 1996, *A&A*, 312, 525
 Perets H. B. et al., 2010, *Nat*, 465, 322
 Pian E. et al., 2006, *Nat*, 442, 1011
 Pignata G. et al., 2011, *ApJ*, 728, 14
 Poole T. S. et al., 2008, *MNRAS*, 383, 627
 Quimby R. M., Aldering G., Wheeler J. C., Höflich P., Akerlof C. W., Rykoff E. S., 2007, *ApJ*, 668, L99
 Quimby R. M. et al., 2011, *Nat*, 474, 487
 Rau A. et al., 2009, *PASP*, 121, 1334
 Richmond M. W. et al., 1996, *AJ*, 111, 327

Sanders N. E. et al., 2012, ApJ, 756, 184
Sauer D. N., Mazzali P. A., Deng J., Valenti S., Nomoto K., Filippenko A.
V., 2006, MNRAS, 369, 1939
Schlegel E., Finkbeiner D. P., Davis M., 1998, ApJ, 500, 525
Soderberg A. M., Nakar E., Berger E., Kulkarni S. R., 2006, ApJ, 638, 930
Soderberg A. M. et al., 2008, Nat, 453, 469
Soderberg A. M. et al., 2010, Nat, 463, 513
Stanek K. Z. et al., 2003, ApJ, 591, L17

Tanaka M. et al., 2008, ApJ, 692, 1131
Tanaka M. et al., 2012, ApJ, 754, 63
Taubenberger S. et al., 2009, MNRAS, 397, 677
van der Horst A. J. et al., 2011, ApJ, 726, 99
Van Dyk S. et al., 2011, ApJ, 741, L28
Woosley S. E., 1993, ApJ, 405, 273

This paper has been typeset from a \TeX/L\AA\TeX file prepared by the author.