

Electron-Induced Conversion of Silylones to Six-Membered Cyclic Silylenes

Sudipta Roy,[†] Kartik Chandra Mondal,[†] Lennard Krause,[†] Peter Stollberg,[†] Regine Herbst-Irmer,[†] Dietmar Stalke,^{*,†} Jann Meyer,[†] A. Claudia Stückl,[†] Bholanath Maity,[‡] Debasis Koley,^{*,‡} Suresh Kumar Vasa,[§] Sheng Qi Xiang,[§] Rasmus Linser,[§] and Herbert W. Roesky^{*,†}

[†]Institut für Anorganische Chemie, Georg-August-Universität, Tammannstraße 4, 37077 Göttingen, Germany

[‡]Department of Chemical Sciences, IISER-Kolkata, Mohanpur Campus, Mohanpur 741252, India

[§]Abteilung für NMR-basierte Strukturbiologie, Max-Planck-Institut für Biophysikalische Chemie, Am Faßberg 11, 37077 Göttingen, Germany


S Supporting Information

ABSTRACT: A silicon atom in the zero oxidation state stabilized by two carbene ligands is known as siladicalcarbene (silylone). There are two pairs of electrons on the silicon atom in silylone. This was recently confirmed by both experimental and theoretical charge density investigations. The silylone is stable up to 195 °C in an inert atmosphere. However, a substoichiometric amount (33 mol%) of potassium metal triggers the activation of the unsaturated C:Si:C backbone, leading to a selective reaction with a tertiary C–H bond in an atom-economical approach to form a six-membered cyclic silylene with three-coordinate silicon atom. Cyclic voltammetry shows that this reaction proceeds via the formation of a silylone radical anion intermediate, which is further confirmed by EPR spectroscopy.

In 1994, West et al.¹ reported on the first N-heterocyclic silylene ($R_2Si:$) which is stable at room temperature (Scheme 1a). After this breakthrough, the chemistry of low-valent silicon was explored more rapidly by various groups.² A silylene in general is a molecule that contains a divalent neutral silicon atom having a lone pair of electrons at the silicon atom.¹ Silylenes with three-coordinate silicon are also known^{2a,c,d,e} (Scheme 1c–f). Some of these compounds can activate small molecules.^{3,4}

In 2009, theoretical calculations on various $(NHC)_2Si$ compounds, coined as silylones, showed that the synthesis of this class of compounds is experimentally feasible.⁵ Approximately two decades after the first report¹ of silylene ($R_2Si:$), the syntheses of silylones $(cAAC)_2Si$ with two-coordinate silicon were reported ($cAAC$ = cyclic alkyl(amino) carbene, Scheme 1g).⁶ Later on, a cyclic silylone $(bNHC)Si$ (Scheme 1h) was prepared by Driess et al.⁷ Silylone features a silicon atom in the oxidation state zero which is stabilized by two carbene ligands. Consequently, this silicon atom contains two pairs of electrons. The silylone (or siladicalcarbene) is the heavier analogue of the carbodicalcarbene (carbene)₂C.^{8–10} Similarly, it also possesses two lone pairs of electrons at the central carbon atom, which are shown to form coordination bonds with acceptor molecules. So far, only metal complexes of carbo-

Scheme 1. (a–f) Representative Silylenes with Two- and Three-Coordinate Silicon, (g) Acyclic Silylones Stabilized by cAAC, and (h) Cyclic Silylones Stabilized by bNHC




dicarbene are developed in connection with their reactivity.^{8–11} In contrast, on the reactivity of silylones, nothing has been known until now.

We have recently reported the synthesis of the dark blue-colored silylone $(cAAC)_2Si$, prepared from the diradical $(cAAC)_2SiCl_2$ under KC_8 reduction,^{12a} and the related charge density investigations.^{12b} The silicon atom of a silylone possesses one non-bonding lone pair, and the second pair of electrons participates in a three-center, two-electron π -bond between two carbene carbon atoms and the central silicon atom (C:Si:C).^{6a} Thus, Si–C bonds of silylones are very strong. This results in an unforeseen stability of these molecules. We have observed that crystalline silylones are stable in an inert atmosphere for about 2 years without any decomposition. They can be characterized by mass spectrometry (see Supporting Information (SI)). Silylones (**1**) are thermally stable up to 195 °C, at which they melt to produce a blue liquid and decompose to a yellow liquid above 220 °C. Silylenes with a two-coordinate silicon atom can react with molecular hydrogen, ammonia, and carbon dioxide.^{3,4} In contrast, $(cAAC)_2Si$ does not react with any of these gases at room temperature. Herein we report the first reactivity study of silylones generating cyclic silylenes with a three-coordinate silicon atom.

Received: October 10, 2014

Considering the fact that cAAC is both a strong σ -donor and a π -acceptor, (cAAC)₂Si might accept an electron at the unsaturated C:Si:C backbone, (Cy-cAAC)₂Si (**1a**, Cy-cAAC = :C(CH₂)(CMe₂)(C₆H₁₀)N-2,6-*i*Pr₂C₆H₃) was investigated by cyclic voltammetry (CV) in 0.1 M [*n*-Bu₄N]PF₆ THF solution. Fortunately, the CV showed a quasi-reversible reduction at $E_{1/2} = -1.55$ V (referred against Cp*₂Fe/Cp*₂Fe⁺), indicating the formation of a radical anion **1a**^{•-}. In addition, the measurement showed three irreversible reductions ($E = -1.07$, -2.39 , and -2.72 V, see SI). Based on the quasi-reversible nature of the CV we anticipated a possible rearrangement of the intermediate radical anion and accordingly a chemical reaction was set up. THF was added at room temperature to a 1:1 molar solid-state mixture of (Cy-cAAC)₂Si (**1a**) and metallic potassium via a cannula. The resultant dark blue solution was stirred for 20 min to obtain a dark green solution which further turned to greenish-yellow solution within the next 15 min of additional stirring. The solution was filtered to separate the unreacted potassium (67 mol%), indicating 33 mol% of potassium was consumed during the reaction. The THF solution was concentrated and stored at -32 °C in a freezer to form bright yellow blocks of **2a** in 80% yield. The analogous compound **2b** was prepared from the precursor (Me₂-cAAC)₂Si (**1b**) (see SI). The X-ray single-crystal diffractions revealed that **2a,b** are isomers of the silylones **1a,b**, respectively. The transformation of **1a,b** into **2a,b** can be rationalized as one of the H-CMe₂ groups (**1a,b**) has selectively reacted (Scheme 2) with one C_{cAAC}-Si bond, producing H-C_{cAAC}-Si(CMe₂) moiety in products **2a,b**, respectively.

Scheme 2. Conversion of Silylone with Two-Coordinate Silicon to Silylene with Three-Coordinate Silicon



The solutions of **2a,b** are stable at 0 °C for about 2 months. The crystals of **2a,b** are stable at room temperature for more than 1 month in an inert atmosphere. They (**2a,b**) melt in the temperature range of 223–224 °C and 211–212 °C, respectively, while the melting range of **1a,b** is 194–195 °C. Compounds **2a,b** were also characterized (see SI) by mass spectrometry (m/z (100%): 678.4(**2a**), 598.4(**2b**)). The ²⁹Si NMR spectra of **2a,b** exhibit resonances at 55.98 and 54.55 ppm, respectively, which are upfield-shifted when compared to those of **1a,b** (71.15 and 66.71 ppm). The ¹³C NMR spectra of **2a,b** show resonances at 69.4(CH)/173.5(C:) and 66.2(CH)/172.1(C:) ppm, respectively, which are upfield-shifted when compared to those of **1a,b** (210.8 and 210.9 ppm).^{6a,b} The absence of ¹H nuclei directly bonded to Si of **2a** is confirmed by a slow solid-state NMR cross-polarization buildup identical to **1a**. A reduced CSA, however, confirms the more symmetrical distribution of electron density around the Si in **2a** than in **1a** (see SI). The unsymmetrical electronic environment of the ²⁹Si center in **1a** resulting from the lone pair creates a strong chemical shift anisotropy (CSA, see SI). The UV-vis spectra of yellow-colored **2a,b**, recorded in *n*-hexane, show absorption bands at 423/420 nm (see SI), while dark blue-colored

silylones **1a,b** exhibit bands at 575/570 and 618/611 nm, respectively.^{6a,b}

2a,b were both characterized by X-ray single-crystal determinations. The structural aspects of **2a** are described herein (see SI for **2b**). Compound **2a** crystallizes in the space group $P2_1/n$. The silicon atom adopts a distorted trigonal pyramidal geometry which is bound to three carbon atoms including the central carbon atom of the former *i*Pr substituent. The selected bond lengths and angles are given in the caption of Figure 1. The structure of **2a** shows that one carbene carbon




Figure 1. Molecular structure of **2a**. H atoms are omitted for clarity, except the freely refined H1. Selected experimental [calculated at the BP86/def2-SVP level for the ground state] bond lengths (Å) and angles (°): Si1–C1 1.952(2) [1.948], Si1–C24 1.824(2) [1.809], Si1–C23 1.961(2) [1.968], C1–N1 1.461(3) [1.465], C24–N2 1.383(3) [1.374]; C23–Si1–C1 101.77(10) [100.67], C2–C1–N1 104.55(17) [105.98], C2–C1–Si1 120.81(16) [118.55], N1–C1–Si1 111.66(15) [111.17], C23–Si1–C24 126.42(10) [127.11], C24–Si1–C1 112.70(10) [114.30], C25–C24–Si1 122.39(15) [122.05], N2–C24–Si1 129.72(17) [129.95], C25–C24–N2 106.16(18) [106.93], C4–N1–C1 114.54(17) [112.91], C21–N1–C1 120.98(17) [122.46], C24–N2–C27 113.51(18) [113.46].

atom (C1) contains one proton, and the silicon atom is additionally bonded to the CMe₂ group (C23) of one of the former isopropyl groups (CHMe₂; C23–Si1 = 1.961(2) Å). This result shows that one of the Me₂(Ar)C–H groups selectively reacted across one of the C_{cAAC}–Si bonds. The C1–Si1 and C1–N1 bonds are elongated from 1.8404(11) to 1.952(2) Å and from 1.3827(15) to 1.461(3) Å, respectively, while the C24(C_{cAAC})–Si1 and C24–N2 bond distances remain very close to the values (C_{cAAC}–Si = 1.8535(12) Å, C24–N2 = 1.3724(14) Å) of silylone **1a**. Finally, a new six-membered ring containing two heteroatoms (shown in purple color in Figure 1) has been formed. The C1–Si1–C23 bond angle of **2a** is 101.77(10)°, while the C1–Si1–C24 angle has changed from 118.16(5)° to 112.70(10)° when compared with that of precursor **1a**.

The Si1 atom is 0.48 Å above the C1–C23–C24 plane, while C24 is located only 0.12 Å above the Si1–C25–N2 plane. All the bond parameters suggest that **2a** might be a singlet diradical (as shown in SI Figure S12) containing one radical each on Si1 and C24 atoms (Si1–C24 = 1.824(2) Å). Similar Si–C bond distances are observed in the monoradical (Cy-cAAC[•])SiCl₃ (181.93(8))¹³ and diradical (Cy-cAAC[•])₂SiCl₂ (1.854(2), 1.843(2) Å). The C_{cAAC}–N bond distance of **2a** is 1.38 Å, close to that of monoradical (Cy-cAAC[•])SiCl₃ (138.27(10)

electron resides in the p orbital of the carbene carbon atom (C24) bonded to the silicon atom (see Mulliken spin density in **IM**₂, SI Figure S14b). Finally, **IM**₂ is oxidized to the product **2b** along with simultaneous reduction of another **1a** species ($\Delta G_L^S = -2.6$ kcal/mol) (SI Table S5).²⁰

In conclusion, we have shown that a substoichiometric amount (33 mol%) of potassium triggers the activation of the unsaturated C_{carbene}:Si:C_{carbene} backbone of very stable silylones **1a,b**, leading to a selective reaction with one intramolecular tertiary C–H bond in an atom-economical approach to form the isomers **2a,b** with a three-coordinate silicon atom. This is the first report on the reactivity studies of silylones. CV suggests that silylone can quasi-reversibly accept an electron to form a very reactive radical anion intermediate, (**1a,b**)^{•−}, which rationalizes the reaction of one of the C–H bonds across C_{carbene}=Si even at room temperature. The energetics of the reaction is computed to be favorable under the given reaction condition, which is important for activation of strong bonds by zero-valent group 14 elements. The electronic structure and bonding properties of the product containing a three-coordinate silicon atom were studied by theoretical calculations, which revealed that it possesses a closed-shell singlet ground state.

■ ASSOCIATED CONTENT

📄 Supporting Information

Syntheses of **2a,b**; UV–vis, crystal structure determination, CV, EPR, and theoretical details. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Authors

dstalke@chemie.uni-goettingen.de

koley@iiserkol.ac.in

hroesky@gwdg.de

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

Dedicated to Prof. Didier Astruc on the occasion of his 68th birthday. We thank Prof. F. Mayer for EPR. H.W.R. thanks the Deutsche Forschungsgemeinschaft (DFG RO 224/60-1) for financial support. D.K. acknowledges the IISER-Kolkata for start-up grant and SERB for DST fellowship (SR/FT/CS-72/2011). D.S. acknowledges partial funding from the DNRF-funded CMC (DNRF93).

■ REFERENCES

(1) Denk, M.; Lennon, R.; Hayashi, R.; West, R.; Belyakov, A. V.; Verne, H. P.; Haaland, A.; Wagner, M.; Metzler, N. *J. Am. Chem. Soc.* **1994**, *116*, 2691–2692.
(2) (a) So, C.-W.; Roesky, H. W.; Magull, J.; Oswald, R. B. *Angew. Chem., Int. Ed.* **2006**, *45*, 3948–3950; *Angew. Chem.* **2006**, *118*, 4052–4054. (b) Driess, M.; Yao, S.; Brym, M.; Wüllen, C. v.; Lentz, D. *J. Am. Chem. Soc.* **2006**, *128*, 9628–9629. (c) Wang, Y.; Xie, Y.; Wei, P.; King, R. B.; Schaefer, H. F., III; Schleyer, P. v. R.; Robinson, G. H. *Science* **2008**, *321*, 1069–1071. (d) Sen, S. S.; Jana, A.; Roesky, H. W.; Schulzke, C. *Angew. Chem., Int. Ed.* **2009**, *48*, 8536–8538; *Angew. Chem.* **2009**, *121*, 8688–8690. (e) Ghadwal, R. S.; Roesky, H. W.; Merkel, S.; Henn, J.; Stalke, D. *Angew. Chem., Int. Ed.* **2009**, *48*, 5683–5686; *Angew. Chem.* **2009**, *121*, 5793–5796. (f) Lee, V. Ya.; Sekiguchi, A. *Organometallic Compounds of Low-Coordinate Si, Ge, Sn and Pb: From Phantom Species to Stable Compounds*; Wiley: New York, 2010.

(3) (a) Mandal, S. K.; Roesky, H. W. *Chem. Commun.* **2010**, 46, 6016–6041. (b) Sen, S. S.; Khan, S.; Nagendran, S.; Roesky, H. W. *Acc. Chem. Res.* **2012**, *45*, 578–587. (c) Bolm, B.; Driess, M. In *Functional Molecular Silicon Compounds II*; Scheschke, D., Ed.; Structure and Bonding 156; Springer: Berlin, 2014; pp 85–123.

(4) Ghadwal, R. S.; Azhakar, R.; Roesky, H. W. *Acc. Chem. Res.* **2013**, *46*, 444–456.

(5) (a) Takagi, N.; Shimizu, T.; Frenking, G. *Chem.—Eur. J.* **2009**, *15*, 8593–8604. (b) Takagi, N.; Shimizu, T.; Frenking, G. *Chem.—Eur. J.* **2009**, *15*, 3448–3456.

(6) (a) Mondal, K. C.; Roesky, H. W.; Schwarzer, M. C.; Frenking, G.; Niepötter, B.; Wolf, H.; Herbst-Irmer, R.; Stalke, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 2963–2967; *Angew. Chem.* **2013**, *125*, 3036–3040. (b) Mondal, K. C.; Samuel, P. P.; Tretiakov, M.; Singh, A. P.; Roesky, H. W.; Stückl, A. C.; Niepötter, B.; Carl, E.; Wolf, H.; Herbst-Irmer, R.; Stalke, D. *Inorg. Chem.* **2013**, *52*, 4736–4743.

(7) Xiong, Y.; Yao, S.; Inoue, S.; Epping, J. D.; Driess, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 7147–7150; *Angew. Chem.* **2013**, *125*, 7287–7291.

(8) Tonner, R.; Öxler, F.; Neumüller, B.; Petz, W.; Frenking, G. *Angew. Chem., Int. Ed.* **2006**, *45*, 8038–8042; *Angew. Chem.* **2006**, *118*, 8206–8211.

(9) Dyker, C. A.; Lavallo, V.; Donnadiou, B.; Bertrand, G. *Angew. Chem.* **2008**, *120*, 3250–3253; *Angew. Chem., Int. Ed.* **2008**, *47*, 3206–3209.

(10) (a) Fürstner, A.; Alcarazo, M.; Goddard, R.; Lehmann, C. W. *Angew. Chem., Int. Ed.* **2008**, *47*, 3210–3214; *Angew. Chem.* **2008**, *120*, 3254–3258. (b) Alcarazo, M.; Lehmann, C. W.; Anoop, A.; Thiel, W.; Fürstner, A. *Nat. Chem.* **2009**, *1*, 295–301.

(11) See SI (S18–S21) for coordination compounds of carbodi-carbene.

(12) (a) Mondal, K. C.; Roesky, H. W.; Schwarzer, M. C.; Frenking, G.; Tkach, I.; Wolf, H.; Kratzert, D.; Herbst-Irmer, R.; Niepötter, B.; Stalke, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 1801–1805; *Angew. Chem.* **2013**, *125*, 1845–1850. (b) Niepötter, B.; Herbst-Irmer, R.; Kratzert, D.; Samuel, P. P.; Mondal, K. C.; Roesky, H. W.; Jerabek, P.; Frenking, G.; Stalke, D. *Angew. Chem., Int. Ed.* **2014**, *53*, 2766–2770; *Angew. Chem.* **2014**, *126*, 2806–2811.

(13) Mondal, K. C.; Roesky, H. W.; Stückl, A. C.; Ehret, F.; Kaim, W.; Dittrich, B.; Maity, B.; Koley, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 11804–11807; *Angew. Chem.* **2013**, *125*, 12020–12023.

(14) Mondal, K. C.; Samuel, P. P.; Roesky, H. W.; Aysin, R. R.; Leites, L. A.; Neudeck, S.; Lübken, J.; Dittrich, B.; Hermann, M.; Frenking, G. *J. Am. Chem. Soc.* **2014**, *136*, 8919–8922.

(15) (a) Martin, D.; Soleilhavoup, M.; Bertrand, G. *Chem. Sci.* **2011**, *2*, 389–399. (b) Martin, C. D.; Soleilhavoup, M.; Bertrand, G. *Chem. Sci.* **2013**, *4*, 3020–3030 and references therein.

(16) Mondal, K. C.; Dittrich, B.; Maity, B.; Koley, D.; Roesky, H. W. *J. Am. Chem. Soc.* **2014**, *136*, 9568–9571.

(17) Bader, R. F. W. *Atoms in Molecules. A Quantum Theory*; Clarendon Press: Oxford, 1990.

(18) Chesnut, D. B. *Chem. Phys.* **2001**, *271*, 9–16.

(19) (a) Mondal, K. C.; Roesky, H. W.; Dittrich, B.; Holzmann, N.; Hermann, M.; Frenking, G.; Meents, A. *J. Am. Chem. Soc.* **2013**, *135*, 15990–15993. (b) Majumdar, M.; Huch, V.; Bejan, I.; Meltzer, A.; Scheschke, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 3516–3520; *Angew. Chem.* **2013**, *125*, 3601–3605.

(20) For the step **IM**₂ → **2a**, the reduction of **1a** is more favored than the step K⁺ → K by 13.2 kcal/mol (ΔG_L^S).