

FLOWERING NEWSLETTER REVIEW

# The (r)evolution of gene regulatory networks controlling *Arabidopsis* plant reproduction: a two-decade history

Alice Pajoro<sup>1,2,\*</sup>, Sandra Biewers<sup>3,\*</sup>, Evangelia Dougalis<sup>4,5,\*</sup>, Felipe Leal Valentim<sup>1,\*</sup>, Marta Adelina Mendes<sup>6,\*</sup>, Aimone Porri<sup>7,\*</sup>, George Coupland<sup>7</sup>, Yves Van de Peer<sup>4,5,8</sup>, Aalt D. J. van Dijk<sup>1,9</sup>, Lucia Colombo<sup>6</sup>, Brendan Davies<sup>3</sup> and Gerco C. Angenent<sup>1,2,†</sup>

<sup>1</sup> Plant Research International (PRI) Droevendaalseweg 1, 6708 PB Wageningen, The Netherlands

<sup>2</sup> Laboratory of Molecular Biology, Wageningen University, Droevendaalseweg 1, 6708 PB Wageningen, The Netherlands

<sup>3</sup> Centre for Plant Sciences, University of Leeds, Leeds LS2 9JT, UK

<sup>4</sup> Department of Plant Systems Biology, VIB, Technologiepark 927, 9052 Ghent, Belgium

<sup>5</sup> Department of Plant Biotechnology and Bioinformatics, Ghent University, Technologiepark 927, 9052 Ghent, Belgium

<sup>6</sup> Dipartimento di BioScienze, Università degli Studi di Milano, Via Celoria 26, 20133, Milan, Italy

<sup>7</sup> Max Planck Institute for Plant Breeding Research, Carl von Linne Weg 10, D-50829 Cologne, Germany

<sup>8</sup> Genomics Research Institute (GRI), University of Pretoria, Private bag X20, Pretoria, 0028, South Africa

<sup>9</sup> Biometris, Wageningen University, Droevendaalseweg 1, 6708 PB Wageningen, The Netherlands

\* These authors contributed equally to the work.

† To whom correspondence should be addressed. E-mail: gerco.angenent@wur.nl

Received 16 March 2014; Revised 16 March 2014; Accepted 28 April 2014

## Abstract

Successful plant reproduction relies on the perfect orchestration of singular processes that culminate in the product of reproduction: the seed. The floral transition, floral organ development, and fertilization are well-studied processes and the genetic regulation of the various steps is being increasingly unveiled. Initially, based predominantly on genetic studies, the regulatory pathways were considered to be linear, but recent genome-wide analyses, using high-throughput technologies, have begun to reveal a different scenario. Complex gene regulatory networks underlie these processes, including transcription factors, microRNAs, movable factors, hormones, and chromatin-modifying proteins. Here we review recent progress in understanding the networks that control the major steps in plant reproduction, showing how new advances in experimental and computational technologies have been instrumental. As these recent discoveries were obtained using the model species *Arabidopsis thaliana*, we will restrict this review to regulatory networks in this important model species. However, more fragmentary information obtained from other species reveals that both the developmental processes and the underlying regulatory networks are largely conserved, making this review also of interest to those studying other plant species.

**Key words:** Flowering time, floral organ development, gene regulation, molecular interactions, networks, transcription factors.

## Plant sexual reproduction

Angiosperms such as *Arabidopsis thaliana* have lifecycles characterized by the alternation of diploid sporophytic and haploid gametophytic stages. Before the sporophyte produces the spores that develop into male and female gametophytes, the plant undergoes several developmental changes that can be considered as phase transitions. The initial vegetative stage can be

subdivided into juvenile and adult phases. Plants in the juvenile phase are able to produce leaves and axillary buds, whereas the initiation of reproductive structures only occurs in the adult vegetative phase. The next phase transition is the switch from vegetative to reproductive growth, when the vegetative shoot apical meristem (SAM) acquires the identity of an inflorescence

meristem (IM) that will then produce floral meristems (FM). Subsequently, floral organs are produced according to a canonical pattern with sepals, petals, stamens, and pistils being formed in sequence, from the periphery to the centre of the floral meristem, to make whorls of organs. Male reproductive development takes place in the third whorl stamens, which are composed of anthers supported on filaments. Inside the anthers the male gametophytes develop by two sequential processes: microsporogenesis and microgametogenesis. Ovules, which form in the fourth whorl, provide structural support to the female gametes and enclose them until seed development, which follows fertilization. The development of the ovule can also be divided in two steps: the specification of the functional megasporangium (megasporogenesis) and the formation of the embryo sac (megagametogenesis) (Reiser and Fischer, 1993; Shi and Yang, 2011). Sexual reproduction requires delivery of the sperm nuclei, via the pollen, to the embryo sac, where a double fertilization occurs and the new diploid sporophyte is formed.

## Gene regulatory networks controlling reproduction

Developmental processes are controlled by tightly coordinated networks of regulators, known as gene regulatory networks (GRNs) that activate and repress gene expression within a spatial and temporal context. In *Arabidopsis thaliana*, the key components of the GRNs controlling major processes in plant reproduction, such as the floral transition and floral organ identity specification, were first identified in loss of function mutants that affect these processes (Blazquez *et al.*, 2006). The interactions between these regulators later began to be revealed through genetic analyses, resulting in the first, mostly linear, GRN maps. These were augmented by reverse genetics, analysis of protein–protein interactions and expression studies in wild-type and mutant plants, resulting in a hierarchical GRN in which master regulators target a subset of genes directing downstream processes (Blazquez *et al.*, 2006) (Fig. 1). Most of these master regulator genes encode transcription factors (TFs), often members of the type II MADS-box family of TFs.

Foreshadowing the complexity that is now becoming very apparent, the early GRN studies already identified several examples of redundancy, where phenotypic consequences that are obvious in mutant combinations, are weak or absent in the corresponding single mutants. For example, individual mutants affecting any of the four *SEPALLATA* genes (*SEPI–4*) have no, or only subtle effects on flowers, whereas the indeterminate flowers of the *sep1sep2sep3* triple mutant are comprised only of sepals (Pelaz *et al.*, 2000) and *sep1sep2sep3sep4* quadruple mutant flowers contain only multiple whorls of leaf-like organs (Ditta *et al.*, 2004).

### GRN for floral transition

Endogenous and environmental signals have their input into the decision to initiate flowering. Several pathways, including photoperiod, autonomous, vernalization, and gibberellin (GA) (Baurle and Dean, 2006; Simpson and Dean, 2002)


converge on a small set of central flowering regulators, including *CONSTANS* (*CO*) and *FLOWERING LOCUS C* (*FLC*), which antagonistically regulate flowering (Fig. 1) (Samach *et al.*, 2000). *FLC* acts as a repressor of flowering and mediates the vernalization and autonomous pathways, whereas *CO* is a floral activator and mediates the photoperiodic pathway. Both genes together regulate the expression of the downstream floral pathway integrator genes *FLOWERING LOCUS T* (*FT*), *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* (*SOC1*), and *LEAFY* (*LFY*) (Blazquez and Weigel, 2000; Simpson and Dean, 2002). The gibberellin pathway influences the phase transition at the SAM by promoting the expression of *SOC1* and *LFY* (Blazquez *et al.*, 1998; Moon *et al.*, 2003) and also acts upstream of *FT* in the leaf, providing evidence for crosstalk between the photoperiod and GA pathways (Porri *et al.*, 2012).

The *FT* protein stimulates flowering by moving from the leaf (where the light is perceived) to the shoot apical meristem (where inflorescence and floral meristems form). Genetic and molecular studies indicate that the *FT* protein comprises part of the inductive signal known as florigen, which promotes flowering in response to photoperiod (Corbesier *et al.*, 2007; Jaeger and Wigge, 2007). *FT* interacts at the SAM with the bZIP TF *FLOWERING LOCUS D* (*FD*) to activate the expression of the floral meristem identity genes *APETALA1* (*API*) and *SOC1*, which in turn activate *LFY*, promoting the floral transition (Wu and Gallagher, 2012). In addition to *LFY*, other genes such as *AGAMOUS LIKE 24* (*AGL24*) also contribute to the up-regulation of *API* (Grandi *et al.*, 2012; Pastore *et al.*, 2011).

The microRNAs miR156 and miR172 have been identified as important regulators of this developmental phase change. Members of the *SQUAMOSA PROMOTER BINDING PROTEIN LIKE* (*SPL*) TF family are targeted by miR156, whereas miR172 targets 6 *APETALA2-LIKE* (*AP2-like*) transcription factors (Wu *et al.*, 2009). Levels of miR156 are high in the juvenile vegetative phase and decrease before onset of the adult vegetative phase, allowing the production of a subset of SPL proteins (*SPL9*, *SPL10*). These SPLs induce the expression of miR172 genes, leading to a gradual increase in miR172 levels during the adult vegetative phase (Aukerman and Sakai, 2003; Jung *et al.*, 2007; Schwarz *et al.*, 2008; Wu and Poethig, 2006; Wu *et al.*, 2009). Increased levels of miR172, in turn, cause a down-regulation of AP2-like genes, which otherwise repress adult traits and flowering (Fig. 1) making the plant competent to flower (Wu *et al.*, 2009).

### GRN for floral organ development

After successful initiation of the floral meristems, *API* and *LFY* activate the organ identity genes (Fig. 1). The way in which this limited set of genes acts to direct formation of the correct floral organs at the appropriate place in the flower is described by the classical ABC model (for reviews see: Alvarez-Buylla *et al.*, 2010; Causier *et al.*, 2010; Immink *et al.*, 2010; O'Maoileidigh *et al.*, 2014). The A-class genes are the MADS-box TF-encoding *API* (Mandel and Yanofsky, 1995) and *APETALA2* (*AP2*), which encodes a member of the AP2/ERF (ethylene



**Fig. 1.** Linear GRN of genes controlling flower formation. The switch from vegetative to reproductive phase is triggered by endogenous and environmental stimuli, some examples of which are illustrated here. These signals converge at the central flowering regulator genes FLC and CO that antagonistically regulate the floral integrator genes at the SAM. The floral integrators activate the meristem identity genes *AP1* and *LFY*, subsequently leading to activation of the ABCDE class genes, specifying organ identity. The endogenous aging pathway involves micro RNAs (miRNAs). At early stages of development, the level of miR156 is high, maintaining the juvenile growth phase. As the plant ages, miR156 levels decrease, allowing the production of a subset of SPL proteins. These SPL proteins induce the expression of MIR172 genes, which are consequently expressed at low levels in the juvenile phase and steadily increase their expression levels in the adult phase. Elevated levels of miR172 cause down-regulation of AP2-like TF factor genes, which otherwise repress flowering. Arrows indicate activation, blocked lines indicate repression, left-right arrows indicate a positive feedback loop. Abbreviations: AGL24, AGAMOUS LIKE 24; AP1, APETALA1; CO, CONSTANS; FD, FLOWERING LOCUS D; FT, FLOWERING LOCUS T; LFY, LEAFY; STK, SEEDSTICK; SEP, SEPALLATA; SHP, SHATTERPROOF; SOC1, SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1; SVP, SHORT VEGETATIVE PHASE; SPL, SQUAMOSA PROMOTER BINDING PROTEIN-LIKE; W1, whorl 1; W2, whorl 2; W3, whorl 3; W4, whorl4.

responsive factor) TF family (Jofuku *et al.*, 1994; Okamuro *et al.*, 1997). The B-class genes are *APETALA3* (*AP3*) (Jack *et al.*, 1992) and *PISTILLATA* (*PI*) (Goto and Meyerowitz, 1994), both of which encode MADS-box TFs. *AGAMOUS* (*AG*), another MADS-box gene, is the only member of the C-class (Yanofsky *et al.*, 1990). The original ABC model was extended by the addition of D- and E-classes, which specify ovule identity (Pinyopich *et al.*, 2003) and floral identity, respectively. The D-class genes include the MADS-box genes *SEEDSTICK* (*STK*) and *SHATTERPROOF1* and 2 (*SHP1/2*), which are redundantly required for the specification of ovule identity (Pinyopich *et al.*, 2003). Four *SEPALLATA* genes (*SEP1–4*), all of which encode MADS-box TFs, comprise the E-class (Ditta *et al.*, 2004; Pelaz *et al.*, 2000; Rounseley *et al.*, 1995) (Fig. 1). Protein–protein interaction studies demonstrated that the ABCDE MADS-box TFs physically interact and bind DNA as tetramers, possibly establishing DNA loops in the promoters of their target genes to activate or repress their transcription

(Egea-Cortines *et al.*, 1999; Melzer and Theissen, 2009; Smacziak *et al.*, 2012b). The E-class SEP proteins are necessary for the formation of higher order complexes involving the A-, B- and C-class TFs and hence mediate their organ identity functions (Honma and Goto, 2001; Pelaz *et al.*, 2001). The resultant hierarchical GRN formed the basis of our understanding of flower development until the advent of new technologies described below.

## The modern version of a GRN

Recently, genome-wide molecular approaches, such as protein assays, expression profiling and chromatin immunoprecipitation (ChIP) have connected the nodes in the GRNs and added novel regulators, interactors, and downstream targets (Hawkins *et al.*, 2010). In contrast to the earlier genetic networks described above, more recent versions show molecular interactions that are independent of mutant phenotypes and are potentially more dynamic in time and space. The

molecular interactions that were added are, for example, protein–protein interactions, which allow combinatorial actions of multiple regulatory factors. In the case of TF complexes, their composition can be an important determinant of DNA binding specificity and affinity and thus interactions will affect target gene regulation. Another layer of molecular interactions recently added to the networks comprises transcription factor binding profiles, i.e. protein–DNA interactions. Large-scale expression analyses, preferably performed with inducible systems (Kaufmann *et al.*, 2010c; Wellmer *et al.*, 2006), add transcriptional activity to the connections ('edges') between nodes in the network. These large-scale data sets have substantially increased the network connectivity and revealed the shortcomings of the classical hierarchical networks. Current GRNs are composed of TFs regulating subsets of genes in a combinatorial fashion and contain multiple regulatory feedback loops, which blur the hierarchical structure (Kaufmann *et al.*, 2010a).

TFs act as process integrators and connect other developmental processes as hubs in the network (Pose *et al.*, 2012). The TF target gene analyses have revealed unexpected connections between processes previously considered to be unrelated, which could not have been predicted by classical genetic approaches. For example, target gene analysis of the SEP3 floral organ identity transcription factor revealed a connection with auxin signalling (Kaufmann *et al.*, 2009). As another striking example, a recent study addressing the role of LFY revealed that in addition to its function in reproductive transition, it is also involved in pathogen responses (Winter *et al.*, 2011).

#### DNA–protein interaction studies provided novel insights into the wiring of GRNs

In the 90s, binding of TFs to DNA sequences was studied by *in vitro* methods, such as electrophoretic mobility shift assays (EMSA) and yeast 1-hybrid studies. Schwarz-Sommer's research group (Tröbner *et al.*, 1992) showed that the Antirrhinum class B homeotic proteins DEFICIENS and GLOBOSA interact with each other and bind *in vitro* to their own promoters, thereby forming an autoregulatory loop. More such autoregulatory loops have been recently identified in GRNs, particularly associated with transcriptional regulation of MADS-box genes (Gomez-Mena *et al.*, 2005). For instance, expression of the MADS-box gene *SOC1* is controlled by the SOC1 protein, which forms heterodimers with AP1 that suppress *SOC1* expression following the successful transition to floral meristem identity (Immink *et al.*, 2012).

SELEX (systematic evolution of ligands by exponential enrichment) is a powerful method to characterise TF binding sites and has been applied to factors in the flowering GRN. For example, Moyroud *et al.* (Moyroud *et al.*, 2011) applied SELEX coupled to next generation sequencing (NGS) to determine the preferred binding sites of LFY.


The development of ChIP was a major breakthrough in the study of DNA–protein interactions, as it allowed the identification of *in vivo* physical interactions between a TF and its target DNA (Gomez-Mena *et al.*, 2005;

Wang *et al.*, 2002). The subsequent introduction of genome-wide arrays (ChIP–ChIP) (Zheng *et al.*, 2009) or large-scale sequencing (ChIP-seq) (Kaufmann *et al.*, 2010b) led to the identification of thousands of target genes for specific TFs. In the past 5 years many genome-wide TF–DNA binding profiles have been generated for TFs involved in plant reproduction (Table 1). The high numbers of interactions that were detected in ChIP-seq experiments revealed a much higher network complexity than expected and demonstrated that these major regulators not only control another layer of regulators, but also genes encoding structural proteins, enzymes and signalling proteins (Fig. 2; Kaufmann *et al.*, 2010a). The ChIP-seq data also confirmed that most TFs involved in plant reproduction bind to their own locus, reinforcing the concept that auto-regulatory loops are a common mechanism of regulation in GRNs. For example, identification of the direct targets of LFY (Winter *et al.*, 2011) and AP1 (Kaufmann *et al.*, 2010c) revealed that LFY is able to promote *API* transcription through direct regulation, and AP1 binds to *LFY* to form a positive feedback loop, leading to a strong and rapid up-regulation of both genes.

Through these genome-wide studies, in particular in combination with transcriptome analysis (Kaufmann *et al.*, 2010c; Wuest *et al.*, 2012), it also became apparent that some TFs act as both activators and repressors. For example, at early stages AP1 represses genes controlling flowering time, whereas at later stages it mainly acts as an activator of floral homeotic genes (Kaufmann *et al.*, 2010c; Pajoro *et al.*, 2014). A similar observation was made for the B-class genes *AP3* and *PI*, which activate genes involved in organogenesis and repress those required for carpel development (Wuest *et al.*, 2012). It is likely that the composition of the TF complexes and their ability to recruit co-factors act together to determine the DNA-binding specificity and the mode of transcriptional action. Detailed analyses of the DNA regions bound by several members of MADS-box TF family using the motif-based sequence analysis tool MEME (Machanick and Bailey, 2011) revealed distinct sequence specificities for each MADS-box TF. Although the consensus binding site for MADS-box proteins, the so-called CArG box, was found at the centre of virtually all binding peaks obtained in ChIP experiments, certain sequence motifs within and flanking the CArG box were preferentially enriched. This corroborates the idea that the DNA sequence specificity is determined to a large extent by the composition of the TF complex. Therefore, protein–protein studies will be required to enable us to understand TF–DNA interaction specificity and hence why specific target genes are recognised by specific transcription factors. Recent large-scale protein–protein and proteomics studies have elucidated the composition of many MADS-box protein complexes involved in flowering time control and floral organ development (Immink *et al.*, 2009; Smaczniak *et al.*, 2012a). The next challenge in these studies will be to unravel the binding specificity of these complexes and how this influences the dynamic control of target gene regulation.

**Table 1.** Overview of genome-wide protein-DNA binding profile studies for TFs involved in plant reproduction

Gene	Family	Function	Approach	Antibody	Tissue	Most relevant targets	Reference	
AGAMOUS	AG	MADS-box transcription factor	ChIP-seq	Anti GFP	Flower buds stage 5	AG, CRC, SHP2, SPL, JAG, SEP3, AP1, AP3, SHP1, SUP, HEC1, HEC2, VDD	O'Maoileidigh et al., 2013	
AGAMOUS-LIKE 15	AGL15	MADS-box transcription factor	Floral transition	ChIP-ChIP	Anti AGL15	Embryonic culture tissue	FLC, SVP, LEC2, FUS3, ABI3, IAA30	Zheng et al., 2009
APETALA1	AP1	MADS-box transcription factor	Repressor of floral transition	ChIP-seq	Anti AP1	Inflorescence meristem; flowers at stage 2, 4, and 8	FD, FDP, LFY, SNZ, TOE1, TOE3, TEM1, TEM2, TFL1, SPL9, SPL15, SEP3, AP2	Kaufmann et al., 2010; Pajoro et al., 2014
APETALA2	AP2	AP2-like family	Floral transition and floral organ development	ChIP-seq	Anti AP2	Inflorescences	AG, SOC1, SEP3, AP1, TOE3, AGL15, ETT, SHP1, SHP2, AGL44, TOE1, RGA-like1, miR156, miR172	Yant et al., 2010
APETALA3	AP3	MADS-box transcription factor	Floral organ development	ChIP-seq	Anti GFP	Flower buds stage 5	CRC, SEP3, SPL, AP1, SUP, AG, UFO, SHP2, RBE, HEC1, HEC2, ALC	Wuest et al., 2012
FLOWERING LOCUS C	FLC	MADS-box transcription factor	Flowering time	ChIP-seq	Anti FLC	12-day-old seedlings	SOC1, FT, SEP3, CBF1, JAZ6, AGL16, SPL15, DIN10, SVP, SPL3, SMZ, TOE3, TEM1, FRI, CIR1, FIO1, LCL1, COL1	Deng et al., 2011
FLOWERING LOCUS M	FLM	MADS-box transcription factor	Flowering time	ChIP-seq	Anti GFP	15-day-old seedlings	SOC1, ATC, TEM2, SMZ, SEP3, AP3, PI, RVE2, FIO1, SHP2, MIR156, AP2, MIR172, AP1	Pose et al., 2013
LEAFY	LFY		Floral transition	ChIP-seq	Anti LFY	Inflorescences	TFL1, AP1, AG, SEP4, LFY, SOC1, PRS, BB, GIS, GOA, STY2, ARR3, GA3OX2	Moyroud et al., 2011
PISTILLATA	PI	MADS-box transcription factor	Floral organ development	ChIP-seq	Anti GFP	Flower buds stage 5	CRC, SEP3, SPL, AP1, SUP, AG, UFO, SHP2, RBE, HEC1, HEC2, ALC	Wuest et al., 2012
SCHLAFMUTZE	SMZ	AP2-like family	Repressor of flowering	ChIP-ChIP	Anti GFP	Seedlings	FT, SMZ, SNZ, AP2, TOE3, SOC1, AP1, TEM1, FRI	Mathieu et al., 2009
SEPALLATA3	SEP3	MADS-box transcription factor	Flower development	ChIP-seq	Anti SEP3	Inflorescences (stage 1–12); flowers at stage 2, 4, and 8	AP1, AP3, SEP1, SEP2, SEP4, AG, SHP1, SHP2, GA1, PIN4, PID, ETT, ARG8, IAA4	Kaufmann et al., 2009; Pajoro et al., 2014
SHORT VEGETATIVE PHASE	SVP	MADS-box transcription factor	Flowering time	ChIP-seq	Anti GFP	2 weeks old seedlings	GI, PRR7, FLK, FLD, CLF, SWN, VNR2, PHYA, STIP, SVP, CLV1, CLV2, PHB, PHV, REV, ATHB8	Gregis et al., 2013
SHORT VEGETATIVE PHASE	SVP	MADS-box transcription factor	Floral organ development	ChIP-seq	Anti GFP	Inflorescences (stage 1–11)	SVP, CLV1, PHB, KAN1, ETT, PIN1, WDR55	Gregis et al., 2013
SUPPRESSOR OF OVEREXPRESSION OF CONSTANS1	SOC1	MADS-box transcription factor	Flowering time	ChIP-seq	Anti GFP	Transition apices	SOC1, CBF1, CBF2, CBF3, mir156, SVP, AGL15, AGL18, TEM2, TOE3, SMZ, SNZ, SEP3, SHP2, AP3, PI, SUP, ETT	Immink et al., 2012



**Fig. 2.** Gene interaction network based on DNA-binding profiles. TFs involved in plant reproduction (Table 1) show a high overlap in target genes. Most TFs bind to their own locus, suggesting that auto-regulatory loops are a common mechanism of regulation. Many pairs of TFs show a reciprocal binding suggesting a mechanism of inter-regulation. Abbreviations: AGAMOUS, AG; AGAMOUS LIKE 24, AGL24; APETALA1, AP1; APETALA2, AP2; APETALA3, AP3; CURLY LEAF, CLF; CLAVATA1, CLV1; COSTANS, CO; ETTIN, ETT; FLOWERING LOCUS C, FLC; FLOWERING LOCUS D, FD; FLOWERING LOCUS T, FT; FLOWERING LOCUS M, FLM; FLOWERING LOCUS KH DOMAIN, FLK; FRUITFULL, FUL; GIGANTEA, GI; LEAFY, LFY; PISTILLATA, PI; SEPALLATA1-4, SEP1-4; SHATTERPROOF1-2, SHP1-2; SQUAMOSA PROMOTER BINDING PROTEIN-LIKE, SPL; SCHLAFMUTZE, SMZ; SCHNARCHZAPPEN, SNZ; SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1, SOC1; SHORT VEGETATIVE PHASE, SVP; SWINGER, SWN; TEMPRANILLO2, TEM2; TERMINAL FLOWER 1, TFL1; TARGET OF EARLY ACTIVATION TAGGED (ET) 3, TOE3; REDUCED VERNALIZATION RESPONSE 2, VRN2.

### SOC1 and SHORT VEGETATIVE PHASE (SVP), two MADS domain proteins acting as hubs in the GRNs

As an example of the progress made regarding the topology and complexity of the GRNs described above, we will highlight some genome-wide analysis of regulators and targets of the MADS domain proteins SVP and SOC1. Owing to its important role as a floral integrator, SOC1 is of particular interest, because it may act as a hub between the GRNs involved in vegetative and reproductive meristems. In this respect, both upstream regulators and downstream targets of *SOC1* are relevant. *SVP* has a dual function, reflected in its expression pattern: in vegetative tissues SVP acts as a floral suppressor, whereas during the first stages of floral meristem development SVP prevents the precocious expression of the floral organ identity genes (Gregis *et al.*, 2006). Recently, the genome-wide SVP binding sites in both tissues were determined and the difference reflects the dual role of this transcription factor (Gregis *et al.*, 2013).

A small set of target genes that act downstream of SOC1 were identified by a microarray expression analysis (Seo *et al.*, 2009) and genome-wide analysis of binding sites based on a ChIP-ChIP experiment using a *SOC1* overexpression line (Tao *et al.*, 2012). More recently, a genome-wide target gene analysis was performed by ChIP-seq on enriched apical meristem tissues expressing GFP-tagged SOC1 protein under the endogenous promoter during floral transition (Immink *et al.*, 2012). This study revealed that floral timing and flower organogenesis networks are heavily interconnected via SOC1 acting as a transition hub and being involved in many positive and negative auto-regulatory feedback loops. Several other MADS-box genes were also found to be bound by SOC1. The floral repressors *SVP*, *AGL15*, and *AGL18* were identified as direct targets of SOC1. Furthermore, SOC1 also binds to the regulatory regions of several floral homeotic genes, such as the MADS-box genes *SEP3*, *AP3*, *PI*, and *SHP2* (Immink *et al.*, 2012). Genes encoding AP2/EREBP transcription factors that are involved in the response to low temperature,

such as CRT/DRE-BINDING FACTOR1 (CBF1), CBF2, and CBF3, were bound by SOC1. Seo and collaborators (Seo *et al.*, 2009) demonstrated that SOC1 down-regulates these CBFs and the ChIP-seq data confirmed that this down-regulation is direct (Immink *et al.*, 2012). SOC1 also regulates a number of other AP2-like genes acting predominantly as suppressors of flowering, such as *TEMPRANILLO2* (*TEM2*), *AP2*, *TARGET OF EAT3* (*TOE3*), *SCHLAFMUTZE* (*SMZ*), and *SCHNARCHZAPFEN* (*SNZ*). Except AP2, these factors suppress *SOC1* expression either directly or indirectly via FT.

Because multiple (auto)regulatory feedback loops exist and because SOC1, as a floral integrator, receives many signals from the different flowering pathways, upstream regulators of *SOC1* were also determined (Immink *et al.*, 2012). A comprehensive matrix-based yeast one-hybrid assay was performed using 135 type II MADS domain protein dimers, available from an *Arabidopsis* MADS dimer collection in yeast (de Folter *et al.*, 2005; Immink *et al.*, 2009; Immink *et al.*, 2012). Although this approach was targeted to only MADS domain proteins, it contributed to a more defined GRN around SOC1. Dimers consisting of proteins involved in floral timing and floral transition, such as SVP–AGL15 and FUL–SOC1, were found to bind to the *SOC1* regulatory sequences. Remarkably, a large number of DNA–TF interactions were also found with dimers containing “ABC-class” homeotic proteins, for instance AG–SEP3, AGL24–AP1, and SOC1–SEP3 (Immink *et al.*, 2012). Previous studies (Kaufmann *et al.*, 2010c; Liu *et al.*, 2008) showed that AP1 is involved in the repression of *SOC1* in the floral meristem, whereas the yeast one-hybrid data confirmed the binding of AP1, in combination with particular dimerization partners, to the *SOC1* promoter. These yeast data were confirmed by *SOC1* reporter lines, which showed down-regulation of the reporter when either *AP1*, *SEP3*, or *AG* expression was induced (Immink *et al.*, 2012).

*SVP* is a key regulator of two developmental phases: during the vegetative phase it represses the floral transition and later it contributes to the specification of floral meristems. To maintain plants in the vegetative phase, SVP represses the expression of *FT* and *TWIN SISTER OF FT* (*TSF*) in the phloem and *SOC1* in the SAM by directly binding to CArG boxes in *FT* and *SOC1* promoter regions (Jang *et al.*, 2009; Lee *et al.*, 2007; Li *et al.*, 2008). SVP interacts with FLC and the FLM- $\beta$  splicing variant, both dimers binding to the *SOC1* promoter as repressor, whereas the SVP/FLM- $\delta$  dimer, which is not able to bind DNA, competes with the SVP/FLM- $\beta$  dimer and acts as a promoter of flowering (Pose *et al.*, 2013). Another competitor of *SVP* is its phylogenetically closest related MADS-box gene *AGL24*, which is a central promoter of flowering (Michaels *et al.*, 2003; Parenicova *et al.*, 2003; Yu *et al.*, 2002). Both SVP and AGL24 directly regulate *SOC1* by binding to the same binding sites in the promoter but they have an opposite effect on *SOC1* expression (Liu *et al.*, 2008). During the floral transition, *SVP* expression gradually decreases in the IM and becomes expressed again at the early stages of flower development (Gregis *et al.*, 2009).

Two recent studies (Gregis *et al.*, 2013; Tao *et al.*, 2012) reported the targets of SVP at a genome-wide scale. A comparison between targets in the SAM and after the floral transition in the FM revealed targets common and specific to its developmental roles (Gregis *et al.*, 2013). SVP binds directly to flowering-time genes involved in the different flowering pathways including the circadian clock and photoperiodic pathway represented by *GIGANTEA* (*GI*) and *PSEUDO-RESPONSE REGULATOR 7* (*PRR7*), the autonomous pathway represented by genes such as *FLOWERING LATE KH MOTIF* (*FLK*) and *FLOWERING LOCUS D* (*FD*), the light signalling pathway represented by *PHYTOCHROME A* (*PHYA*), as well as to genes that encode components of chromatin-associated complexes such as *CURLY LEAF* (*CLF*), *SWINGER* (*SWN*), and *VERNALIZATION2* (*VRN2*). During the early stages of flower development (stages 1 and 2) *SVP* controls the expression of homeotic genes to maintain the floral meristem in an undifferentiated state (Gregis *et al.*, 2006). This role is executed together with AP1 and together this dimer recruits the SEU-LUG repressor complex. Comparison of regions bound by AP1 and SVP revealed many common targets, such as *CLAVATA1* (*CLV1*) and *AG*, which are down-regulated by SVP (Gregis *et al.*, 2013). *CLV1* and *AG* are both suppressors of *WUSCHEL* (*WUS*), which maintains the meristematic activity in the FM and controls the size of the FM. At late stage 2, when SVP expression is switched off, AP1 interacts with SEP3 and the repression of the homeotic genes is removed.


## Connecting GRNs and hormonal pathways in plant reproduction

The ChIP-seq data published recently demonstrate that MADS-domain TFs are able to bind to thousands of different genomic regions, several of which belong to genes involved in different hormonal pathways (Table 1). To understand the connection between MADS-domain TFs and the different hormone pathways, we analysed each ChIP-seq dataset available for these proteins and grouped them into specific hormone-related subclasses. Among the targets of these transcription factors involved in flowering time and floral development are genes involved in auxin, gibberellin, jasmonic acid, ethylene, brassinosteroid, salicylic acid and abscisic acid pathways. Different functional sub-categories can be recognized for each hormone pathway (Table 2).

*SEP3*, *AP1*, and *AG* are the TFs that score the highest number of sub-categories, meaning that they are involved in almost all hormone pathways. This information is in accordance with previous findings, where *SEP3* was found to integrate and modulate different growth-related and hormone pathway genes, particularly those connected with auxin signalling (Kaufmann *et al.*, 2009). *AP1* directly controls the expression of genes with known functions in the control of organ growth such as genes involved in the biosynthesis and response to gibberellin (GA) (Kaufmann *et al.*, 2010c). A recent publication about *AG* direct targets illustrated that approximately half of the 225 identified genes encode

**Table 2.** Targets of TFs based on ChIP datasets divided into different hormone-related sub-categories

For SVP, two different tissues were used for the identification of target genes: inflorescences and seedlings.



proteins with regulatory functions, including many transcription factors, (receptor) kinases, putative ligands, and proteins involved in different phytohormone signalling pathways (O'Maoileidigh *et al.*, 2013).

Genes belonging to “response to abscisic acid (ABA) stimulus”, and “ABA-mediated signalling pathway” categories are bound by almost all TFs analysed. ABA is known to play important roles in many aspects of seed development, such as accumulation of storage compounds, acquisition of desiccation tolerance, induction of seed dormancy, and suppression of precocious germination (Kanno *et al.*, 2010). Most likely this hormone plays unknown functions during flower development. GA and ABA also play important roles in regulating the floral transition. For instance, Shan and colleagues (Shan *et al.*, 2012) showed that an increase in sensitivity to

ABA delays flowering time. Gibberellins promote flowering in *Arabidopsis* through the activation of the floral integrator genes *SOC1*, *LFY*, and *FT* in the inflorescence and floral meristems, and in leaves, respectively.

Among the many roles of auxin is its function in the outgrowth and development of lateral organs, including floral organs, which has been suggested based on mutant phenotypes (Immink *et al.*, 2012) and *SEP3* targets (Kaufmann *et al.*, 2009). ETTIN (ETT) imparts regional identity in the FM by affecting perianth organ number spacing, stamen formation, and regional differentiation of stamens and gynoecium (Sessions *et al.*, 1997). The ChIP-seq analysis with SVP (Gregis *et al.*, 2013) also identified target genes involved in the auxin signalling pathway, such as *BIG*, which encodes a putative auxin transporter required for normal auxin efflux

and inflorescence development (Gil *et al.*, 2001; Yamaguchi *et al.*, 2007).

These examples show the integration of the flowering and flower development GRNs in hormone signalling, although it is far from understood how these TFs control the hormone pathways and lead to the initiation and differentiation of the flower.

## Tissue-specific characterization of gene regulatory networks: Spatial action and short/long distance movement


The current GRNs are still relatively static and lack detailed information about the spatial and temporal expression dynamics of genes that are part of the network. The networks active in the leaf and the SAM are relevant to the floral transition and spatial expression patterns of a number of key regulators are available for these tissues. Examples include members of the *SPL* gene family, which encode transcription factors that regulate flowering in *Arabidopsis* (Wang *et al.*, 2009; Wu *et al.*, 2009). The temporal and spatial expression patterns of *SPL* genes were studied using laser capture microdissection coupled to transcriptome sequencing and *in situ* hybridization (Cardon *et al.*, 1999; Porri *et al.*, 2012; Torti *et al.*, 2012; Wang *et al.*, 2009). These studies showed that several *SPL* genes, including *SPL3*, *SPL4*, *SPL5*, and *SPL9* are expressed in both leaf and shoot apical meristems of *Arabidopsis*. *SPL4* and *SPL5* expression rises in the centre of the meristem of wild-type plants during the transition to flowering, when plants are exposed to inductive long-day conditions through the *FT* pathway (Torti *et al.*, 2012). In contrast, *SPL9* mRNA is detectable at the flanks of the inflorescence meristem (Porri *et al.*, 2012; Wang *et al.*, 2009). Recently it was shown that *SPL* proteins are regulated by interaction with the transcriptional repressor DELLA proteins that is degraded in response to GA, indicating that these transcription factors function as important integrators of distinct flowering pathways (Yu *et al.*, 2012). In the presence of active GA, DELLA proteins are ubiquitinated and targeted for degradation by the 26S proteasome (Dill *et al.*, 2004; Willige *et al.*, 2007). As DELLA proteins bind *in vitro* and *in vivo* to several members of the *SPL* family, such as *SPL2*, *SPL9* and *SPL10*, it was suggested that the presence of active GA is an essential step to produce functional *SPL* proteins by releasing them from DELLA repressors (Yu *et al.*, 2012). Downstream events of *SPL* gene activity include the activation of *FT* transcription through the module miR172/AP2-like, a process that occurs specifically in the leaf (Fig. 3A). *SPL9* and *SPL10* directly activate transcription of *MIR172*, which produces a mature miRNA that cleaves, or prevents translation of, transcripts encoding AP2-like transcription factors (Mathieu *et al.*, 2009). One of these factors is SMZ, which binds the *FT* locus 1.5 kb downstream of the coding region to repress its expression (Mathieu *et al.*, 2009). Therefore, in the presence of active GA, *SPL* proteins lead to *FT* activation by reducing *AP2-like* transcript levels by promoting *MIR172* expression (Fig. 3A). At the shoot apical meristem *SPLs* transcription is also dependent on GA and GA signalling (Galvao *et al.*, 2012; Porri *et al.*, 2012).

Transgenic plants expressing a GA catabolic enzyme specifically at the SAM, showed significant delay in *SPL* gene transcriptional activation when plants were exposed to inductive LD (Porri *et al.*, 2012).

Downstream events of *SPL* genes include the activation of key flowering genes. *SPL9* and *SPL3* bind To *SOC1* and *LFY* loci respectively, providing a direct mechanism by which *SPLs* act at the meristem to promote the switch from the vegetative to reproductive phase (Wang *et al.*, 2009; Yamaguchi *et al.*, 2009) (Fig. 3B). On the contrary, *SOC1* was found to bind *SPL3*, *SPL4*, and *SPL5* promoters, suggesting the presence of a positive feedback loop between *SOC1* and *SPLs* in flowering time control (Jung *et al.*, 2012). Furthermore, *SOC1* and the related transcription factor *FRUITFUL* (*FUL*) were shown genetically to be required for *SPL4* activation during floral transition (Torti *et al.*, 2012). Thus, *SPL* genes have different genetic and molecular functions during floral transition, and these are spatially separated between leaf and shoot apex (Fig. 3A, B).

Advances in understanding GRNs also provide information about the local and long-distance movement of hormones, proteins, or RNAs, which may be produced in specific cell types and move to other cells to regulate pathways. In this way, mobile factors can act as a hub between spatially separated GRNs. A striking example is *FT*, which encodes a small protein that functions as a master regulator of *Arabidopsis* flowering (Kardailsky *et al.*, 1999; Kobayashi *et al.*, 1999). *FT* was initially identified through a late flowering mutant (Koornneef *et al.*, 1991). Genetic interaction between *ft* and other flowering mutants identified important components of the *FT* pathway (Koornneef *et al.*, 1991). These include the upstream circadian clock component *GI* and the downstream acting gene *CO*, whose product activates directly *FT* transcription in response to long-day (Sawa and Kay, 2011; Schaffer *et al.*, 1998; Suarez-Lopez *et al.*, 2001) (Fig. 3A). More recently, studies revealed the exact tissues where *FT* is expressed. A *GUS* reporter gene fused to the *FT* promoter sequence showed GUS activity specifically in the companion cells of the leaf throughout the vasculature (Adrian *et al.*, 2010; Takada and Goto, 2003). Although *FT* is expressed in the vasculature, it promotes flowering at the shoot apex, because the *FT* protein moves from companion cells to the apex to trigger flowering (Corbesier *et al.*, 2007; Jaeger and Wigge, 2007; Mathieu *et al.*, 2007). Thus, *FT* is a long-distance signalling molecule that connects flowering promoting signals in the leaves with the GRN that is active in the SAM (Fig. 3). In addition, *FT* acts redundantly with a closely related gene *TSF*, but whether *TSF* protein also travels from leaf to the SAM is not yet understood (Jang *et al.*, 2009; Yamaguchi *et al.*, 2005).

Plant hormones are mostly small organic molecules with the ability to move from cell to cell through specific transmembrane transporters. Auxin, cytokinin and GA were shown to affect flowering of *Arabidopsis* through interconnected genetic pathways (Richter *et al.*, 2013). Although specific cell-cell transporters have not yet been identified for GA, recent experimental evidence suggests that this hormone moves among plant tissues. Grafting experiments performed



**Fig. 3.** Mobile factors are crucial for flowering induction in *Arabidopsis*. (A) *FT* expression regulation in the leaf. The photoperiod pathway activates transcription of *FT* through CO and GI to promote flowering specifically under long days. CO protein is stabilized by light and it activates *FT* expression in the companion cells of the leaf. Gibberellin (GA) activates the transcription of *FT* through the *SPL/miR172* module. GI positively regulates the abundance of miRNA172 independently of CO. The *FT* protein is uploaded in the sieve element and then moves towards the SAM where it triggers the transition to flowering. Movement of *FT* occurs through the phloem system probably by a passive transport mechanism mediated by the plasmodesmata. (B) *FT* protein is downloaded at the base of the SAM. *FT* forms a heterodimer complex with the b-ZIP transcription factor FD, which activates transcription of the floral integrator SOC1. SOC1 activates in turn transcription of *SPLs* including *SPL3*, *SPL4*, and *SPL5*. Activation of *SPLs* expression is crucial for floral transition by activating the floral meristem identity genes LFY and AP1, two genes that specify floral primordia at the flanks of the SAM. GA is also involved in the activation of *SPLs* both at the transcriptional and post-transcriptional levels. Arrows indicate activation and blocked lines indicate repression. Arrows with dashed line represent indirect interaction. Abbreviations: Gibberellin, GA; APETALA1, AP1; CONSTANS, CO; FLOWERING LOCUS T, FD; FLOWERING LOCUS T, FT; GIGANTEA, GI; LEAFY, LFY; SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1, SOC1; SHORT VEGETATIVE PHASE, SVP; SQUAMOSA PROMOTER BINDING PROTEIN-LIKE, SPL.

in *Arabidopsis*, suggested that GA is the mobile signal that induces thickness of the hypocotyl by increasing the production of xylem tissue (Ragni *et al.*, 2011). Xylem formation is associated with the transition to flowering and with increasing expression of the *GIBBERELLIN 3-OXIDASE 1*

(*GA3ox1*) (which encodes an enzyme involved in GA biosynthesis) in the SAM. This generates increased levels of GA that migrates from the shoot to the hypocotyl to induce xylem formation, as demonstrated by a grafting experiment, in which a *Landsberg* scion rescued the defective xylem expansion of the

GA mutant *gal-3* (Ragni *et al.*, 2011). Long distance movement of active GA may occur from the leaf to the shoot apex. Labelled GA<sub>4</sub> applied to the leaf could be detected at the shoot apex of *Arabidopsis* (Eriksson *et al.*, 2006), implying that the hormone is capable of long-range movement, presumably through the vasculature.

Recently, fluorescein labelled GA (Fl-GA) was used to trace GA distribution in the root system (Shani *et al.*, 2013). Application of Fl-GA<sub>3</sub> in *Arabidopsis* roots showed specific GA accumulation patterns in the endodermal cells of the elongation zone. In addition, *REPRESSOR OF GA* (*RGA*) marker (*GFP:RGA*), which encodes a DELLA protein that is degraded in the presence of active GA, showed decreasing signal in the endodermal cells, indicating that GA accumulates in these cells in wild-type plants (Shani *et al.*, 2013). These new experimental approaches could be useful to further investigate a possible role of GA movement in flowering time control of *Arabidopsis*.

## Modelling plant reproduction GRNs

An important aspect of understanding GRNs is how perturbations in one part of the network are transmitted to other parts of the network, and ultimately how this results in changes in flowering time. The modern version of *Arabidopsis* GRNs as presented above involves highly-connected, non-linear networks. Given this complexity, it is not possible to predict the effect of gene perturbations on e.g. flowering time in an intuitive way. Therefore, mathematical modelling plays an important role in providing a quantitative understanding of GRNs.

One of the pioneering models for cell-fate determination during the formation of floral organ primordia in *Arabidopsis* provided insights into the semi-quantitative relationships between the genes in the reproduction GRN (Espinosa-Soto *et al.*, 2004). Based on published data, regulatory relationship between 15 ABC and key non-ABC genes were translated into a discrete Boolean network model. The state of each gene is updated according to the states of the genes that directly regulate it, via a set of logical rules derived from a survey of molecular genetic experimental data. Model simulations for all possible starting states showed that the network converges to a few steady states that correspond to expression patterns observed in each of the primordial cell types (i.e. inflorescence, sepals, petals, stamens, or carpels) and are in agreement with the phenotypes predicted by the ABC genetic model for both wild type and mutants. Analysis of the simulation results and the logical rules derived from published data led the authors to speculate that *AG* is involved in a positive feedback loop to maintain its own expression; this prediction was experimentally confirmed by a later study (Gomez-Mena *et al.*, 2005).

A quantitative model that captures not only the regulatory relationships but also the kinetics of MADS domain complex formation was later proposed to represent a GRN for organ-fate determination in *Arabidopsis* (van Mourik *et al.*, 2010). For this work, transcriptional regulation of six genes representing the five gene classes in the ABCDE model were

modelled using ordinary differential equations (ODEs). For each gene, there is an ODE describing how the gene expression level is influenced by the concentrations of its regulators. Based on gene expression data, whorl-specific concentrations were estimated, which were then used to estimate the various model parameters. These parameters describe the interaction affinity of the various MADS-domain protein dimers, the binding affinity of these dimers to target promoters and the decay of gene products into non-functional components. The model generates continuous time-course expressions for the involved genes in the different floral whorls that reasonably match experimental data. It has to be noted that such a model provides more detailed information about the network's dynamics than the pioneering discrete-network model (Espinosa-Soto *et al.*, 2004), but contains many more parameters that have to be fitted using experimental data.

Kaufmann *et al.* (2011) presented a model based on Petri nets that simulates floral quartet concentration dynamics. This model showed that complex formation attenuates stochastic fluctuations in gene expression thus enabling more robust organ-specific expression patterns.

With respect to the modelling of GRNs involved in flowering time control, some of the modelling approaches mentioned above, such as Boolean networks, in which time is not explicitly present, are less suitable. An ODE approach can serve as a framework for modelling the GRN underlying flowering time control. In particular, the control of flowering by photoperiod depends on an integration of external signals captured by photoreceptors and endogenous rhythm controlled by the circadian clock (Hayama and Coupland, 2003; Song *et al.*, 2013). This makes mathematical models for the *Arabidopsis* clock gene circuit, as recently reviewed, very relevant (Bujdoso and Davis, 2013). One important aspect is the regulation of *FT*, which is a major direct target of the clock-regulated gene *CO* (Yanovsky and Kay, 2002). Salazar *et al.* (2009) used an ODE-based modelling to reconstruct the rhythmic regulation of *CO* by the circadian clock, and the subsequent effect of *CO* expression on the regulation of *FT*. To account for the effect of photoperiod on gene expression, the model assumed an explicit role of post-translational regulation: *CO* protein is stabilized during the daytime, but rapidly degrades during the night. Thus only the peak of *CO* mRNA that occurs in the light leads to *CO* protein accumulation and, therefore, *FT* activation. Interestingly, the expression levels of *FT* were simulated for different photoperiod cycles (light/dark), which indicated a non-linear relationship between observed flowering time and the amount of *FT* transcribed over a cycle-period. In the leaves, *FT* is not only regulated by the photoperiod sensors but also by temperature-related cues (Song *et al.*, 2013) via an *FLC*-mediated mechanism. Recently, the regulatory relationship between *FLC* and *FT* under the influence of temperature and photoperiod was modelled for the perennial *Arabidopsis halleri* (Satake *et al.*, 2013). Explicit information about temperature and photoperiod was used to simulate ODEs describing gene expression levels. Assessment of gene expression simulated for different temperatures allowed changes in the perennial flowering cycle to be forecast under a climate change scenario.

Finally, a recent ODE model for the transcriptional regulation of five key integrators of flowering time (Jaeger *et al.*, 2013) was used to explore mechanistically how different feedback loops affect flowering time.

These examples show that GRN modelling enables various qualitative and quantitative predictions on network output and demonstrates the importance of such modelling approaches in understanding how the complex GRNs in plant reproduction fulfil their function.

## Conclusion and perspective

The introduction of next-generation sequencing and genome-wide approaches has changed our view of gene regulation and GRNs. We have moved from linear genetic interactions towards global highly connected gene networks. Many genome-wide expression compendia have become available as well as protein–DNA binding profile data that rapidly increased our knowledge of transcriptional regulation and network wiring. Information on epigenetic regulation of gene expression remains poorly represented in the current GRN models and long-distance signalling also needs further consideration. Another complexity concerns spatial and temporal determinants. We know that genes can have different functions, depending on the tissue and stage at which they are expressed. For example, AP1 acts as a repressor of flowering time genes during flower initiation, whereas at a later stages it activates genes involved in organ formation. Information about tissue and stage-specific gene expression (Jiao and Meyerowitz, 2010; Wellmer *et al.*, 2006) and protein–DNA binding (Pajoro *et al.*, 2014) will generate a better understanding of gene function and network dynamics. Most likely, more tissue and stage-specific information about protein–DNA binding will become available in the near future and will allow reconstruction of stage-specific GRNs, highlighting the dynamics of the network topology and interactions. To achieve this resolution, novel technologies such as the INTACT system (Deal and Henikoff, 2011) and single cell approaches (Shapiro *et al.*, 2013) will be important.

Recent studies of TF DNA-binding profiles and gene expression analyses have shown that there is only a weak correlation between binding of a TF and changes in expression of its target genes (O'Maoileidigh *et al.*, 2014). An explanation could be that multiple TF binding events or co-factors are needed for gene regulation. In such a scenario only a specific combination of TFs binding will trigger changes in expression. Sequential ChIP analysis (Oh *et al.*, 2012) could be used to identify TF co-occupancy and obtain a better insight into the regulation of gene expression. Another explanation could be that a single binding event to a *cis*-regulatory element is not sufficient to drive expression, whereas binding of a TF to multiple sites, allowing a conformational change of the DNA, is needed to regulate gene expression. For instance, conformational changes are triggered by the binding of STK to multiple sites in the promoter of its target gene *VERDANDI* (Mendes *et al.*, 2013).

New techniques, such as chromatin capture (Stadhouders *et al.*, 2013) and ChIA-Pet (Zhang *et al.*, 2012) (Fullwood *et al.*, 2009), can be used to characterise *cis*-regulatory element interactions and their role in gene regulation.

The use of mathematical modelling will facilitate a better insight into GRN complexity, allowing the consequences of network perturbations to be predicted. Current models are primarily based on gene expression profiles and information about non-coding RNAs, protein–DNA, and protein–protein interactions is increasing and becoming incorporated into GRNs. Nevertheless, the experimentally validated protein–protein interactome is still quite sparse (Chatr-Aryamontri *et al.*, 2013; Smaczniak *et al.*, 2012b) and therefore we still rely on predicting the *Arabidopsis* interactome using orthology relationships, gene ontology, and co-expression (De Bodt *et al.*, 2009; Lee *et al.*, 2010; Lin *et al.*, 2011).

In conclusion, advances in genome-wide studies resulted in the confirmation of previously known genetic interactions at the molecular level, but also identified many novel regulatory interactions. We are just at the beginning of the genome-wide characterization of GRNs and new factors and interactions are undoubtedly waiting to be discovered. A challenge for the near future will be to unravel the spatial and temporal regulation of the genes in the current networks. This will allow further development and application of mathematical models, which will contribute to our understanding of the role and importance of genes during plant reproduction.

## Acknowledgements

The authors wish to thank the Marie-Curie Actions for support via the Marie-Curie-ITN network grant SYSFLO (FP7/2007–2011, grant agreement no. 237909).

## References

- Adrian J, Farrona S, Reimer JJ, Albani MC, Coupland G, Turck F.** 2010. *cis*-Regulatory elements and chromatin state coordinately control temporal and spatial expression of *FLOWERING LOCUS T* in *Arabidopsis*. *The Plant Cell* **22**, 1425–1440.
- Alvarez-Buylla ER, Azpeitia E, Barrio R, Benitez M, Padilla-Longoria P.** 2010. From ABC genes to regulatory networks, epigenetic landscapes and flower morphogenesis: making biological sense of theoretical approaches. *Seminars in Cell & Developmental Biology* **21**, 108–117.
- Aukerman MJ, Sakai H.** 2003. Regulation of flowering time and floral organ identity by a MicroRNA and its *APETALA2*-like target genes. *The Plant Cell* **15**, 2730–2741.
- Baurle I, Dean C.** 2006. The timing of developmental transitions in plants. *Cell* **125**, 655–664.
- Blazquez MA, Weigel D.** 2000. Integration of floral inductive signals in *Arabidopsis*. *Nature* **404**, 889–892.
- Blazquez MA, Ferrandiz C, Madueno F, Parcy F.** 2006. How floral meristems are built. *Plant Molecular Biology* **60**, 855–870.
- Blazquez MA, Green R, Nilsson O, Sussman MR, Weigel D.** 1998. Gibberellins promote flowering of *Arabidopsis* by activating the LEAFY promoter. *The Plant Cell* **10**, 791–800.
- Bujdoso N, Davis SJ.** 2013. Mathematical modeling of an oscillating gene circuit to unravel the circadian clock network of *Arabidopsis thaliana*. *Frontiers in Plant Science* **4**, 3.
- Cardon G, Hohmann S, Klein J, Nettesheim K, Saedler H, Huijser P.** 1999. Molecular characterisation of the *Arabidopsis* SBP-box genes. *Gene* **237**, 91–104.

- Causier B, Schwarz-Sommer Z, Davies B.** 2010. Floral organ identity: 20 years of ABCs. *Seminars in Cell & Developmental Biology* **21**, 73–79.
- Chatr-Aryamontri A, Breitkreutz BJ, Heinicke S, et al.** 2013. The BioGRID interaction database: 2013 update. *Nucleic Acids Research* **41**, D816–823.
- Corbesier L, Vincent C, Jang S, et al.** 2007. FT protein movement contributes to long-distance signaling in floral induction of *Arabidopsis*. *Science* **316**, 1030–1033.
- De Bodt S, Proost S, Vandepoele K, Rouze P, Van de Peer Y.** 2009. Predicting protein–protein interactions in *Arabidopsis thaliana* through integration of orthology, gene ontology and co-expression. *BMC Genomics* **10**, 288.
- de Folter S, Immink RG, Kieffer M, et al.** 2005. Comprehensive interaction map of the *Arabidopsis* MADS Box transcription factors. *The Plant Cell* **17**, 1424–1433.
- Deal RB, Henikoff S.** 2011. The INTACT method for cell type-specific gene expression and chromatin profiling in *Arabidopsis thaliana*. *Nature Protocols* **6**, 56–68.
- Deng W, Ying H, Hellwell CA, Taylor JM, Peacock WJ, Dennis ES.** 2011. *FLOWERING LOCUS C (FLC)* regulates development pathways throughout the life cycle of *Arabidopsis*. *Proceedings of the National Academy of Science, USA* **108**, 6680–6685.
- Dill A, Thomas SG, Hu J, Steber CM, Sun TP.** 2004. The *Arabidopsis* F-box protein SLEEPY1 targets gibberellin signaling repressors for gibberellin-induced degradation. *The Plant Cell* **16**, 1392–1405.
- Ditta G, Pinyopich A, Robles P, Pelaz S, Yanofsky MF.** 2004. The *SEP4* gene of *Arabidopsis thaliana* functions in floral organ and meristem identity. *Current Biology* **14**, 1935–1940.
- Egea-Cortines M, Saedler H, Sommer H.** 1999. Ternary complex formation between the MADS-box proteins SQUAMOSA, DEFICIENS and GLOBOSA is involved in the control of floral architecture in *Antirrhinum majus*. *The EMBO Journal* **18**, 5370–5379.
- Eriksson S, Bohlenius H, Moritz T, Nilsson O.** 2006. GA4 is the active gibberellin in the regulation of *LEAFY* transcription and *Arabidopsis* floral initiation. *The Plant Cell* **18**, 2172–2181.
- Espinosa-Soto C, Padilla-Longoria P, Alvarez-Buylla ER.** 2004. A gene regulatory network model for cell-fate determination during *Arabidopsis thaliana* flower development that is robust and recovers experimental gene expression profiles. *The Plant Cell* **16**, 2923–2939.
- Fullwood MJ, Liu MH, Pan YF, et al.** 2009. An oestrogen-receptor-alpha-bound human chromatin interactome. *Nature* **462**, 58–64.
- Galvao VC, Horrer D, Kuttner F, Schmid M.** 2012. Spatial control of flowering by DELLA proteins in *Arabidopsis thaliana*. *Development* **139**, 4072–4082.
- Gil P, Dewey E, Friml J, Zhao Y, Snowden KC, Putterill J, Palme K, Estelle M, Chory J.** 2001. BIG: a calossin-like protein required for polar auxin transport in *Arabidopsis*. *Genes & Development* **15**, 1985–1997.
- Gomez-Mena C, de Folter S, Costa MM, Angenent GC, Sablowski R.** 2005. Transcriptional program controlled by the floral homeotic gene AGAMOUS during early organogenesis. *Development* **132**, 429–438.
- Goto K, Meyerowitz EM.** 1994. Function and regulation of the *Arabidopsis* floral homeotic gene *PISTILLATA*. *Genes & Development* **8**, 1548–1560.
- Grandi V, Gregis V, Kater MM.** 2012. Uncovering genetic and molecular interactions among floral meristem identity genes in *Arabidopsis thaliana*. *The Plant Journal* **69**, 881–893.
- Gregis V, Andres F, Sessa A, et al.** 2013. Identification of pathways directly regulated by SHORT VEGETATIVE PHASE during vegetative and reproductive development in *Arabidopsis*. *Genome Biology* **14**, R56.
- Gregis V, Sessa A, Colombo L, Kater MM.** 2006. AGL24, SHORT VEGETATIVE PHASE, and APETALA1 redundantly control AGAMOUS during early stages of flower development in *Arabidopsis*. *The Plant Cell* **18**, 1373–1382.
- Gregis V, Sessa A, Dorca-Fornell C, Kater MM.** 2009. The *Arabidopsis* floral meristem identity genes *AP1*, *AGL24* and *SVP* directly repress class B and C floral homeotic genes. *The Plant Journal* **60**, 626–637.
- Hawkins RD, Hon GC, Ren B.** 2010. Next-generation genomics: an integrative approach. *Nature Reviews Genetics* **11**, 476–486.
- Hayama R, Coupland G.** 2003. Shedding light on the circadian clock and the photoperiodic control of flowering. *Current Opinion in Plant Biology* **6**, 13–19.
- Honma T, Goto K.** 2001. Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* **409**, 525–529.
- Immink RG, Kaufmann K, Angenent GC.** 2010. The ‘ABC’ of MADS domain protein behaviour and interactions. *Seminars in Cell & Developmental Biology* **21**, 87–93.
- Immink RG, Tonaco IA, de Folter S, Shchennikova A, van Dijk AD, Busscher-Lange J, Borst JW, Angenent GC.** 2009. SEPALLATA3: the ‘glue’ for MADS box transcription factor complex formation. *Genome Biology* **10**, R24.
- Immink RG, Pose D, Ferrario S, et al.** 2012. Characterization of SOC1’s central role in flowering by the identification of its upstream and downstream regulators. *Plant Physiology* **160**, 433–449.
- Jack T, Brockman LL, Meyerowitz EM.** 1992. The homeotic gene *APETALA3* of *Arabidopsis thaliana* encodes a MADS box and is expressed in petals and stamens. *Cell* **68**, 683–697.
- Jaeger KE, Wigge PA.** 2007. FT protein acts as a long-range signal in *Arabidopsis*. *Current Biology* **17**, 1050–1054.
- Jaeger KE, Pullen N, Lamzin S, Morris RJ, Wigge PA.** 2013. Interlocking feedback loops govern the dynamic behavior of the floral transition in *Arabidopsis*. *The Plant Cell* **25**, 820–833.
- Jang S, Torti S, Coupland G.** 2009. Genetic and spatial interactions between FT, TSF and SVP during the early stages of floral induction in *Arabidopsis*. *The Plant Journal* **60**, 614–625.
- Jiao Y, Meyerowitz EM.** 2010. Cell-type specific analysis of translating RNAs in developing flowers reveals new levels of control. *Molecular Systems Biology* **6**, 419.
- Jofuku KD, den Boer BG, Van Montagu M, Okamuro JK.** 1994. Control of *Arabidopsis* flower and seed development by the homeotic gene *APETALA2*. *The Plant Cell* **6**, 1211–1225.
- Jung JH, Ju Y, Seo PJ, Lee JH, Park CM.** 2012. The SOC1-SPL module integrates photoperiod and gibberellin acid signals to control flowering time in *Arabidopsis*. *The Plant Journal* **69**, 577–588.
- Jung JH, Seo YH, Seo PJ, Reyes JL, Yun J, Chua NH, Park CM.** 2007. The *GIGANTEA*-regulated microRNA172 mediates photoperiodic flowering independent of CONSTANS in *Arabidopsis*. *The Plant Cell* **19**, 2736–2748.
- Kanno Y, Jikumaru Y, Hanada A, Nambara E, Abrams SR, Kamiya Y, Seo M.** 2010. Comprehensive hormone profiling in developing *Arabidopsis* seeds: examination of the site of ABA biosynthesis, ABA transport and hormone interactions. *Plant and Cell Physiology* **51**, 1988–2001.
- Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harrison MJ, Weigel D.** 1999. Activation tagging of the floral inducer FT. *Science* **286**, 1962–1965.
- Kaufmann K, Pajoro A, Angenent GC.** 2010a. Regulation of transcription in plants: mechanisms controlling developmental switches. *Nature Reviews Genetics* **11**, 830–842.
- Kaufmann K, Nagasaki M, Jauregui R.** 2011. Modelling the molecular interactions in the flower developmental network of *Arabidopsis thaliana*. *Studies in Health Technology and Informatics* **162**, 279–297.
- Kaufmann K, Muino JM, Osteras M, Farinelli L, Krajewski P, Angenent GC.** 2010b. Chromatin immunoprecipitation (ChIP) of plant transcription factors followed by sequencing (ChIP-SEQ) or hybridization to whole genome arrays (ChIP-ChIP). *Nature Protocols* **5**, 457–472.
- Kaufmann K, Muino JM, Jauregui R, Aioldi CA, Smaczniak C, Krajewski P, Angenent GC.** 2009. Target genes of the MADS transcription factor SEPALLATA3: integration of developmental and hormonal pathways in the *Arabidopsis* flower. *PLoS Biology* **7**, e1000090.
- Kaufmann K, Wellmer F, Muino JM, et al.** 2010c. Orchestration of floral initiation by APETALA1. *Science* **328**, 85–89.
- Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T.** 1999. A pair of related genes with antagonistic roles in mediating flowering signals. *Science* **286**, 1960–1962.

- Koornneef M, Hanhart CJ, van der Veen JH.** 1991. A genetic and physiological analysis of late flowering mutants in *Arabidopsis thaliana*. *Molecular and General Genetics* **229**, 57–66.
- Lee I, Ambaru B, Thakkar P, Marcotte EM, Rhee SY.** 2010. Rational association of genes with traits using a genome-scale gene network for *Arabidopsis thaliana*. *Nature Biotechnology* **28**, 149–156.
- Lee JH, Yoo SJ, Park SH, Hwang I, Lee JS, Ahn JH.** 2007. Role of SVP in the control of flowering time by ambient temperature in *Arabidopsis*. *Genes & Development* **21**, 397–402.
- Li D, Liu C, Shen L, Wu Y, Chen H, Robertson M, Hellwell CA, Ito T, Meyerowitz E, Yu H.** 2008. A repressor complex governs the integration of flowering signals in *Arabidopsis*. *Developmental Cell* **15**, 110–120.
- Lin M, Shen X, Chen X.** 2011. PAIR: the predicted *Arabidopsis* interactome resource. *Nucleic Acids Research* **39**, D1134–1140.
- Liu C, Chen H, Er HL, Soo HM, Kumar PP, Han JH, Liou YC, Yu H.** 2008. Direct interaction of AGL24 and SOC1 integrates flowering signals in *Arabidopsis*. *Development* **135**, 1481–1491.
- Machanick P, Bailey TL.** 2011. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics* **27**, 1696–1697.
- Mandel MA, Yanofsky MF.** 1995. A gene triggering flower formation in *Arabidopsis*. *Nature* **377**, 522–524.
- Mathieu J, Warthmann N, Kuttner F, Schmid M.** 2007. Export of FT protein from phloem companion cells is sufficient for floral induction in *Arabidopsis*. *Current Biology* **17**, 1055–1060.
- Mathieu J, Yant LJ, Murdter F, Kuttner F, Schmid M.** 2009. Repression of flowering by the miR172 target SMZ. *PLoS Biology* **7**, e1000148.
- Melzer R, Theissen G.** 2009. Reconstitution of ‘floral quartets’ in vitro involving class B and class E floral homeotic proteins. *Nucleic Acids Research* **37**, 2723–2736.
- Mendes MA, Guerra RF, Berns MC, Manzo C, Masiero S, Finzi L, Kater MM, Colombo L.** 2013. MADS domain transcription factors mediate short-range DNA looping that is essential for target gene expression in *Arabidopsis*. *The Plant Cell* **25**, 2560–2572.
- Michaels SD, Ditta GS, Gustafson-Brown C, Pelaz S, Yanofsky M, Amasino RM.** 2003. AGL24 acts as a promoter of flowering in *Arabidopsis* and is positively regulated by vernalization. *The Plant Journal* **33**, 867–874.
- Moon J, Suh SS, Lee H, Choi KR, Hong CB, Paek NC, Kim SG, Lee I.** 2003. The SOC1 MADS-box gene integrates vernalization and gibberellin signals for flowering in *Arabidopsis*. *The Plant Journal* **35**, 613–623.
- Moyroud E, Minguez EG, Ott F, et al.** 2011. Prediction of regulatory interactions from genome sequences using a biophysical model for the *Arabidopsis* LEAFY transcription factor. *The Plant Cell* **23**, 1293–1306.
- O’Maoileidigh DS, Graciet E, Wellmer F.** 2014. Gene networks controlling *Arabidopsis thaliana* flower development. *New Phytologist* **201**, 16–30.
- O’Maoileidigh DS, Wuest SE, Rae L, et al.** 2013. Control of reproductive floral organ identity specification in *Arabidopsis* by the C function regulator AGAMOUS. *The Plant Cell* **25**, 2482–2503.
- Oh E, Zhu JY, Wang ZY.** 2012. Interaction between BZR1 and PIF4 integrates brassinosteroid and environmental responses. *Nature Cell Biology* **14**, 802–809.
- Okamuro JK, Caster B, Villarroel R, Van Montagu M, Jofuku KD.** 1997. The AP2 domain of APETALA2 defines a large new family of DNA binding proteins in *Arabidopsis*. *Proceedings of the National Academy of Science, USA* **94**, 7076–7081.
- Pajoro A, Madrigal P, Muino JM, et al.** 2014. Dynamics of chromatin accessibility and gene regulation by MADS-domain transcription factors in flower development. *Genome Biology* **15**, R41.
- Parenicova L, de Folter S, Kieffer M, et al.** 2003. Molecular and phylogenetic analyses of the complete MADS-box transcription factor family in *Arabidopsis*: new openings to the MADS world. *The Plant Cell* **15**, 1538–1551.
- Pastore JJ, Limpuangthip A, Yamaguchi N, Wu MF, Sang Y, Han SK, Malaspina L, Chavdaroff N, Yamaguchi A, Wagner D.** 2011. LATE MERISTEM IDENTITY2 acts together with LEAFY to activate APETALA1. *Development* **138**, 3189–3198.
- Pelaz S, Ditta GS, Baumann E, Wisman E, Yanofsky MF.** 2000. B and C floral organ identity functions require SEPALLATA MADS-box genes. *Nature* **405**, 200–203.
- Pelaz S, Gustafson-Brown C, Kohalmi SE, Crosby WL, Yanofsky MF.** 2001. APETALA1 and SEPALLATA3 interact to promote flower development. *The Plant Journal* **26**, 385–394.
- Pinyopich A, Ditta GS, Savidge B, Liljegren SJ, Baumann E, Wisman E, Yanofsky MF.** 2003. Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature* **424**, 85–88.
- Porri A, Torti S, Romera-Branchat M, Coupland G.** 2012. Spatially distinct regulatory roles for gibberellins in the promotion of flowering of *Arabidopsis* under long photoperiods. *Development* **139**, 2198–2209.
- Pose D, Yant L, Schmid M.** 2012. The end of innocence: flowering networks explode in complexity. *Current Opinion in Plant Biology* **15**, 45–50.
- Pose D, Verhage L, Ott F, Yant L, Mathieu J, Angenent GC, Immink RG, Schmid M.** 2013. Temperature-dependent regulation of flowering by antagonistic FLM variants. *Nature* **503**, 414–417.
- Ragni L, Nieminen K, Pacheco-Villalobos D, Sibout R, Schwechheimer C, Hardtke CS.** 2011. Mobile gibberellin directly stimulates *Arabidopsis* hypocotyl xylem expansion. *The Plant Cell* **23**, 1322–1336.
- Reiser L, Fischer RL.** 1993. The Ovule and the embryo sac. *The Plant Cell* **5**, 1291–1301.
- Richter R, Behringer C, Zourelidou M, Schwechheimer C.** 2013. Convergence of auxin and gibberellin signaling on the regulation of the GATA transcription factors GNC and GNL in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **110**, 13192–13197.
- Rounseley SD, Ditta GS, Yanofsky MF.** 1995. Diverse roles for MADS box genes in *Arabidopsis* development. *The Plant Cell* **7**, 1259–1269.
- Salazar JD, Saithong T, Brown PE, Foreman J, Locke JC, Halliday KJ, Carre IA, Rand DA, Millar AJ.** 2009. Prediction of photoperiodic regulators from quantitative gene circuit models. *Cell* **139**, 1170–1179.
- Samach A, Onouchi H, Gold SE, Ditta GS, Schwarz-Sommer Z, Yanofsky MF, Coupland G.** 2000. Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*. *Science* **288**, 1613–1616.
- Satake A, Kawagoe T, Saburi Y, Chiba Y, Sakurai G, Kudoh H.** 2013. Forecasting flowering phenology under climate warming by modelling the regulatory dynamics of flowering-time genes. *Nature Communications* **4**, 2303.
- Sawa M, Kay SA.** 2011. GIGANTEA directly activates Flowering Locus T in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences, USA* **108**, 11698–11703.
- Schaffer R, Ramsay N, Samach A, Corden S, Putterill J, Carre IA, Coupland G.** 1998. The late elongated hypocotyl mutation of *Arabidopsis* disrupts circadian rhythms and the photoperiodic control of flowering. *Cell* **93**, 1219–1229.
- Schwarz S, Grande AV, Bujdoso N, Saedler H, Huijser P.** 2008. The microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in *Arabidopsis*. *Plant Molecular Biology* **67**, 183–195.
- Seo E, Lee H, Jeon J, Park H, Kim J, Noh YS, Lee I.** 2009. Crosstalk between cold response and flowering in *Arabidopsis* is mediated through the flowering-time gene SOC1 and its upstream negative regulator FLC. *The Plant Cell* **21**, 3185–3197.
- Sessions A, Nemhauser JL, McColl A, Roe JL, Feldmann KA, Zambryski PC.** 1997. ETTIN patterns the *Arabidopsis* floral meristem and reproductive organs. *Development* **124**, 4481–4491.
- Shan H, Chen S, Jiang J, et al.** 2012. Heterologous expression of the chrysanthemum R2R3-MYB transcription factor CmMYB2 enhances drought and salinity tolerance, increases hypersensitivity to ABA and delays flowering in *Arabidopsis thaliana*. *Molecular Biotechnology* **51**, 160–173.
- Shani E, Weinstain R, Zhang Y, Castillejo C, Kaiserli E, Chory J, Tsien RY, Estelle M.** 2013. Gibberellins accumulate in the elongating endodermal cells of *Arabidopsis* root. *Proceedings of the National Academy of Sciences, USA* **110**, 4834–4839.

- Shapiro E, Biezuner T, Linnarsson S.** 2013. Single-cell sequencing-based technologies will revolutionize whole-organism science. *Nature Reviews Genetics* **14**, 618–630.
- Shi DQ, Yang WC.** 2011. Ovule development in *Arabidopsis*: progress and challenge. *Current Opinion in Plant Biology* **14**, 74–80.
- Simpson GG, Dean C.** 2002. *Arabidopsis*, the Rosetta stone of flowering time? *Science* **296**, 285–289.
- Smaczniak C, Immink RG, Angenent GC, Kaufmann K.** 2012a. Developmental and evolutionary diversity of plant MADS-domain factors: insights from recent studies. *Development* **139**, 3081–3098.
- Smaczniak C, Immink RG, Muino JM, et al.** 2012b. Characterization of MADS-domain transcription factor complexes in *Arabidopsis* flower development. *Proceedings of the National Academy of Sciences, USA* **109**, 1560–1565.
- Song YH, Ito S, Imaizumi T.** 2013. Flowering time regulation: photoperiod- and temperature-sensing in leaves. *Trends in Plant Science* **18**, 575–583.
- Stadhouders R, Kolovos P, Brouwer R, et al.** 2013. Multiplexed chromosome conformation capture sequencing for rapid genome-scale high-resolution detection of long-range chromatin interactions. *Nature Protocols* **8**, 509–524.
- Suarez-Lopez P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G.** 2001. CONSTANS mediates between the circadian clock and the control of flowering in *Arabidopsis*. *Nature* **410**, 1116–1120.
- Takada S, Goto K.** 2003. TERMINAL FLOWER2, an *Arabidopsis* homolog of HETEROCHROMATIN PROTEIN1, counteracts the activation of FLOWERING LOCUS T by CONSTANS in the vascular tissues of leaves to regulate flowering time. *The Plant Cell* **15**, 2856–2865.
- Tao Z, Shen L, Liu C, Liu L, Yan Y, Yu H.** 2012. Genome-wide identification of SOC1 and SVP targets during the floral transition in *Arabidopsis*. *The Plant Journal* **70**, 549–561.
- Torti S, Fornara F, Vincent C, Andres F, Nordstrom K, Gobel U, Knoll D, Schoof H, Coupland G.** 2012. Analysis of the *Arabidopsis* shoot meristem transcriptome during floral transition identifies distinct regulatory patterns and a leucine-rich repeat protein that promotes flowering. *The Plant Cell* **24**, 444–462.
- Tröbner W, Ramirez L, Motte P, Hue I, Huijser P, Lonnig WE, Saedler H, Sommer H, Schwarz-Sommer Z.** 1992. GLOBOSA: a homeotic gene which interacts with DEFICIENS in the control of *Antirrhinum* floral organogenesis. *The EMBO Journal* **11**, 4693–4704.
- van Mourik S, van Dijk AD, de Gee M, Immink RG, Kaufmann K, Angenent GC, van Ham RC, Molenaar J.** 2010. Continuous-time modeling of cell fate determination in *Arabidopsis* flowers. *BMC Systems Biology* **4**, 101.
- Wang H, Tang W, Zhu C, Perry SE.** 2002. A chromatin immunoprecipitation (ChIP) approach to isolate genes regulated by AGL15, a MADS domain protein that preferentially accumulates in embryos. *The Plant Journal* **32**, 831–843.
- Wang JW, Czech B, Weigel D.** 2009. miR156-regulated SPL transcription factors define an endogenous flowering pathway in *Arabidopsis thaliana*. *Cell* **138**, 738–749.
- Wellmer F, Alves-Ferreira M, Dubois A, Riechmann JL, Meyerowitz EM.** 2006. Genome-wide analysis of gene expression during early *Arabidopsis* flower development. *PLoS Genetics* **2**, e117.
- Willige BC, Ghosh S, Nill C, Zourelidou M, Dohmann EM, Maier A, Schwechheimer C.** 2007. The DELLA domain of GA INSENSITIVE mediates the interaction with the GA INSENSITIVE DWARF1A gibberellin receptor of *Arabidopsis*. *The Plant Cell* **19**, 1209–1220.
- Winter CM, Austin RS, Blanvillain-Baufume S, et al.** 2011. LEAFY target genes reveal floral regulatory logic, cis motifs, and a link to biotic stimulus response. *Developmental Cell* **20**, 430–443.
- Wu G, Poethig RS.** 2006. Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target SPL3. *Development* **133**, 3539–3547.
- Wu G, Park MY, Conway SR, Wang JW, Weigel D, Poethig RS.** 2009. The sequential action of miR156 and miR172 regulates developmental timing in *Arabidopsis*. *Cell* **138**, 750–759.
- Wu S, Gallagher KL.** 2012. Transcription factors on the move. *Current Opinion in Plant Biology* **15**, 645–651.
- Wuest SE, O'Maileidigh DS, Rae L, Kwasniewska K, Raganelli A, Hanczaryk K, Lohan AJ, Loftus B, Graciet E, Wellmer F.** 2012. Molecular basis for the specification of floral organs by APETALA3 and PISTILLATA. *Proceedings of the National Academy of Sciences, USA* **109**, 13452–13457.
- Yamaguchi A, Kobayashi Y, Goto K, Abe M, Araki T.** 2005. TWIN SISTER OF FT (TSF) acts as a floral pathway integrator redundantly with FT. *Plant and Cell Physiology* **46**, 1175–1189.
- Yamaguchi A, Wu MF, Yang L, Wu G, Poethig RS, Wagner D.** 2009. The microRNA-regulated SBP-Box transcription factor SPL3 is a direct upstream activator of LEAFY, FRUITFULL, and APETALA1. *Developmental Cell* **17**, 268–278.
- Yamaguchi N, Suzuki M, Fukaki H, Morita-Terao M, Tasaka M, Komeda Y.** 2007. CRM1/BIG-mediated auxin action regulates *Arabidopsis* inflorescence development. *Plant and Cell Physiology* **48**, 1275–1290.
- Yanofsky MF, Ma H, Bowman JL, Drews GN, Feldmann KA, Meyerowitz EM.** 1990. The protein encoded by the *Arabidopsis* homeotic gene agamous resembles transcription factors. *Nature* **346**, 35–39.
- Yanovsky MJ, Kay SA.** 2002. Molecular basis of seasonal time measurement in *Arabidopsis*. *Nature* **419**, 308–312.
- Yant L, Mathieu J, Dinh TT, Ott F, Lanz C, Wollmann H, Chen X, Schmid M.** 2010. Orchestration of the floral transition and floral development in *Arabidopsis* by the bifunctional transcription factor APETALA2. *The Plant Cell* **22**, 2156–2170.
- Yu H, Xu Y, Tan EL, Kumar PP.** 2002. AGAMOUS-LIKE 24, a dosage-dependent mediator of the flowering signals. *Proceedings of the National Academy of Science, USA* **99**, 16336–16341.
- Yu S, Galvao VC, Zhang YC, Horrer D, Zhang TQ, Hao YH, Feng YQ, Wang S, Schmid M, Wang JW.** 2012. Gibberellin regulates the *Arabidopsis* floral transition through miR156-targeted SQUAMOSA promoter binding-like transcription factors. *The Plant Cell* **24**, 3320–3332.
- Zhang J, Poh HM, Peh SQ, et al.** 2012. ChIA-PET analysis of transcriptional chromatin interactions. *Methods* **58**, 289–299.
- Zheng Y, Ren N, Wang H, Stromberg AJ, Perry SE.** 2009. Global identification of targets of the *Arabidopsis* MADS domain protein AGAMOUS-Like15. *The Plant Cell* **21**, 2563–2577.