

^1H , ^{13}C and ^{15}N resonance assignments of the Calmodulin-Munc13-1 peptide complex

Fernando Rodríguez-Castañeda · Nicolas Coudeville · Stefan Becker · Nils Brose · Teresa Carlomagno · Christian Griesinger

Received: 5 October 2009 / Accepted: 2 December 2009
© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract Ca^{2+} -Calmodulin binding to the variable N-terminal region of the diacylglycerol/phorbol ester-binding UNC13/Munc13 family of proteins modulates the short-term synaptic plasticity characteristics in neurons. Here, we report the sequential backbone and side chain resonance assignment of the Ca^{2+} -Calmodulin/Munc13-1^{458–492} peptide complex at pH 6.8 and 35°C (BMRB No. 15470).

Keywords Calcium · Calmodulin · Munc13 · Neurotransmitter release · Synaptic plasticity · Phorbol esters and vesicle priming

Biological context

UNC13/Munc13 family of proteins are diacylglycerol/phorbol ester-binding proteins specifically localized to presynaptic

active zones, where the synaptic vesicles dock and release their neurotransmitters to the intersynaptic cleft for the chemical signal transmission among neurons (Junge et al. 2004; Brose et al. 1995). In *Caenorhabditis elegans*, *unc-13* is essential for coordinated movement (Brenner 1974). At a molecular level, UNC13/Munc13 proteins are essential in the maturation process of synaptic vesicles, remodeling the SNARE complex of proteins leading to fusion competent molecular states. Upon deletion of UNC13/Munc13 proteins neurotransmitter release is completely impaired (Richmond et al. 2001).

Ca^{2+} -Calmodulin (CaM) is a highly conserved ubiquitous eukaryotic signaling protein, regulating the activity of numerous proteins like protein kinases, phosphodiesterases, ion channels and pumps, and nitric-oxide synthases. Various studies have shown that CaM activates these proteins interacting with regulatory elements possessing an amphiphilic α -helix character (Crivici and Ikura 1995).

It has been shown (Junge et al. 2004) that some of the Munc13 protein isoforms contain a highly conserved CaM binding motif in their variable N-terminal region. However, this binding motif does not belong to any of the classical CaM binding motifs previously described (Rhoads and Friedberg 1997). Synthetic peptides of the Munc13-1 and ubMunc13-2 isoforms belonging to this CaM binding domain form high-affinity complexes with CaM in vitro. To unravel the CaM binding mechanism to Munc13 proteins, we carried out NMR experiments on the CaM/Munc13-1^{458–492} peptide complex.

Electronic supplementary material The online version of this article (doi:10.1007/s12104-009-9204-2) contains supplementary material, which is available to authorized users.

F. Rodríguez-Castañeda · N. Coudeville · S. Becker · T. Carlomagno · C. Griesinger (✉)
NMR-Based Structural Biology Department, Max Planck Institute for Biophysical Chemistry, Am Faßberg 11, 37077 Göttingen, Germany
e-mail: cigr@nmr.mpibpc.mpg.de

N. Brose
Molecular Neurobiology Department, Max Planck Institute for Experimental Medicine, Hermann-Rein-Str. 3, 37075 Göttingen, Germany

Present Address:
T. Carlomagno
European Molecular Biology Laboratory,
Meyerhofstrasse 1, 69117 Heidelberg, Germany

Methods and experiments

NMR sample preparation

The costs of ^{15}N , ^{13}C -uniformly labeled synthetic peptides are prohibitive. Therefore, we chose a recombinant approach

to obtain the Munc13-1^{458–492} peptide and will be described elsewhere. Briefly, the CaM binding domain (458–492) coding sequence of *munc13-1* was cloned into the pGEX-2T expression vector (*GE Healthcare*). The uniformly labeled (¹⁵N, ¹³C) GST-Munc13-1^{458–492} fusion protein was over-expressed in M9 minimal medium containing ¹⁵NH₄Cl and ¹³C-glucose as solely source of nitrogen and carbon, respectively. Due to solubility problems, the GST-Munc13-1^{458–492} fusion protein was co-expressed with CaM using resistance to two different antibiotics. The GST-Munc13-1^{458–492}/CaM complex was co-purified via affinity chromatography using GST-Sepharose (*GE-Healthcare*) according to the vendor instructions. The GST fusion was cleaved with thrombin and the Munc13-1 peptide purified by reverse-phase HPLC. The eluted peptide was lyophilized, reconstituted with 250 µl of the NMR-sample aqueous buffer (20 mM Bis-Tris, 150 mM KCl, 10 mM CaCl₂, pH 6.8) and mixed with a 1.2 M excess of unlabeled CaM dissolved in (*Millipore*) water to give a total final volume of 2 ml. The CaM/Munc13-1^{459–492} complex was concentrated by vacuum centrifugation in a *Speedvac* to a final volume of 250 µl giving a final concentration of approximately 0.5 mM. In order to get a lock signal in the NMR spectrometer, we added 10 µl of 99% D₂O.

The (U-¹⁵N, ¹³C) labeled CaM was obtained as described (Guerini et al. 1984; Haberbz et al. 2006). Briefly, CaM was over-expressed from a pET28a-CaM construct in *E. coli* BL21(DE3) in labeled M9 minimal medium. The cells were lysed by ultrasound sonication and the cell debris removed by centrifugation at 15,000 g for 45 min. The supernatant was precipitated with 2.84% trichloroacetic acid and subjected to hydrophobic interaction chromatography on a phenylsepharose column; CaM was eluted with an EGTA containing buffer. To prepare the complex sample, 8 mg of lyophilized (U-¹⁵N, ¹³C) labeled CaM were weighted and dissolved in 2 ml of (*Millipore*) water. Around 2.4 mg of a synthetic natural abundance Munc13-1^{459–492} peptide (1.2 equivalents) were weighted and dissolved in 250 µl of the NMR-sample aqueous buffer (20 mM Bis-Tris, 150 mM KCl, 10 mM CaCl₂, pH 6.8) and added to the CaM solution; the mixture was concentrated by vacuum centrifugation in a *Speedvac* to give a final volume of 250 µl. 10 µl of 99% D₂O were added for the NMR-lock signal. An additional (U-¹⁵N, ¹³C) labeled CaM/Munc13-1^{459–492} complex sample dissolved in 99% D₂O was prepared in the same manner. The three samples were placed in 280 µl *Shigemi* microcells covered with their glass plunger and sealed with *Parafilm*.

NMR experiments

All NMR experiments were carried out at 35°C on the following *Bruker* spectrometers: *Avance* 600 MHz

equipped with a z-gradient cryogenic HCN probe head; *DRX* 600 MHz with a z-gradient HCN probe head; *Avance* 700 MHz with a triple-axis gradient HCN probe head; *DMX* 800 MHz with a z-gradient cryogenic HCN probe head; and *Avance* 900 MHz with a z-gradient cryogenic HCN probe head.

The experiments made use of pulse field gradient coherence selection. Quadrature detection in the indirect dimensions was achieved either with the States-TPPI or Echo/antiecho methods. The chemical shift referencing was achieved internally by the addition of 0.4 mM DSS. The acquisition parameters for the experiments are given in Table 1. The spectra were processed with XWIN-NMR, Felix (*Accelrys, Inc.*), NMR-PIPE (Delaglio et al. 1995) and analyzed with SPARKY (Goddard and Kneller 1999).

Assignments and data deposition

The sequential backbone and side chain resonance assignment is nearly complete and had been deposited in the Biological Magnetic Resonance Data Bank (<http://www.bmrb.wisc.edu>) under the accession number 15,470. The ¹H-¹⁵N-HSQC spectrum of the (U-¹⁵N, ¹³C) CaM/Munc13-1^{459–492} complex sample is shown in Fig. 1. Its general appearance is consistent with that one of Ca²⁺-CaM, however, many NH cross-peaks in the N-terminal domain (1–76) of CaM had large line-widths mainly in helices A and D. This gave rise to an assignment completeness of 95.7% for the backbone resonances (CA, CB, C, N, H, HA and HB). In contrast, the assignment of the C-terminal domain (83–148) of the Ca²⁺-CaM/peptide complex was nearly complete (98.8% Backbone resonances). This suggested that the N-terminal domain of Ca²⁺-CaM experiences conformational exchange, whereas the C-terminal domain consists of a single conformer under these conditions. In Fig. 2, the ¹H-¹⁵N-HSQC spectrum of the (U-¹⁵N, ¹³C) Munc13-1^{458–492}/CaM complex is shown. An overlay of this spectrum with that one of a (U-¹⁵N, ¹³C)-Munc13-1^{458–492} peptide in 8 M urea solution (supplementary material) clearly revealed an increase in the amide resonance dispersion of the peptide, characteristic of secondary structure formation in this amphiphilic α-helix motif upon complex formation with Ca²⁺-CaM.

The assignment of the labeled Munc13-1^{458–492} peptide in complex with Ca²⁺-CaM was completed to 99.2% of the backbone resonances. The chemical shifts of the ε-methyl protons in the methionines of Ca²⁺-CaM in complex with the Munc13-1^{458–492} peptide were compared with those of free Ca²⁺-CaM. Most of the ε-methyl protons of the eight methionines in Ca²⁺-CaM were shifted upfield upon binding to the peptide. Significant changes (>0.15 ppm) were found for M51 (−0.3 ppm), M71 (−0.31 ppm), M109

Table 1 NMR data acquisition parameters

Sample	Experiment	Spectral frequency MHz	Complex points			
			D_1	D_2	D_3	
(U- ^{15}N , ^{13}C) CaM/Munc13-1 ^{459–492} in 90% H ₂ O, 10% D ₂ O	^1H - ^{15}N HSQC	900	2,048	128	–	
	^1H - ^{15}N NOESY-HSQC, t_{mix} 100 ms	900	2,048	48	128	
	HNCO	700	2,048	46	46	
	HN(CA)CO	600	1,024	60	60	
	HNCACB	700	1,024	40	25	
	CBCA(CO)NH	600	1,024	58	24	
	H(CC)(CO)NH-TOCSY	700	2,048	30	64	
	(H)CC(CO)NH-TOCSY	700	2,048	70	30	
	^1H - ^{13}C HSQC	600	2,048	128	–	
	(U- ^{15}N , ^{13}C) CaM/Munc13-1 ^{459–492} in 99% D ₂ O	^1H - ^{13}C NOESY-HSQC t_{mix} 120 ms	700	2,048	48	58
	HCCH-TOCSY	600	2,048	128	64	
^1H - ^{13}C HSQC	700	2,048	128	–		
^1H - ^{13}C NOESY-HSQC (aromatic region) t_{mix} 130 ms	700	2,048	48	40		
HBCB(Caro)HG	600	2,048	46	–		
HBCB(Caro)HE	600	2,048	46	–		
^1H - ^{13}C HMBC (methionines)	700	2,048	48	120		
(U- ^{15}N , ^{13}C) Munc13-1 ^{458–492} /CaM in 90% H ₂ O, 10% D ₂ O	^1H - ^{15}N HSQC	600	1,024	128	–	
	^1H - ^{15}N NOESY-HSQC and ^1H - ^{13}C NOESY-HSQC shared version, t_{mix} 120 ms	600	2,048	180	114	
	HNCO	900	1,024	64	32	
	HN(CA)CO	600	1,024	48	46	
	HNCACB	600	1,024	50	100	
	CBCA(CO)NH	600	1,024	29	60	
	H(CC)(CO)NH-TOCSY	600	1,024	26	50	
	(H)CC(CO)NH-TOCSY	600	1,024	30	52	
	HCCH-TOCSY	600	2,048	32	128	



Fig. 1 ^1H - ^{15}N -HSQC spectrum of the (U- ^{15}N , ^{13}C) CaM/Munc13-1^{459–492} complex measured at 35°C at 900 MHz. The highly overlapped central region is enlarged on the right side

Fig. 2 ^1H - ^{15}N -HSQC spectrum of the (U - ^{15}N , ^{13}C) Munc13-1^{458–492}/CaM complex recorded at 35°C at 600 MHz



(-0.17 ppm), M124 (-1.17 ppm), M144 (-0.3 ppm) and M145 (-0.65 ppm). The effect on the chemical shift for the methionines is attributed to deshielding effects caused by the aromatic rings of the various aromatic amino acids present in the peptide complexed to Ca^{2+} -CaM.

Acknowledgments The authors would like to thank Dr. Pierre Montaville for useful discussions in the protein preparation part; Volker Klaukien, Kerstin Overkamp and Gerhard Wolff for peptide synthesis and HPLC purification; and Dr. Vinesh Vijayan and Dr. Christophe Fères for assistance in the NMR-measurements. This work has been financed by the Max-Planck Society.

Ethical standards The experiments comply with the german current laws for ethical standards.

Conflict of interests The authors declare that they have no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Brenner S (1974) The genetics of *Caenorhabditis elegans*. Genetics 77:71–94
- Brose N, Hofmann K, Hata Y, Südhof TC (1995) Mammalian homologues of *Caenorhabditis elegans unc-13* gene define novel family of C_2 -domain proteins. J Biol Chem 270:25273–25280

- Crivici A, Ikura M (1995) Molecular and structural basis of target recognition by calmodulin. Annu Rev Biophys Biomol Struct 24:85–116
- Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A (1995) NMRPipe: a multidimensional spectral processing system based on UNIX pipes. J Biomol NMR 6:277–293
- Goddard TD, Kneller DG (1999) SPARKY 3. University of California, San Francisco
- Guerini D, Krebs J, Carafoli E (1984) Stimulation of the purified erythrocyte Ca^{2+} -ATPase by tryptic fragments of calmodulin. J Biol Chem 259:15172–15177
- Haberz P, Rodríguez-Castañeda F, Junker J, Becker S, Leonov A, Griesinger C (2006) Two new chiral EDTA-based metal chelates for weak alignment of proteins in solution. Org Lett 8:1275–1278
- Junge HJ, Rhee J-S, Jahn O, Varoquaux F, Spiess J, Waxham MN, Rosenmund C, Brose N (2004) Calmodulin and Munc13 form a Ca^{2+} sensor/effector complex that controls short-term synaptic plasticity. Cell 118:389–401
- Rhoads AR, Friedberg F (1997) Sequence motifs for calmodulin recognition. FASEB J 11:331–340
- Richmond JE, Weimer RM, Jorgensen EM (2001) An open form of syntaxin bypasses the requirement for *unc-13* in vesicle priming. Nature 412:338–341