

Functional renormalisation group for the $U(1)$ - T_5^6 TGFT with closure constraint

Vincent Lahoche^{a1} and Dine Ousmane Samary^{b,c2}

- a) Laboratoire de Physique Théorique, CNRS-UMR 8627, Université Paris-Sud 11, 91405 Orsay Cedex, France
- b) Max Planck Institute for Gravitational Physics, Albert Einstein Institute, Am Mühlenberg 1, 14476, Potsdam, Germany
- c) Faculté des Sciences et Techniques/ ICMIPA-UNESCO Chair, Université d'Abomey- Calavi, 072 BP 50, Benin

Abstract

This paper is focused on the functional renormalization group applied to the T_5^6 tensor model on the Abelian group $U(1)$ with closure constraint. In a first time we derive the flow equations for the couplings and mass parameters in a suitable truncation around the marginal interactions with respect to the perturbative power counting. In a second time, we study the behavior around the Gaussian fixed point, and show that the theory is non-asymptotically free. Finally, we discuss the UV completion of the theory. We show the existence of several non-trivial fixed points, study the behavior of the renormalization group flow around them, and point out evidence in favor of an asymptotically safe theory.

1 Introduction

Tensor models (TM) generalize matrix models and are considered as a convenient formalism for studying random geometries [1]-[17]. TM also offer an alternative to other approaches dealing with quantum gravity (QG) which is based on new mathematical/statistical tools, for example, the $1/N$ expansion recently discovered [16]-[17]. On the other hand, group field theory (GFT) is a quantum field theory over group manifolds and is considered as a second quantization of loop quantum gravity [18]-[22]. Both TM and GFT belong to the so-called background-independent scenario for QG. They aim at describing a rudimentary phase of the geometry of spacetime, namely when this geometry is hypothetically still in a discrete form, or at least not yet continuous. It is also named "pre-geometric" phase of our spacetime. Recently TM and GFT have been combined to provide a new class of field theories so called tensorial group field theory (TGFT). TGFTs improve the GFTs in order to allow for renormalization [23]-[32]. Moreover, it has been shown that several TGFTs models are asymptotically free in the UV, in other words, near the Gaussian fixed point [33]-[43].

The renormalization group (RG) method formulated first by Wilson [47]-[48] is a nonperturbative method which allows to interpolate smoothly between the UV laws and the IR phenomena in physical systems. The RG has a vast range of applications. A particularization of the RG, the functional renormalization group (FRG) is a realization of the RG concept in the framework of quantum and/or statistical field theory and is one of the best candidates for studying quantum fluctuations [49]. An

¹vincent.lahoche@th.u-psud.fr

²dine.ousmane.samary@aei.mpg.de

important property of this method is that the FRG could be used in regimes where perturbative calculations are invalid, for instance, at the vicinity of nontrivial fixed points in the infrared regime.

Recently much interest were focused on the FRG equation of various Matrix and TGFT models [39]-[46]. The differential equations of the flow were derived using Wetterich's equation [49]. The fixed points were given and further evidence of asymptotically safety and asymptotically freedom were derived around these fixed points in the UV.

The TGFT of the form T_5^6 on the $U(1)$ group with closure constraint is proved to be renormalizable [30]. The proof of this claim is performed using multi-scale analysis. The closure constraint also called gauge invariance condition can help to define the emergence of the metric on spacetime after phase transition and therefore makes this type of model relevant for the understanding the quantum theory of gravitation. The perturbative computation of the β -functions of such model is given in [38]. This paper aims at giving the FRG analysis to a renormalizable tensor model T_5^6 with closure constraint and for improving the conclusion given in [38]. In a truncation containing all relevant and marginal interactions, we find nontrivial fixed points. The FRG flows for coupling constants and for the mass parameter are solved numerically.

The paper is organized as follows. In section (2) we first present the model which is analysed in this paper, namely the T_5^6 model with closure constraints. In section (3) we give the flow equations of the coupling constants and mass parameter by using the dimensional renormalization parameters. In section (4) we give the nontrivial fixed points and provide the numerical solution of the flow equations. In section (5) the validity of the choice of the truncation of the effective action is discussed. The behavior of our model in the vicinity of these fixed points is also given. The conclusion and discussion are made in section (6).

2 The T_5^6 TGFT model

This section is devoted to a short review of the particular TGFT that we present in this paper. First, we define and give some properties of our model. Next, we discuss the canonical dimension that allows to make sense of the exponentiation of the functional action in the partition function.

2.1 The model

We consider fields ψ and $\bar{\psi}$ acting over the d copies of the group $U(1)$, i.e. $\psi, \bar{\psi} : U(1)^d \rightarrow \mathbb{C}$. In the Fourier representation, the fields variables $T_{\vec{p}}, \bar{T}_{\vec{p}}, \vec{p} \in \mathbb{Z}^d$ are maps $\bar{T}, T : \mathbb{Z}^d \rightarrow \mathbb{C}$, such that,

$$\psi(\vec{g}) = \sum_{\vec{p}} T_{\vec{p}} e^{i\vec{\theta} \cdot \vec{p}}, \quad \vec{g} = (\mathbf{g}_1, \dots, \mathbf{g}_d), \quad \vec{p} = (p_1, \dots, p_d), \quad \vec{\theta} = (\theta_1, \dots, \theta_d), \quad \theta_j \in [0, 2\pi). \quad (1)$$

The parameter θ_k with $\mathbf{g}_k \equiv e^{i\theta_k}$ is related to the parametrization of the $U(1)$ group such that $U(1) \approx S^1$. The theory we consider is described by its generating function or the vacuum-vacuum transition amplitude:

$$Z_\Lambda[J, \bar{J}] = e^{W_\Lambda[J, \bar{J}]} = \int d\mu_{C_\Lambda}(\bar{T}, T) e^{S_{int}[\bar{T}, T] + \langle \bar{J}, T \rangle + \langle \bar{T}, J \rangle} \quad (2)$$

where the notation $\langle \cdot, \cdot \rangle$ means: $\langle \bar{J}, T \rangle = \sum_{\vec{p} \in \mathbb{Z}^d} \bar{J}_{\vec{p}} T_{\vec{p}}$, $d\mu_{C_\Lambda}$ is the Gaussian measure with the covariance C_Λ such that:

$$\int d\mu_{C_\Lambda} T_{\vec{p}} \bar{T}_{\vec{p}'} = \frac{e^{-(\vec{p}^2 + m^2)/\Lambda^2}}{\vec{p}^2 + m^2} \delta\left(\sum_{i=1}^d p_i\right) \delta_{\vec{p}\vec{p}'} = C_\Lambda(\vec{p}, \vec{p}'), \quad (3)$$

$$\mathcal{W}^{(\ell)}(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3, \mathbf{g}_4) = \text{Diagram (8)} \quad (8)$$

$$\mathcal{X}^{(\ell)}(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3, \mathbf{g}_4, \mathbf{g}_5, \mathbf{g}_6) = \text{Diagram (9)} \quad (9)$$

$$\mathcal{Y}^{(\ell)}(\mathbf{g}_1, \mathbf{g}_2, \mathbf{g}_3, \mathbf{g}_4, \mathbf{g}_5, \mathbf{g}_6) = \text{Diagram (10)} \quad (10)$$

and the delta $\delta(\sum_{i=1}^d p_i)$ implements the closure constraint, see [23, 28]. Λ is the UV cutoff which will impose that the modulus of momentum vectors remains less than Λ , namely, $|\vec{p}| \leq \Lambda$. We keep in mind that we will eventually take the limit $\Lambda \rightarrow \infty$. We define a model by its action at a high (UV) energy scale. The classical action S_{int} is defined as a sum of tensorial invariances [1], [3]:

$$S_{int}[\bar{T}, T] = \sum_{b \in \mathcal{B}} \lambda_b \text{Tr}_b[\bar{T}, T]. \quad (4)$$

A tensor invariant is a polynomial in the tensor T and its conjugate \bar{T} which is invariant under the action of the tensor product of d independent copies of the unitary group $U(N)$. The sum is taken over a finite set \mathcal{B} of such invariants d -bubbles [1] associated with the couplings λ_b .

The interaction (4) of a tensor field theory in dimension $d = 5$ [30] is

$$S_{int}[\bar{T}, T] = \frac{\lambda_1}{2} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1, \dots, 4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} + \frac{\lambda_2}{3} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1, \dots, 6}} \mathcal{X}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \quad (5)$$

$$+ \lambda_3 \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1, \dots, 6}} \mathcal{Y}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6}, \quad (6)$$

where the symbols $\mathcal{W}^{(\ell)}$, $\mathcal{X}^{(\ell)}$ and $\mathcal{Y}^{(\ell)}$ are products of delta functions associated to tensor invariant interactions, and $\lambda_i(\Lambda)$ are coupling constants. For instance:

$$\mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} = \delta_{p_1 \ell p_4 \ell} \delta_{p_2 \ell p_3 \ell} \prod_{j \neq \ell} \delta_{p_{1j} p_{2j}} \delta_{p_{3j} p_{4j}}. \quad (7)$$

Such a kernel is called *bubble* [1], and can be pictured graphically as a 6-colored bipartite regular graph, with black and white vertices corresponding respectively to the fields T and \bar{T} , and each line corresponding to a Kronecker delta. As an example, the 4-valent bubble associated to the kernel $\mathcal{W}^{(1)}$ is depicted on Figure (1) below, and in the same way, all the interaction bubbles involved in the action S_{int} are defined as:

where the index ℓ takes values from 1 to 5, and refers to the single color characterizing each bubbles.




Figure 1: An example of bubble of order 4

2.2 Canonical dimension

Another important definition for our purpose concern the notion of *canonical dimension*. We will only give the essential here, and the reader interested in the details may consult [37]. In our model, the divergence degree for an arbitrary Feynman graph \mathcal{G} is given by [23, 30]

$$\omega(\mathcal{G}) = -2L(\mathcal{G}) + (F(\mathcal{G}) - R(\mathcal{G})) \quad (11)$$

where L is the number of propagator, F the number of faces. Let us pick an arbitrary orientation for all of the edges e and for all of the faces f . Then R is the rank of the incidence matrix ϵ_{fe} :

$$\epsilon_{fe} = \begin{cases} 1 & \text{if } e \in f \text{ and their orientation match} \\ -1 & \text{if } e \in f \text{ and their orientation do not match} \\ 0 & \text{otherwise.} \end{cases} \quad (12)$$

Note that the rank R does not depends on the chosen of the orientations. Denoting by $n_i(\mathcal{G})$ the number of vertices in \mathcal{G} with $2i$ external lines, the divergences subgraphs are said to be melonic [16, 23], if and only if they satisfy the following relation:

$$F(\mathcal{G}) - R(\mathcal{G}) = 3(L(\mathcal{G}) - \sum_i n_i(\mathcal{G}) + 1) \quad (13)$$

which, together with the topological relation $L(\mathcal{G}) = \sum_i i n_i(\mathcal{G}) - N(\mathcal{G})/2$ leads to:

$$\omega(\mathcal{G}) = 3 - \frac{N(\mathcal{G})}{2} - 2n_1(\mathcal{G}) - n_2(\mathcal{G}), \quad (14)$$

where $N(\mathcal{G})$ denote the number of external lines of \mathcal{G} . For $N = 4$, $\omega \leq 1$, the value 1 corresponding to melonic graphs with only 6-point interactions bubble. This conclusion indicates that perturbatively around the Gaussian Fixed Point (GFP), the coupling constant λ_1 scaling as Λ for some cut-off Λ , and we associate a canonical dimension $[\lambda_1] = 1$ to this constant. In the same way, we deduce that for a generic coupling λ_b , associated to a melonic bubble with N_b external lines:

$$[\lambda_b] = 3 - \frac{N_b}{2} \quad (15)$$

giving explicitly:

$$[m] = 1 \quad [\lambda_1] = 1 \quad [\lambda_2] = [\lambda_3] = 0. \quad (16)$$

3 Functional renormalization group with closure constraint

In this section we discuss the physical consequences of the renormalization group flow by truncating the space of actions. The procedure is standard, and consist in a systematic projection of the renormalization group flow into a finite dimensional subspace of generalized couplings. The approximated trajectory $\{\Gamma_k\}$ is then described by several functions, which are solution of a finite coupled system of differential equations, the so-called β -functions. The difficult point of this approach is to justify the choice of the truncation. For our purpose, we use a standard dimensional argument, and neglect all the interactions up to the marginal coupling with respect to the perturbative power counting. As discussed in this section, such a truncation make sense as long as the anomalous dimension remains small, giving a consistency condition for the validity of the truncation, which can be easily cheeked. However, we will see that, truncation may be introduce singular artefact, as lines of fixed points, which depends on the truncation.

We start this section by evaluating he Wetterich equation, and find the system of β -functions studied for our model. The asymptotic behavior in the UV is also provided.

3.1 Truncation and Regularization

The FRG method is based on the following deformation to our original partition function given in the equation (2) i.e.

$$Z_k[J, \bar{J}] = \int d\mu_{C_\Lambda}(\bar{T}, T) e^{S_{int}[\bar{T}, T] - \Delta S_k[\bar{T}, T] + \langle \bar{J}, T \rangle + \langle \bar{T}, J \rangle} \quad (17)$$

where we have added to the original action a IR cut-off $\Delta S_k[\bar{T}, T]$, defined as:

$$\Delta S_k[\bar{T}, T] = \sum_{\vec{p} \in \mathbb{Z}^5} R_k(|\vec{p}|) \bar{T}_{\vec{p}} T_{\vec{p}}. \quad (18)$$

The cut-off function R_k , depend on the real parameter k playing the role of a running cut-off, and is chosen such that:

- $R_s(\vec{p}) \geq 0$ for all $\vec{p} \in \mathbb{Z}^d$ and $s \in (-\infty, +\infty)$.
- $\lim_{s \rightarrow -\infty} R_s(\vec{p}) = 0$, implying: $\mathcal{Z}_{s=-\infty}[\bar{J}, J] = \mathcal{Z}[\bar{J}, J]$. This condition ensures that the original model is in the family (17). Physically, it means that the original model is recovered when all the fluctuations are integrated out.
- $\lim_{s \rightarrow \ln \Lambda} R_s(\vec{p}) = +\infty$, ensuring that all the fluctuations are frozen when $e^s = \Lambda$. As a consequence, the bare action will be represented by the initial condition for the flow at $s = \ln \Lambda$.
- For $-\infty < s < \ln \Lambda$, the cutoff R_s is chosen so that $R_s(|p| > e^s) \ll 1$, a condition ensuring that the UV modes $|p| > e^s$ are almost unaffected by the additional cutoff term, while $R_s(|p| < e^s) \sim 1$, or $R_s(|p| < e^s) \gg 1$, will guarantee that the IR modes $|p| < e^s$ are decoupled.

- $\frac{d}{d|\vec{p}|} R_s(\vec{p}) \leq 0$, for all $\vec{p} \in \mathbb{Z}^d$ and $s \in (-\infty, +\infty)$, which means that high modes should not be suppressed more than low modes.

The equation describing the flow of the couplings, the so called Wetterich equation has been established in [49] in the case of a theory with closure constraint: For a given cut-off R_k , the effective average action satisfies the following first order partial differential equation:

$$\partial_k \Gamma_k = \sum_{\vec{p} \in \mathbb{Z}^5} \partial_k R_k(|\vec{p}|) \cdot [\Gamma_k^{(2)} + R_k]^{-1}(\vec{p}, \vec{p}) \delta \left(\sum_{i=1}^5 p_i \right). \quad (19)$$

where Γ_k , the effective average action and is defined as the Legendre transform of the free energy $W_k := \ln[Z_k]$ as :

$$\Gamma_k[\bar{T}, T] + \sum_{\vec{p} \in \mathbb{Z}^5} R_k(|\vec{p}|) \bar{T}_{\vec{p}} T_{\vec{p}} := \langle \bar{J}, T \rangle + \langle \bar{T}, J \rangle - W_k[J, \bar{J}] \quad (20)$$

and

$$\Gamma_k^{(2)}(\vec{p}, \vec{p}') := \frac{\partial^2 \Gamma_k}{\partial T_{\vec{p}} \partial \bar{T}_{\vec{p}'}} \quad (21)$$

where T denote the mean field $T_{\vec{p}} := \frac{\partial W_k}{\partial J_{\vec{p}}}$, and is a gauge invariant field in the sens that: $T_{\vec{p}} = T_{\vec{p}} \delta \left(\sum_{i=1}^5 p_i \right)$.

The Wetterich flow equation is an exact differential equation which must be truncated, i.e. it must be projected to functions of few variables or even onto some finite-dimensional sub-theory space. However as in nonperturbative analysis [49], and discussed in introduction of this section the question of error estimate is very important and nontrivial in functional renormalization. One way to estimate the error in FRG is to improve the truncation in successive steps, i.e. to enlarge the sub-theory space by including more and more running couplings. The difference in the flows for different truncations gives a good estimate of the error. In addition, one can use different regulator functions R_k in a given (fixed) truncation and determine the difference of the RG flows in the infrared for the respective regulator choices. In this section, we adopt the simplest truncation, consisting in a restriction to the essential and marginal coupling with respect to the perturbative power counting (i.e. whose canonical dimension is upper or equal to zero). As mentioned before, such a truncation make sense as long as the anomalous dimension remains small, and a qualitative argument is the following. Let us define the anomalous dimension $\eta := \partial_s \ln(Z)$ (see equation (25) below). In the vicinity of a fixed point, η can reach to a non-zero value η_* . As a result, the effective propagator becomes:

$$\frac{Z^{-1}}{\vec{p}^2 + (m_s^2/Z)} \approx \frac{e^{-\eta_* s}}{\vec{p}^2 + m_*^2} \quad , \quad (22)$$

and then modifies the power counting (14), which becomes in the melonic sector (all the star-quantities refers to the non-Gaussian fixed point that we consider):

$$\omega_*(\mathcal{G}) = -(2 + \eta_*)L(\mathcal{G}) + (F(\mathcal{G}) - R(\mathcal{G})) = 3 - \frac{N}{2}(1 - \eta_*) - 3\eta_* n_3 - (1 + 2\eta_*)n_2 - (2 + \eta_*)n_1. \quad (23)$$

As a result, the canonical dimension (16) turn to be

$$[t_b]_* = 3 - \frac{N_b}{2}(1 - \eta_*) = [t_b] + \frac{N_b}{2}\eta_* \quad , \quad (24)$$

from which one can argue that, as long as $\eta_* \ll 1$, the classification in term of essential, inessential and marginal couplings remains unchanged, and the truncation around marginal couplings with respect to the perturbative power counting make sense.

Up to these consideration, our choice of truncation is the following:

$$\begin{aligned} \Gamma_k[\bar{T}, T] = & \sum_{\vec{p} \in \mathbb{Z}^5} \left(Z(k)\vec{p}^2 + m^2(k) \right) T_{\vec{p}} \bar{T}_{\vec{p}} + \frac{\lambda_1(k)}{2} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} \\ & + \frac{\lambda_2(k)}{3} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{X}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \\ & + \lambda_3(k) \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{Y}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6}. \end{aligned} \quad (25)$$

Note that we have adopted an additional restriction concerning the degree of the differential operator for the kinetic term, which can be viewed as the first term in the *derivative expansion*. One more time, a consistency check must be to introduce the next contribution, and evaluate its relative contribution. We will not considered this question in this paper.

We move on to the extraction of the truncated flow equations for m^2 , Z and λ_i from the full Wetterich equation (19). We write the second derivative of Γ_k as:

$$\Gamma_k^{(2)}[\bar{T}, T](\vec{p}, \vec{p}') = \left(-Z(k)\vec{p}^2 + m^2(k) \right) \delta \left(\sum_{i=1}^5 p_i \right) \delta_{\vec{p}\vec{p}'} + F_{k,(1)}[\bar{T}, T]_{\vec{p}, \vec{p}'} + F_{k,(2)}[\bar{T}, T]_{\vec{p}, \vec{p}'}$$

in such a way that all the field-dependent terms of order $2n$ are in $F_{k,(n)}$. In particular, $F_{k,(1)}$ depends on $\lambda_1(k)$, while $F_{k,(2)}$ depends on $\lambda_2(k)$ and $\lambda_3(k)$.

For the regulator R_k , we adopt the Litim's cut-off [51]:

$$R_k(|\vec{p}|) = Z(k)(k^2 - \vec{p}^2)\Theta(k^2 - \vec{p}^2), \quad (26)$$

and computing the first derivative with respect to k , we find:

$$k\partial_k R_k(|\vec{p}|) = \{k\partial_k Z(k)(k^2 - \vec{p}^2) + 2Z(k)k^2\}\Theta(k^2 - \vec{p}^2). \quad (27)$$

Hence, we are now in position to extract the flow equations for each couplings, which is the subject of the next section.

3.2 Flow equations in the UV regime

We will deduce the flow equation in the UV regime. In this regime, all the sums can be replaced by integration following the arguments of [41], essentially because the divergences of the integral approximations are the same as the exact sums. The method consist to a formal expansion of the r.h.s of the Wetterich equation (19) in power of couplings, and identification of the corresponding terms in the l.h.s. The r.h.s involve in general some contractions between the $F_{k,(n)}$ and the effective propagator $\partial_k R_k$. And in this UV regime, only the melonic graphs contribute.

Expanding the r.h.s and the l.h.s of the flow equation, we obtain the following relations (in matrix notations):

$$k\partial_k \Gamma_{k,(1)} = -\text{Tr}_{GI} \left[\partial_k R_k \mathcal{K}_k^{-1} F_{k,(1)} \mathcal{K}_k^{-1} \right], \quad (28)$$




Figure 2: Typical contribution to the 2-points observable. The dashed line represent the contraction with the propagator $\partial_k R_k$.

$$k\partial_k \Gamma_{k,(2)} = -\text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1}] + \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^2], \quad (29)$$

$$k\partial_k \Gamma_{k,(3)} = 2\text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(1)} \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1}] - \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^3]. \quad (30)$$

where the subscript GI means "Gauge Invariant" sums, in the sense that all the terms summing involve a product with a delta $\delta\left(\sum_{i=1}^5 p_i\right)$, $\Gamma_{k(n)}$ means the term of order n in the truncation of equation (25), and:

$$\mathcal{K}_k^{-1} := \frac{1}{Z(k)\vec{p}^2 + m^2(k) + R_k(|\vec{p}|)}. \quad (31)$$

3.2.1 Flow equations for Z and m^2

Expanding the trace in the r.h.s of the equation (28), we find, using the expressions (26) and (27)

$$\text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(1)} \mathcal{K}_k^{-1}] = \sum_{\vec{p}^2 \leq k^2} \frac{k\partial_k Z(k)(k^2 - \vec{p}^2) + 2Z(k)k^2}{[Z(k)k^2 + m^2(k)]^2} F_{k(1)}(\vec{p}, \vec{p}) \delta\left(\sum_{i=1}^5 p_i\right). \quad (32)$$

Graphically, this contribution can be pictured as in Figure (2), where the dashed line represent the contraction with the propagator $\partial_k R_k$. The operator $F_{k(1)}$ is a sum over the intermediate colors labeling each 4-valent bubbles: $F_{k(1)} = \sum_{i=1}^5 F_{k(1)}^{(i)}$, and each term contribute separately to the wave function and mass flow. As a result, we focus our attention on the computation of the trace for the case $i = 1$. As explained in [41], we will identify the contribution to the coupling of a given bubble in the l.h.s by expanding the r.h.s around its local approximation i.e. around the value $q = 0$, where q denote the "external momentum" shared by the red lines in the Figure (2). The two-points case is in a sense the more interesting, because for the wave function contribution, the local expansion require the first deviation to the exact local approximation, corresponding to the mass-term. This first deviation is proportional to q^2 . Then, identifying the terms in front of each powers of q^2 , we find:

$$k\partial_k m^2(k) = -5\lambda_1(k) \frac{k\partial_k Z(k)(S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^2} \quad (33)$$

$$k\partial_k Z(k) = \frac{-2\lambda_1(k)Z(k)k^2 S_1''}{[Z(k)k^2 + m^2(k)]^2 + \lambda_1(k)(S_1''k^2 - S_2'')} \quad (34)$$

where the factor 5 in front of the equation (33) takes into account the contributions for each i , S_i'' denote the coefficient in front of q^2 in the expansion of $S_i(q)$ in power of q , and the sums S_1 and S_2 are :

$$S_1(q) = \sum_{\vec{p} \in \mathbb{Z}^4} \delta \left(\sum_{i=1}^4 p_i + q \right) \theta(k^2 - q^2 - \vec{p}^2), \quad (35)$$

$$S_2(q) = \sum_{\vec{p} \in \mathbb{Z}^4} \vec{p}^2 \delta \left(\sum_{i=1}^4 p_i + q \right) \theta(k^2 - q^2 - \vec{p}^2). \quad (36)$$

Since we will be mostly interested in the large- k limit, we can approximate the sums by integrals, replacing the Kronecker deltas by Dirac deltas. The support of the integrals is in the intersection of the hyperplane of equation $q + \sum_{l=1}^4 p_l = 0$ and the 4-ball of radius $\sqrt{k^2 - q^2}$. Note that the Kronecker delta of the closure constraint can be rewritten as $q + \vec{p} \cdot \vec{n} = 0$, where $\vec{n} = (1, 1, 1, 1) \in \mathbb{R}^4$ is the vector with all components equals to 1. Using the rotational invariance of our integral, we can choose one of our coordinate axis to be in the direction \vec{n} . If we choose the axis 1 in this direction, our constraint writes as $\delta(q + p'_1 |\vec{n}|) = \delta(q + 2p'_1) = \delta(q/2 + p'_2)/2$, and we find the following integral approximation:

$$S_1(q) \simeq \frac{1}{2} \Omega_3 \left[k^2 - \frac{5q^2}{4} \right]^{\frac{3}{2}} \quad (37)$$

$$S_2(q) \simeq \frac{1}{2} \left[\frac{5q^2}{4} + \frac{3}{5} \left(k^2 - \frac{5q^2}{4} \right) \right] \Omega_3 \left(k^2 - \frac{5q^2}{4} \right)^{\frac{3}{2}}, \quad (38)$$

where $\Omega_d := \pi^{d/2} / \Gamma(d/2 + 1)$ is the volume of the unit d -ball, with the special value $\Omega_3 = 4\pi/3$. Using this integral approximation, we obtain:

$$S_1(0) = \frac{2}{3} \pi k^3, \quad S_2(0) = \frac{2}{5} \pi k^5, \quad S_1'' = -\frac{5}{4} \pi k, \quad S_2'' = -\frac{5}{12} \pi k^3 \quad (39)$$

giving:

$$k \partial_k m^2(k) = -\frac{4\pi}{3} \lambda_1(k) \frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^2} k^5 \quad (40)$$

$$\eta(k) = \frac{5\pi}{2} \lambda_1(k) \frac{k^3}{[Z(k)k^2 + m^2(k)]^2 - \lambda_1(k) \frac{5}{6} \pi k^3} \quad (41)$$

with the anomalous dimension $\eta(k)$ defined as:

$$\eta(k) := k \partial_k \ln(Z(k)). \quad (42)$$

Note that, in this case, and for the other computations, the extraction of the local approximation in the UV limit brings up a very nice property of the melonic sector, called *traciality*. Traciality is a concept firstly introduced in a perturbative renormalization framework, ensuring that local approximation of high subgraphs make sense in the TFGT context [23], [25].




Figure 3: Contributions coming from the 6-points interactions to the 4-point interaction.




Figure 4: Contribution to the 4-point interaction involved two vertices.

3.2.2 Flow equation for λ_1

The flow equation for λ_1 (29) involves two traces that we will compute separately. The first trace $\text{Tr}_{GI}[\partial_k R_k \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1}]$ involves two typical contributions that we have pictured in Figure (3) below. Note that the two diagrams pictured on this Figure have the same connectivity as the 4-valent interaction with intermediate line of color red, and we will only consider the flow equation for the coupling attached to one of the five color.

The computation of these two contributions follows exactly the same way as the extraction of the flow equation for the mass parameter. We expand with respect to the external momentum q (i.e. the momentum around the external face sharing the same line as the four internal faces) around $q = 0$, the first term of the expansion giving the relevant contribution. Because the two sums involve a loop of length one, they can be expressed in terms of the two sums S_1 and S_2 , and we find:

$$\begin{aligned} \text{Tr}_{GI} [k \partial_k R_k \mathcal{K}_k^{-1} F_{k,(2)(3)a} \mathcal{K}_k^{-1}] &= 4\lambda_3 \frac{k \partial_k Z(k) (S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^2} \\ &\times \sum_{\{\vec{p}_i\}_{i=1,\dots,4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} + \mathcal{O}(q), \end{aligned} \quad (43)$$

$$\begin{aligned} \text{Tr}_{GI} [k \partial_k R_k \mathcal{K}_k^{-1} F_{k,(2)(3)b} \mathcal{K}_k^{-1}] &= \lambda_2 \frac{k \partial_k Z(k) (S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^2} \\ &\times \sum_{\{\vec{p}_i\}_{i=1,\dots,4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} + \mathcal{O}(q). \end{aligned} \quad (44)$$

The contribution of the last trace $\text{Tr}_{GI}[\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^2]$ is graphically pictured in Figure (4), where the dotted line means contraction with Kronecker delta. Expanding in local approximation, we find, for the extra-local contribution with $q = 0$:

$$\text{Tr}_{GI} [k \partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1),(4)} \mathcal{K}_k^{-1})^2] \approx \lambda_1^2(k) \frac{k \partial_k Z(k) (S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^3} \quad (45)$$

$$\times \sum_{\{\vec{p}_i\}_{i=1,\dots,4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4}. \quad (46)$$




Figure 5: Contribution to the flow of λ_3

Hence, summing the contributions (43), (44) and (45), and using the integral approximation for the sums, we find:

$$k\partial_k\lambda_1(k) = -(\lambda_2 + 4\lambda_3)\frac{4\pi}{15}\frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^2}k^5 + \lambda_1^2(k)\frac{4\pi}{15}\frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^3}k^5 \quad (47)$$

3.2.3 Flow equations for λ_2 and λ_3

The only contribution to λ_3 is pictured in Figure (5) below, corresponding to the trace:

$$\text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(1)} \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1}].$$

Indeed, it is the only contraction in the melonic sector with the same connectivity as the 6-point interaction associated with the coupling λ_3 . The local approximation of the diagram can be computed exactly as for the contribution (3) to λ_1 , and we obtain:

$$\begin{aligned} \text{Tr}_{fig(5)}(q=0) &= \lambda_1\lambda_3 \frac{k\partial_k Z(k)(S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^3} \\ &\times \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{Y}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \end{aligned} \quad (48)$$

and:

$$k\partial_k\lambda_3(k) = \frac{16\pi}{15}\lambda_1\lambda_3 \frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^3}k^5. \quad (49)$$

In the same way, the contribution to the local approximation of the diagram (6)a to the flow equation for λ_2 writes as:

$$\begin{aligned} \text{Tr}_{fig(6)a}(q=0) &= \lambda_1\lambda_2 \frac{k\partial_k Z(k)(S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^3} \\ &\times \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{X}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \end{aligned} \quad (50)$$

Finally, the contribution coming from the diagram (6)b in Figure (6) involves a loop with two delta propagators, and:

$$\text{Tr}_{fig(6)b}(q=0) = \lambda_1^3(k) \frac{k\partial_k Z(k)(S_1(0)k^2 - S_2(0)) + 2Z(k)k^2 S_1(0)}{[Z(k)k^2 + m^2(k)]^4}$$




Figure 6: Contribution to the flow of λ_2

$$\times \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{X}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6}. \quad (51)$$

Grouping together the contributions (50) and (51), we find:

$$k\partial\lambda_2 = \frac{24\pi}{15} \lambda_1 \lambda_2 \frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^3} k^5 - \frac{12\pi}{15} \lambda_1^3 \frac{\eta(k) + 5}{[Z(k)k^2 + m^2(k)]^4} k^5. \quad (52)$$

3.2.4 Dimensionless renormalized parameters

Taking into account the canonical dimension define in Section (2.2), the renormalized dimensionless couplings are defined as:

$$m = \sqrt{Z} k \bar{m} \quad \lambda_1 = Z^2 k \bar{\lambda}_1 \quad \lambda_2 = Z^3 \bar{\lambda}_2 \quad \lambda_3 = Z^3 \bar{\lambda}_3 \quad (53)$$

Using the flow equations (41), (40), (47), (52) and (49), we find for the dimensionless renormalized couplings the following autonomous system:

$$\eta(k) = \frac{5\pi}{2} \bar{\lambda}_1(k) \frac{1}{[1 + \bar{m}^2(k)]^2 - \bar{\lambda}_1(k) \frac{5}{6}\pi} \quad (54)$$

$$\beta_{m^2} = -(2 + \eta) \bar{m}^2(k) - \frac{4\pi}{3} \bar{\lambda}_1(k) \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^2} \quad (55)$$

$$\beta_{\lambda_1} = -(1 + 2\eta) \bar{\lambda}_1(k) - (\bar{\lambda}_2 + 4\bar{\lambda}_3) \frac{4\pi}{15} \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^2} + \bar{\lambda}_1^2(k) \frac{4\pi}{15} \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^3} \quad (56)$$

$$\beta_{\lambda_3} = -3\eta \bar{\lambda}_3(k) + \frac{16\pi}{15} \bar{\lambda}_1 \bar{\lambda}_3 \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^3}. \quad (57)$$

$$\beta_{\lambda_2} = -3\eta \bar{\lambda}_2(k) + \frac{24\pi}{15} \bar{\lambda}_1 \bar{\lambda}_2 \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^3} - \bar{\lambda}_1^3 \frac{12\pi}{15} \frac{\eta(k) + 5}{[1 + \bar{m}^2(k)]^4}, \quad (58)$$

with the definition : $\beta_i := k\partial_k \bar{i}$, $i \in \{m^2, \lambda_1, \lambda_2, \lambda_3\}$.

4 Fixed points in the UV regime

At vanishing β -functions we obtain a fixed points. But these fixed points does not get any quantum correction and is called the Gaussian fixed point. In the neighborhood of these fixed points, the stability is determined by the linearized system of β -functions. All these points are studied in detail in this section.

4.1 Vicinity of the Gaussian fixed point

The autonomous system describing the flow of the dimensionless couplings admits a trivial fixed point for the values $\bar{\lambda}_1 = \bar{\lambda}_2 = \bar{\lambda}_3 = \bar{m} = 0$ called Gaussian Fixed Point (GFP). Expanding our equations around this points, we find the reduced autonomous system:

$$\begin{cases} \beta_{m^2} & \approx -2\bar{m}^2 - \frac{20\pi\bar{\lambda}_1}{3}, \\ \beta_{\lambda_1} & \approx -\bar{\lambda}_1 - \frac{4\pi}{3}(\bar{\lambda}_2 + 4\bar{\lambda}_3)\left(1 + \frac{\pi}{2}\bar{\lambda}_1 + 2\bar{m}^2\right) - \frac{11\pi}{3}\bar{\lambda}_1^2, \\ \beta_{\lambda_2} & \approx \frac{\pi}{2}\bar{\lambda}_1\bar{\lambda}_2, \\ \beta_{\lambda_3} & \approx -\frac{13\pi}{6}\bar{\lambda}_1\bar{\lambda}_3 \end{cases} \quad (59)$$

and the anomalous dimension:

$$\eta(k) \approx \frac{5\pi\bar{\lambda}_1}{2}. \quad (60)$$

These equations give the qualitative behavior of the RG trajectories around the GFP. In order to study its stability, we compute the *stability matrix* $\beta_{ij} := \partial_i \beta_j$ $i \in \{m^2, \lambda_1, \lambda_2, \lambda_3\}$, and evaluate each coefficient at the GFP. We find:

$$\beta_{ij}^{GFP} := \begin{pmatrix} -2 & 0 & 0 & 0 \\ -\frac{20\pi}{3} & -1 & 0 & 0 \\ 0 & -\frac{4\pi}{3} & 0 & 0 \\ 0 & -\frac{16\pi}{3} & 0 & 0 \end{pmatrix}, \quad (61)$$

with eigenvalues $(-2, -1, 0, 0)$ and eigenvectors $e_1^{GFP} = (\frac{9}{160\pi^2}, \frac{3}{8\pi}, \frac{1}{4}, 1)$; $e_2^{GFP} = (0, \frac{3}{16\pi}, \frac{1}{4}, 1)$; $e_3^{GFP} = (0, 0, 0, 1)$; $e_4^{GFP} = (0, 0, 1, 0)$. One recall that the *critical exponents* are the opposite values of the eigenvalues of the β_{ij} , and that the fixed point can be classified following the sign of their critical exponents. Hence, we have two relevant directions in the UV, with critical exponents 2 and 1, and two marginal couplings with zero critical exponents. Moreover, note that the critical exponents are equal to the canonical dimension around the GFP. Finally, note that the previous system of equations admits other fixed points, or a *line of fixed points* in addition to the Gaussian one, for the values: $\bar{\lambda}_1 = \bar{m} = 0$; $\bar{\lambda}_3 = -\bar{\lambda}_2/4$. This fixed point occurs as well as in the non-perturbation analysis, and we will return on this subject in the next Section.

For the moment, we are in position to discuss the qualitative flow diagram around the Gaussian fixed point. First of all, note that all the coefficients of the beta function of the system (59) are not negative definite. This fact seems to be a special feature of this model, meaning that the weight of the anomalous dimension does not dominate the vertex contribution. This fact is a first difference with respect to the similar non-Abelian ϕ^6 model studied in [37]. However, the analysis providing in this reference remains true, and the model is not asymptotically free. We will not repeat the complete analysis given in [37], but a qualitative argument is the following. Exploiting the fact that

the hyperplans $\bar{\lambda}_2 = 0$ and $\bar{\lambda}_3 = 0$ are invariant under the flow, we can look at only a two-dimensional reduction of the complete system (59). We choose $\bar{\lambda}_2 = 0$, and plot the numerical integration of the reduced flow equation in Figure (7) (on the left) below. In the domain, $\bar{\lambda}_3 > 0$, even if a given




Figure 7: Phase portrait in the plans $(\bar{\lambda}_1, \bar{\lambda}_3)$ for $\bar{\lambda}_2 = 0$ (one the left) and in the plan $(\bar{\lambda}_1, \bar{\lambda}_2)$, for $\bar{\lambda}_3 = 0$.

trajectory approaches of the Gaussian fixed point, $\bar{\lambda}_1$ reaches a negative value, and it is ultimately repelled for k sufficiently large. The same phenomenon occurs for $\bar{\lambda}_2$ in the plan $\bar{\lambda}_2 < 0$ (see the Figure(7) one the right). The issue of the UV completion of a theory which is non-Asymptotically free is one of the difficult question that we address to the non-perturbative renormalization group machinery, and the rest of this paper is essentially devoted to this one.

4.2 Non-Gaussian fixed points

Solving numerically the system (40)-(49), we find some non-Gaussian fixed points, whose relevant characteristics are summarized in the Table (8) below. In addition to these non-Gaussian fixed point, the system admit a *line of fixed points, LFP*, for the the values:

$$LFP = \{\bar{m}^2 = 0, \bar{\lambda}_1 = 0, \bar{\lambda}_2 = -4\bar{\lambda}_3\}, \quad (62)$$

with critical exponents:

$$\left\{ \begin{array}{l} \theta^{(1)} = -2, \\ \theta^{(2)} = 0, \\ \theta^{(3)} = -\frac{1}{2}(1 + \sqrt{1 - \frac{128}{9}\pi^2\bar{\lambda}_2}), \\ \beta_{\lambda_3} = -\frac{1}{2}(1 - \sqrt{1 - \frac{128}{9}\pi^2\bar{\lambda}_2}). \end{array} \right. \quad (63)$$

The denominator of η , $D := [1 + \bar{m}^2(k)]^2 - \bar{\lambda}_1(k)\frac{5}{6}\pi$ introduce a singularity in the flow. At the Gaussian fixed point, and in a sufficiently small domain around, $D > 0$. But further away from the GFP, D may cancel, creating in the $(\bar{\lambda}_1, \bar{m}^2)$ -plan a singularity line. The area below this line where $D < 0$ is thus disconnected from the region $D > 0$ connected to GFP. Then, we ignore for

FP	\bar{m}^2	$\bar{\lambda}_1$	$\bar{\lambda}_2$	$\bar{\lambda}_3$	η	$\theta^{(1)}$	$\theta^{(2)}$	$\theta^{(3)}$	$\theta^{(4)}$
FP_1	-0.3	0.005	0.0009	-0.0002	-6.3	-299	56.1	-11.7	5.8
FP_2	-0.7	0.008	0.0006	-0.0002	0.76	-7.4-1.9i	-7.4+1.9i	3.34	-0.12
FP_3	-0.9	0.0007	$3.32 \cdot 10^{-6}$	0.	1.3	-66.7	-42.63	-27.7	1.80
FP_4	-0.8	0.04	-0.02	0.	-5.9	-144.8	-14.4	-7.5	-5.4
FP_5	0.06	-0.006	0.002	0.	-0.04	1.9	1.09	-0.04	-0.01
FP_6	1.32	-0.5	-0.06	0.	-0.6	3.0	-1.23	-1.13	-0.39

Figure 8: Summary of the properties of the non-Gaussian fixed points. Again, the critical exponents θ^i are the opposite values of the eigenvalues of the stability matrix: $\beta_* =: \text{diag}(-\theta_*^1, -\theta_*^2, -\theta_*^3, -\theta_*^4)$.

our purpose the fixed points in the disconnected region, for which $D < 0$. A direct computation show that only the fixed points FP_2 , FP_3 , FP_5 and FP_6 are relevant for an analysis in the domain connected to the Gaussian fixed point.

- The fixed points FP_2 and FP_3 are very similar. They have three irrelevant directions and one relevant direction in the UV. For each of these fixed points, the three irrelevant directions span a three dimensional manifold on which the trajectory runs toward the fixed point in the IR, while the trajectories out side are repelled of this critical surface, as pictured on Figure (9). This picture, the existence of a separatrix between two connected region of the phase space is reminiscent of a critical behavior, with phase transition between a broken and a symmetric phase, and these separatrix are *IR-critical surfaces*. This interpretation is highlighted for the two fixed points in the zero momenta limit. Indeed, in both cases, the contributions in the effective action of the terms proportionals to $\bar{\lambda}_2$ and $\bar{\lambda}_3$ can be neglected in comparison with the contributions of the two first terms, leading in the first approximation a Ginsburg-Landau equation for ϕ^4 scalar complex theory. Note that for FP_2 two critical exponents are complex, providing some oscillations of the trajectories, and implying that the fixed point is an IR-attractor in the two-dimensional manifold spanned by the eigenvectors corresponding to these two critical exponents. Moreover, the fixed point FP_5 appears to be an IR fixed point, with coordinates of opposite sign.

- The fixed point FP_5 has two relevant and two irrelevant directions in the UV. The relevant directions in the UV span a two-dimensional manifold corresponding to a *UV-multicritical surface*. Such a surface is interesting for the UV-completion of the theory. Indeed, all the trajectories in the surface are oriented toward the fixed point in the UV, while the dimension of the surface give an interesting number of physical parameter, providing an evidence in favor of the *asymptotic safety* of the model in the UV [52].

- Finally, we have the line of fixed point, for which we will distinguish four cases:

- i* In the domain $d_1 = \{\bar{\lambda}_2 < 0\}$ we have two relevant, one marginal and one irrelevant directions.
- ii* At the point $d_2 = \{\bar{\lambda}_2 = 0\}$, we recover the GFP, with two relevant and two marginal directions.
- iii* In the domain $d_3 = \{\bar{\lambda}_2 \in]0, (\frac{3}{8\pi})^2\}$ we have three relevant and one marginal directions. One more time, this section of the critical line is interesting in view of the UV-completion of the theory and provide a supplementary evidence in favor of asymptotic safety. Indeed, in each points, the rele-




Figure 9: Qualitative behavior of the RG trajectories around an IR fixed point. The critical surface (in grey) is spanned by the relevant directions in the IR, and the arrows are oriented toward the IR direction.

vant directions in the UV span a three dimensional UV-critical surface, in favor of the existence of a non-trivial asymptotically safe theory with three independent physical parameters. This line of fixed point has been recently discussed in [43] for a similar model improved by unconnected interaction bubbles.

iv In the domain $d_4 = \{\bar{\lambda}_2 > (\frac{3}{8\pi})^2\}$ The situation is very reminiscent of the previous one. We have three eigenvalues with negative real part and one equal to zero. Hence, we have three relevant and one marginal directions. The only difference in comparison with the domain d_3 is that the eigenvalue have non-zero imaginary parts, giving some oscillations and attractor phenomena in the trajectories.

Finally, we briefly discuss the values of the anomalous dimensions. With our conventions, the couplings of the relevant operator are suppressed as a power of k in the UV limit $k \rightarrow \infty$. The couplings decreases when the trajectory goes away from the UV regime. However, the power law behavior is limited to the attractive region of the fixed point, far from its scaling regime it can deviate from the power law one. And we can evaluate this deviation. For instance, in the vicinity of FP_5 , one deduce from (24) that the canonical dimension becomes:

$$[t_b]_{FP_5} \approx 3 - 1.6 \frac{N_b}{2}, \quad (64)$$

from which we deduce that all the interaction of valence up or equal to four becomes inessential. The same phenomenon occurs in the vicinity of FP_4 , where all the interactions up to these of valence four become inessential. In contrary, at the fixed points FP_2 and FP_4 the anomalous dimension is positive, meaning that the power counting is improved with respect to the Gaussian one, and irrelevant operators are enhanced in the UV.

5 Truncation with an interaction of valence 8

This section aims to identify how the adding of interaction of the valence 8 may modify the flow equation and the fixed point of our model. This means that the effective action is now truncate to

satisfy the following form:

$$\begin{aligned}
\Gamma_k[\bar{T}, T] &= \sum_{\vec{p} \in \mathbb{Z}^5} \left(Z(k) \vec{p}^2 + m^2(k) \right) \bar{T}_{\vec{p}} T_{\vec{p}} + \frac{\lambda_1(k)}{2} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,4}} \mathcal{W}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} + \\
&+ \frac{\lambda_2(k)}{3} \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{X}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \\
&+ \lambda_3(k) \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,6}} \mathcal{Y}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6}^{(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} \\
&+ \sum_{i=1}^4 \kappa_i \lambda_{4,i}(k) \sum_{\ell=1}^3 \sum_{\{\vec{p}_i\}_{i=1,\dots,8}} \mathcal{Z}_{\vec{p}_1, \vec{p}_2, \vec{p}_3, \vec{p}_4, \vec{p}_5, \vec{p}_6, \vec{p}_7, \vec{p}_8}^{i,(\ell)} T_{\vec{p}_1} \bar{T}_{\vec{p}_2} T_{\vec{p}_3} \bar{T}_{\vec{p}_4} T_{\vec{p}_5} \bar{T}_{\vec{p}_6} T_{\vec{p}_7} \bar{T}_{\vec{p}_8}, \quad (65)
\end{aligned}$$

where we assume that the last term of the action (65) takes into account all contributions of melonic interactions of the form T^8 with combinatorial factor κ_i , $i = 1, 2, 3, 4$, and the coupling constants $\lambda_{4,i}$, $i = 1, 2, 3, 4$ are related to the vertex $V_{4,i}$ see figures (10). We get for $\Gamma_{k,(3)}$ and $\Gamma_{k,(4)}$ the flow




Figure 10: Graphical representation of the vertices of valence 8

equations:

$$\begin{aligned}
k \partial_k \Gamma_{k,(3)} &= 2 \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(1)} \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1}] - \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^3] \\
&- \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(3)} \mathcal{K}_k^{-1}], \quad (66)
\end{aligned}$$

and

$$\begin{aligned}
k \partial_k \Gamma_{k,(4)} &= -\text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(4)} \mathcal{K}_k^{-1}] + \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(2)} \mathcal{K}_k^{-1})^2] \\
&+ \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^4] \\
&- 3 \text{Tr}_{GI} [\partial_k R_k \mathcal{K}_k^{-1} F_{k,(2)} \mathcal{K}_k^{-1} (F_{k,(1)} \mathcal{K}_k^{-1})^2], \quad (67)
\end{aligned}$$

Setting $\kappa_i = 1, \forall i$, and taking into account the dimensionless renormalized parameter, and grouping all melonic contributions, the flow equations of the coupling constants $\bar{\lambda}_2$, $\bar{\lambda}_3$ and $\bar{\lambda}_{4,i}$, are:

$$\left\{ \begin{array}{l}
\beta_{m^2} = -(2 + \eta) \bar{m}^2 - \frac{4\pi}{3} \bar{\lambda}_1 \frac{\eta+5}{[1+\bar{m}^2]^2}, \\
\beta_{\lambda_1} = -(1 + 2\eta) \bar{\lambda}_1 - (\bar{\lambda}_2 + 4\bar{\lambda}_3) \frac{4\pi}{15} \frac{\eta+5}{[1+\bar{m}^2]^2} + \bar{\lambda}_1^2 \frac{24\pi}{15} \frac{\eta+5}{[1+\bar{m}^2]^3}, \\
\beta_{\lambda_2} = -3\eta \bar{\lambda}_2 + \frac{24\pi}{15} \bar{\lambda}_1 \bar{\lambda}_2 \frac{\eta+5}{[1+\bar{m}^2]^3} - \bar{\lambda}_3^3 \frac{12\pi}{15} \frac{\eta+5}{[1+\bar{m}^2]^4} - \frac{48\pi}{15} \bar{\lambda}_{4,2} \frac{\eta+5}{[1+\bar{m}^2]^2} - \frac{12\pi}{15} \bar{\lambda}_{4,4} \frac{\eta+5}{[1+\bar{m}^2]^2}, \\
\beta_{\lambda_3} = -3\eta \bar{\lambda}_3 + \frac{16\pi}{15} \bar{\lambda}_1 \bar{\lambda}_3 \frac{\eta+5}{[1+\bar{m}^2]^3} - \frac{8\pi}{15} \bar{\lambda}_{4,1} \frac{\eta+5}{[1+\bar{m}^2]^2} - \frac{36\pi}{15} \bar{\lambda}_{4,3} \frac{\eta+5}{[1+\bar{m}^2]^2} - \frac{8\pi}{15} \bar{\lambda}_{4,4} \frac{\eta+5}{[1+\bar{m}^2]^2}, \\
\beta_{\lambda_{4,1}} = (1 - 4\eta) \bar{\lambda}_{4,1} + \frac{16\pi}{15} \bar{\lambda}_{4,1} \bar{\lambda}_1 \frac{\eta+5}{[1+\bar{m}^2]^3} + \frac{4\pi}{15} \bar{\lambda}_3^2 \frac{\eta+5}{[1+\bar{m}^2]^3} - \frac{24\pi}{15} \bar{\lambda}_1^2 \bar{\lambda}_3 \frac{\eta+5}{[1+\bar{m}^2]^4} + \frac{4\pi}{15} \bar{\lambda}_1^4 \frac{\eta+5}{[1+\bar{m}^2]^5}, \\
\beta_{\lambda_{4,2}} = (1 - 4\eta) \bar{\lambda}_{4,2} + \frac{32\pi}{15} \bar{\lambda}_{4,2} \bar{\lambda}_1 \frac{\eta+5}{[1+\bar{m}^2]^3} + \frac{4\pi}{15} \bar{\lambda}_2^2 \frac{\eta+5}{[1+\bar{m}^2]^3} - \frac{12\pi}{15} \bar{\lambda}_1^2 \bar{\lambda}_2 \frac{\eta+5}{[1+\bar{m}^2]^4}, \\
\beta_{\lambda_{4,3}} = (1 - 4\eta) \bar{\lambda}_{4,3} + \frac{24\pi}{15} \bar{\lambda}_{4,3} \bar{\lambda}_1 \frac{\eta+5}{[1+\bar{m}^2]^3}, \\
\beta_{\lambda_{4,4}} = (1 - 4\eta) \bar{\lambda}_{4,4} + \frac{24\pi}{15} \bar{\lambda}_{4,4} \bar{\lambda}_1 \frac{\eta+5}{[1+\bar{m}^2]^3} + \frac{8\pi}{15} \bar{\lambda}_2 \bar{\lambda}_3 \frac{\eta+5}{[1+\bar{m}^2]^3} - \frac{24\pi}{15} \bar{\lambda}_1^2 \bar{\lambda}_3 \frac{\eta+5}{[1+\bar{m}^2]^4}
\end{array} \right. \quad (68)$$

FP	FP_1	FP_2	FP_3	FP_4	FP_5	FP_6	FP_7	FP_8	FP_9	FP_{10}
\bar{m}^2	-1.07	-0.91	-0.84	-0.84	-1.22	-0.75	-0.76	-0.74	-0.59	1.45
λ_1	0.004	0.005	0.02	0.02	0.009	0.006	0.045	0.05	0.14	-0.65
$10^2\lambda_2$	0.1	0.03	0.7	-0.4	0.2	-0.01	2	1	-40	-16
$10^3\lambda_3$	-0.3	0.04	-3	0.	-0.7	0.02	-10	-10	-0.6	-3
$10^2\lambda_{4,1}$	0.01	-0.04	0.1	0.01	0.001	-230	0.3	0.3	0.09	-0.3
$10^3\lambda_{4,2}$	-0.04	-0.01	0.	-1	-0.06	-2000.	-0.9	-1	100	-10
$10^3\lambda_{4,3}$	0.	0.	-0.3	-0.9	0.	0.	0.	0.	0.	0.
$10^2\lambda_{4,4}$	0.003	690	0.02	0.4	0.006	670	0.05	0.2	0.03	-0.03
η	-6.8	-6.3	-6.0	-6.0	2.6	0.9	-5.9	-5.6	-0.65	-0.66
$\theta^{(1)}$	307.8	289.8	179.6	180.0	-110.7	24.7	142.6	137.9	121.6	3.2
$\theta^{(2)}$	-245.8	-112.0	-12+25i	-45.9	-50+13i	14.5	19+6i	20+4i	30	3.1
$\theta^{(3)}$	173.0	-67.8	-12-25i	31	-50-13i	12.2	19-6i	20-4i	23.6	-3.0
$\theta^{(4)}$	113.6	31+6.7i	21+10i	26	-39	7.7	11.7	13+1.7i	18.9	2.8
$\theta^{(5)}$	77+19i	31-6.6i	21-10i	-22	-33	-6	1.2+10i	13-1.7i	17.2	2.7
$\theta^{(6)}$	77-19i	-28.5	-15	10	24.8	5.7	1.2-10i	11	9.8	1.0
$\theta^{(7)}$	-67	-19	6	6.1	-18	-5	9.8	-5.7	6.3	0.7
$\theta^{(8)}$	7.2	5.8	2.2	4	-2.3	-1.3	5.5	5.4	5.2	0.3

Figure 11: Summary of the properties of the non-Gaussian fixed points in the ϕ^8 truncation.

with the anomalous dimension given by equation (54). One more time, the system can be solved numerically, and the fixed points as well as their essential properties are summarized in the Table (11).

Interestingly, note that the line of fixed points has disappeared, that is not a surprise, because such line of fixed points are generally a pathology of the crude truncation. Among the fixed points listed in the table, only FP_5 , FP_6 and FP_{10} have $D > 0$.

- The fixed point FP_5 have seven irrelevant directions and one relevant direction in the UV, and seems to be an IR fixed point, whose irrelevant direction span an IR-critical surface with seven dimensions.
- The fixed point FP_6 have five relevant directions and three irrelevant directions in the UV. The relevant directions span an UV-multicritical surface of dimension five. The existence of a such sub-manifold is in accordance with the asymptotic safety of the theory.
- The fixed point FP_{10} have seven relevant directions and one irrelevant direction in the UV. It correspond to an UV fixed point whose relevant directions span a seven-dimensional UV-critical surface. One more time, the existence of a such manifold seems to be in accordance with a non-trivial asymptotically safe theory.

At this stage, it is not obvious to make contact with the fixed points obtained in the previous truncation. The standard way to highlight these relation is to consider truncation with higher and higher valence, and seek convergence of the fixed points. But in our case, the difficulty of a such computations is very improved by the non-locality of the interactions, and these conclusions have to

be confirmed by more finer analysis.

6 Discussion and conclusion

In this paper the renormalization group analysis is applied for just renormalizable T_5^6 TGFT model in the deep UV limit. Using the simplest approximation consisting in a truncation around the marginal interactions with respect to the perturbative power counting i.e. around the Gaussian fixed point, we have derived the flow equations for each couplings. Because we have focused our attention on the UV sector, the leading contributions to the flow equations provide to the melonic sector, a consideration which considerably simplify the computation of the flow equations. In a second time, using of the appropriate notion of canonical dimension in the UV, we have translated our flow equation in an autonomous system of differential equations, whose we have computed numerically the fixed points, as well as the behavior of the flow's trajectory around each of them.

We have find two type of fixed points. The IR fixed points, whose relevant directions in the IR span an IR-critical surface, a picture in favor of phase-transitions. This is supported, for two of these IR fixed points, by the negative value of their mass-parameter, and the fact that in their vicinity, the effective action turn to be a Ginsburg-Landau like equation for a ϕ^4 scalar complex theory, advocating a condensed phase transition interpretation. In opposition, the second type of fixed points are UV critical, and their relevant directions in the UV span critical surface with dimensions higher or equal to two, a picture in accordance with a well-defined and non-trivial behavior in the UV for asymptotically safe theories. In all the case, we observe that anomalous dimensions enhanced or weaken the UV-power counting for relevant operators with respect to the perturbative power-counting, a phenomena which seems to indicate a break-down of our crude truncation in dome domains of the phase-space. Moreover, the presence of pathological effects as a line of fixed point seems to confirm this suspicions, as well as its disappearance in a higher-truncation, while our conclusions about asymptotic safety and IR fixed points remains true. The connection between the new fixed point and these ones obtained in the first truncation remains however unclear at this stage without more control over the approximation procedure.

Finally, note that in the complementary IR regime, the flow equations receive non-melonic contributions. This is due to the fact that, for a very small cut-off, the sums take values 0 or 1. As pointed in [41], the appropriate rescalling is provided by the standard power counting, and the flow equation turn as well to be an autonomous system, which can be solved numerically. However, we have to keep in mind that our model is defined on a compact manifold ($U(1)^6$ in the referenced paper, $U(1)^5$ in our case). Then, no-phase transition can be occurs, and all the non-Gaussian fixed point reach to the Gaussian one when the cut-off tend to 0, except if the radius of the circles tend simultaneously to the infinity.

7 Acknowledgments

D O Samary research at Max-Planck Institute is supported by the Alexander von Humboldt foundation.

References

- [1] R. Gurau and J. P. Ryan, “Colored Tensor Models - a review,” SIGMA **8**, 020 (2012) doi:10.3842/SIGMA.2012.020 [arXiv:1109.4812 [hep-th]].
- [2] R. Gurau, “Colored Group Field Theory,” Commun. Math. Phys. **304**, 69 (2011) doi:10.1007/s00220-011-1226-9 [arXiv:0907.2582 [hep-th]].
- [3] V. Rivasseau, “Constructive Tensor Field Theory,” arXiv:1603.07312 [math-ph].
- [4] V. Rivasseau, “Random Tensors and Quantum Gravity,” arXiv:1603.07278 [math-ph].
- [5] V. Rivasseau, “The Tensor Theory Space,” Fortsch. Phys. **62**, 835 (2014) doi:10.1002/prop.201400057 [arXiv:1407.0284 [hep-th]].
- [6] V. Rivasseau, “The Tensor Track, III,” Fortsch. Phys. **62**, 81 (2014) doi:10.1002/prop.201300032 [arXiv:1311.1461 [hep-th]].
- [7] V. Rivasseau, “The Tensor Track, IV,” arXiv:1604.07860 [hep-th].
- [8] V. Rivasseau, “The Tensor Track: an Update,” arXiv:1209.5284 [hep-th].
- [9] V. Bonzom, R. Gurau, J. P. Ryan and A. Tanasa, “The double scaling limit of random tensor models,” JHEP **1409**, 051 (2014) doi:10.1007/JHEP09(2014)051 [arXiv:1404.7517 [hep-th]].
- [10] S. Dartois, R. Gurau and V. Rivasseau, “Double Scaling in Tensor Models with a Quartic Interaction,” JHEP **1309**, 088 (2013) doi:10.1007/JHEP09(2013)088 [arXiv:1307.5281 [hep-th]].
- [11] T. Delepouve and R. Gurau, “Phase Transition in Tensor Models,” JHEP **1506**, 178 (2015) doi:10.1007/JHEP06(2015)178 [arXiv:1504.05745 [hep-th]].
- [12] D. Benedetti and R. Gurau, “Symmetry breaking in tensor models,” Phys. Rev. D **92**, no. 10, 104041 (2015) doi:10.1103/PhysRevD.92.104041 [arXiv:1506.08542 [hep-th]].
- [13] A. Baratin, S. Carrozza, D. Oriti, J. Ryan and M. Smerlak, “Melonic phase transition in group field theory,” Lett. Math. Phys. **104**, 1003 (2014) doi:10.1007/s11005-014-0699-9 [arXiv:1307.5026 [hep-th]].
- [14] D. Benedetti and R. Gurau, “Phase Transition in Dually Weighted Colored Tensor Models,” Nucl. Phys. B **855**, 420 (2012) doi:10.1016/j.nuclphysb.2011.10.015 [arXiv:1108.5389 [hep-th]].
- [15] V. Bonzom, R. Gurau, A. Riello and V. Rivasseau, “Critical behavior of colored tensor models in the large N limit,” Nucl. Phys. B **853**, 174 (2011) doi:10.1016/j.nuclphysb.2011.07.022 [arXiv:1105.3122 [hep-th]].
- [16] R. Gurau, “The complete $1/N$ expansion of colored tensor models in arbitrary dimension,” Annales Henri Poincare **13**, 399 (2012) doi:10.1007/s00023-011-0118-z [arXiv:1102.5759 [gr-qc]].
- [17] R. Gurau, “The $1/N$ expansion of colored tensor models,” Annales Henri Poincare **12**, 829 (2011) doi:10.1007/s00023-011-0101-8 [arXiv:1011.2726 [gr-qc]].

- [18] D. Oriti, J. P. Ryan and J. Thurigen, “Group field theories for all loop quantum gravity,” *New J. Phys.* **17**, no. 2, 023042 (2015) doi:10.1088/1367-2630/17/2/023042 [arXiv:1409.3150 [gr-qc]].
- [19] D. Oriti, “A Quantum field theory of simplicial geometry and the emergence of spacetime,” *J. Phys. Conf. Ser.* **67**, 012052 (2007) doi:10.1088/1742-6596/67/1/012052 [hep-th/0612301].
- [20] C. Rovelli, “Quantum gravity,” *Scholarpedia* **3**, no. 5, 7117 (2008). doi:10.4249/scholarpedia.7117
- [21] C. Rovelli, “Zakopane lectures on loop gravity,” *PoS QGQGS 2011*, 003 (2011) [arXiv:1102.3660 [gr-qc]].
- [22] C. Rovelli, “Loop quantum gravity: the first twenty five years,” *Class. Quant. Grav.* **28**, 153002 (2011) doi:10.1088/0264-9381/28/15/153002 [arXiv:1012.4707 [gr-qc]].
- [23] S. Carrozza, D. Oriti and V. Rivasseau, “Renormalization of Tensorial Group Field Theories: Abelian U(1) Models in Four Dimensions,” *Commun. Math. Phys.* **327**, 603 (2014) doi:10.1007/s00220-014-1954-8 [arXiv:1207.6734 [hep-th]].
- [24] S. Carrozza, “Tensorial methods and renormalization in Group Field Theories,” doi:10.1007/978-3-319-05867-2 arXiv:1310.3736 [hep-th].
- [25] S. Carrozza, D. Oriti and V. Rivasseau, “Renormalization of a SU(2) Tensorial Group Field Theory in Three Dimensions,” *Commun. Math. Phys.* **330**, 581 (2014) doi:10.1007/s00220-014-1928-x [arXiv:1303.6772 [hep-th]].
- [26] J. Ben Geloun, “Renormalizable Models in Rank $d \geq 2$ Tensorial Group Field Theory,” *Commun. Math. Phys.* **332**, 117 (2014) doi:10.1007/s00220-014-2142-6 [arXiv:1306.1201 [hep-th]].
- [27] V. Lahoche and D. Oriti, “Renormalization of a tensorial field theory on the homogeneous space SU(2)/U(1),” arXiv:1506.08393 [hep-th].
- [28] V. Lahoche, D. Oriti and V. Rivasseau, “Renormalization of an Abelian Tensor Group Field Theory: Solution at Leading Order,” *JHEP* **1504**, 095 (2015) doi:10.1007/JHEP04(2015)095 [arXiv:1501.02086 [hep-th]].
- [29] J. Ben Geloun and E. R. Livine, “Some classes of renormalizable tensor models,” *J. Math. Phys.* **54**, 082303 (2013) doi:10.1063/1.4818797 [arXiv:1207.0416 [hep-th]].
- [30] D. Ousmane Samary and F. Vignes-Tourneret, “Just Renormalizable TGFT’s on $U(1)^d$ with Gauge Invariance,” *Commun. Math. Phys.* **329**, 545 (2014) doi:10.1007/s00220-014-1930-3 [arXiv:1211.2618 [hep-th]].
- [31] J. Ben Geloun and V. Rivasseau, “A Renormalizable 4-Dimensional Tensor Field Theory,” *Commun. Math. Phys.* **318**, 69 (2013) doi:10.1007/s00220-012-1549-1 [arXiv:1111.4997 [hep-th]].
- [32] J. Ben Geloun and V. Bonzom, “Radiative corrections in the Boulatov-Ooguri tensor model: The 2-point function,” *Int. J. Theor. Phys.* **50**, 2819 (2011) doi:10.1007/s10773-011-0782-2 [arXiv:1101.4294 [hep-th]].

- [33] J. Ben Geloun and D. O. Samary, “3D Tensor Field Theory: Renormalization and One-loop β -functions,” *Annales Henri Poincaré* **14**, 1599 (2013) doi:10.1007/s00023-012-0225-5 [arXiv:1201.0176 [hep-th]].
- [34] J. Ben Geloun, “Two and four-loop β -functions of rank 4 renormalizable tensor field theories,” *Class. Quant. Grav.* **29**, 235011 (2012) doi:10.1088/0264-9381/29/23/235011 [arXiv:1205.5513 [hep-th]].
- [35] V. Rivasseau, “Why are tensor field theories asymptotically free?,” *Europhys. Lett.* **111**, no. 6, 60011 (2015) doi:10.1209/0295-5075/111/60011 [arXiv:1507.04190 [hep-th]].
- [36] R. C. Avohou, V. Rivasseau and A. Tanasa, “Renormalization and Hopf algebraic structure of the five-dimensional quartic tensor field theory,” *J. Phys. A* **48**, no. 48, 485204 (2015) doi:10.1088/1751-8113/48/48/485204 [arXiv:1507.03548 [math-ph]].
- [37] S. Carrozza, “Discrete Renormalization Group for SU(2) Tensorial Group Field Theory,” *Ann. Inst. Henri Poincaré Comb. Phys. Interact.* **2** (2015), 49-112 doi:10.4171/AIHPD/15 [arXiv:1407.4615 [hep-th]].
- [38] D. Ousmane Samary, “Beta functions of $U(1)^d$ gauge invariant just renormalizable tensor models,” *Phys. Rev. D* **88**, no. 10, 105003 (2013) doi:10.1103/PhysRevD.88.105003 [arXiv:1303.7256 [hep-th]].
- [39] J. B. Geloun, R. Martini and D. Oriti, “Functional Renormalisation Group analysis of Tensorial Group Field Theories on \mathbb{R}^d ,” arXiv:1601.08211 [hep-th].
- [40] J. B. Geloun, R. Martini and D. Oriti, “Functional Renormalization Group analysis of a Tensorial Group Field Theory on \mathbb{R}^3 ,” *Europhys. Lett.* **112**, no. 3, 31001 (2015) doi:10.1209/0295-5075/112/31001 [arXiv:1508.01855 [hep-th]].
- [41] D. Benedetti and V. Lahoche, “Functional Renormalization Group Approach for Tensorial Group Field Theory: A Rank-6 Model with Closure Constraint,” arXiv:1508.06384 [hep-th].
- [42] D. Benedetti, J. Ben Geloun and D. Oriti, “Functional Renormalisation Group Approach for Tensorial Group Field Theory: a Rank-3 Model,” *JHEP* **1503**, 084 (2015) doi:10.1007/JHEP03(2015)084 [arXiv:1411.3180 [hep-th]].
- [43] J. B. Geloun and T. A. Koslowski, “Nontrivial UV behavior of rank-4 tensor field models for quantum gravity,” arXiv:1606.04044 [gr-qc].
- [44] P. Donà, A. Eichhorn, P. Labus and R. Percacci, “Asymptotic safety in an interacting system of gravity and scalar matter,” *Phys. Rev. D* **93**, no. 4, 044049 (2016) doi:10.1103/PhysRevD.93.044049 [arXiv:1512.01589 [gr-qc]].
- [45] P. Donà, A. Eichhorn and R. Percacci, “Consistency of matter models with asymptotically safe quantum gravity,” *Can. J. Phys.* **93**, no. 9, 988 (2015) doi:10.1139/cjp-2014-0574 [arXiv:1410.4411 [gr-qc]].
- [46] A. Eichhorn and T. Koslowski, “Continuum limit in matrix models for quantum gravity from the Functional Renormalization Group,” *Phys. Rev. D* **88**, 084016 (2013) doi:10.1103/PhysRevD.88.084016 [arXiv:1309.1690 [gr-qc]].

- [47] K. G. Wilson, “Renormalization group and critical phenomena. 1. Renormalization group and the Kadanoff scaling picture,” *Phys. Rev. B* **4**, 3174 (1971). doi:10.1103/PhysRevB.4.3174
- [48] K. G. Wilson, “Renormalization group and critical phenomena. 2. Phase space cell analysis of critical behavior,” *Phys. Rev. B* **4**, 3184 (1971). doi:10.1103/PhysRevB.4.3184
- [49] C. Wetterich, “Average Action and the Renormalization Group Equations,” *Nucl. Phys. B* **352**, 529 (1991). doi:10.1016/0550-3213(91)90099-J
- [50] N. Tetradis and D. F. Litim, “Analytical solutions of exact renormalization group equations,” *Nucl. Phys. B* **464**, 492 (1996) doi:10.1016/0550-3213(95)00642-7 [hep-th/9512073].
- [51] D. F. Litim, “Optimization of the exact renormalization group,” *Phys. Lett. B* **486**, 92 (2000) doi:10.1016/S0370-2693(00)00748-6 [hep-th/0005245].
- [52] R. Percacci, “Asymptotic Safety”, arXiv: 0709.3851, [hep-th], 2008.