

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-1-(3-Ethoxy-2-hydroxybenzylidene)-thiosemicarbazideAmir Adabi Ardakani,^{a*} Hadi Kargar,^b Reza Kia^{c,d} and Muhammad Nawaz Tahir^{e*}

^aArdakan Branch, Islamic Azad University, Ardakan, Iran, ^bDepartment of Chemistry, Payame Noor University, PO BOX 19395-3697 Tehran, I. R. of IRAN, ^cX-ray Crystallography Lab., Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran, ^dDepartment of Chemistry, Science and Research Branch, Islamic Azad University, Tehran, Iran, and ^eDepartment of Physics, University of Sargodha, Punjab, Pakistan
Correspondence e-mail: A.Adabi@iauardakan.ac.ir, dmntahir_uos@yahoo.com


Received 27 December 2011; accepted 5 January 2012

Key indicators: single-crystal X-ray study; $T = 291$ K; mean $\sigma(\text{C}-\text{C}) = 0.007$ Å; R factor = 0.056; wR factor = 0.119; data-to-parameter ratio = 18.5.

The title compound, $\text{C}_{10}\text{H}_{13}\text{N}_3\text{O}_2\text{S}$, crystallizes with two independent molecules (*A* and *B*) in the asymmetric unit. In the crystal, the *A* and *B* molecules are linked *via* pairs of $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{S}$ hydrogen bonds, forming dimers with $R_2^2(14)$ and $R_2^2(6)$ ring motifs. These dimers are linked *via* a pair of $\text{N}-\text{H}\cdots\text{S}$ hydrogen bonds with an $R_2^2(8)$ ring motif, forming chains propagating along the *c*-axis direction. The crystal was refined as an inversion twin with a final BASF ratio of 0.54 (11):0.46 (11).

Related literature

For standard bond lengths, see: Allen *et al.* (1987). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For background to thiosemicarbazones in coordination chemistry, see: Casas *et al.* (2000). For their biological applications, see: for example, Maccioni *et al.* (2003); Ferrari *et al.* (2000). For related structures, see: Kargar *et al.* (2010a,b).



Experimental

Crystal data

$\text{C}_{10}\text{H}_{13}\text{N}_3\text{O}_2\text{S}$
 $M_r = 239.29$
 Monoclinic, $P2_1$
 $a = 6.0728$ (3) Å
 $b = 16.1595$ (8) Å
 $c = 12.8490$ (6) Å
 $\beta = 90.238$ (3)°
 $V = 1260.91$ (11) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.25$ mm⁻¹
 $T = 291$ K
 $0.24 \times 0.14 \times 0.08$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.800$, $T_{\max} = 0.926$
 12062 measured reflections
 5428 independent reflections
 2303 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.075$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$
 $wR(F^2) = 0.119$
 $S = 0.92$
 5428 reflections
 293 parameters
 1 restraint
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³
 Absolute structure: Flack (1983),
 2232 Friedel pairs
 Flack parameter: 0.54 (11)

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> -H... <i>A</i>	<i>D</i> -H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> -H... <i>A</i>
O1-H1O...S2 ⁱ	0.83	2.53	3.180 (4)	135
O3-H3O...S1 ⁱⁱ	0.83	2.43	3.143 (4)	145
N2-H2N...O3 ⁱ	0.90	2.20	2.954 (6)	142
N5-H5N...O1 ⁱⁱ	0.87	2.17	3.009 (5)	160
N3-H3NB...S2 ⁱⁱⁱ	0.90	2.53	3.403 (4)	166
N6-H6NB...S1 ^{iv}	0.88	2.55	3.398 (5)	161

Symmetry codes: (i) $x + 1, y, z$; (ii) $x - 1, y, z$; (iii) $x + 1, y, z - 1$; (iv) $x - 1, y, z + 1$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

AAA thanks the Islamic Azad University, Ardakan Branch (this paper was extracted from the research project). HK thanks PNU for financial support. MNT thanks Sargodha University for research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2360).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
 Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
 Casas, J. S., Garcia-Tasende, M. S. & Sordo, J. (2000). *Coord. Chem. Rev.* **209**, 197–261.
 Ferrari, M. B., Capacchi, S., Reffo, G., Pelosi, G., Tarasconi, P., Albertini, R., Pinelli, S. & Lunghi, P. (2000). *J. Inorg. Biochem.* **81**, 89–97.
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.

Kargar, H., Kia, R., Akkurt, M. & Büyükgüngör, O. (2010*a*). *Acta Cryst.* **E66**, o2999.
Kargar, H., Kia, R., Akkurt, M. & Büyükgüngör, O. (2010*b*). *Acta Cryst.* **E66**, o2982.

Maccioni, E., Cardia, M. C., Distinto, S., Bonsignore, L. & De Logu, A. (2003). *Farmaco* **58**, 951–959.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

supporting information

Acta Cryst. (2012). E68, o340–o341 [doi:10.1107/S1600536812000487]

(E)-1-(3-Ethoxy-2-hydroxybenzylidene)thiosemicarbazide

Amir Adabi Ardakani, Hadi Kargar, Reza Kia and Muhammad Nawaz Tahir

S1. Comment

Thiosemicarbazones constitute an important class of N,S donor ligands due to their propensity to react with a wide range of metals (Casas *et al.*, 2000). Thiosemicarbazones exhibit various biological activities and have therefore attracted considerable pharmaceutical interest (Maccioni *et al.*, 2003; Ferrari *et al.*, 2000). Herein, we report on the crystal structure of the new title thiosemicarbazone compound.

The title compound crystallized with two independent molecules (A and B) in the asymmetric unit, Fig. 1. The bond lengths (Allen *et al.*, 1987) and angles are within the normal ranges and are comparable to those observed for related structures (Kargar *et al.*, 2010*a,b*).

In the crystal, the A and B molecules are linked *via* pairs of N-H \cdots O and O-H \cdots S hydrogen bonds (Table 1 and Fig. 2) to form dimers, with R²₂(14) and R²₂(6) ring motifs (Bernstein *et al.*, 1995). These dimers are further linked *via* a pair of N-H \cdots S hydrogen bonds, with an R²₂(8) ring motif, to form chains that extend in direction [0 0 1] (Table 1 and Fig. 2).


The crystal was refined as an inversion twin with a final refined BASF ratio of 0.54 (11)/0.46 (11) for 2232 Friedel pairs.

S2. Experimental


A mixture of 3-ethoxysalicylaldehyde (0.01 mol) and hydrazinecarbothioamide (0.01 mol) in 20 ml of ethanol was refluxed for about 2 h. On cooling, the solid separated was filtered and recrystallized from ethanol. Colourless plate-like crystals of the title compound, suitable for X-ray diffraction, were obtained by slow evaporation of a solution in ethanol.

S3. Refinement

O- and N-bound H atoms were located in a difference Fourier map and were initially refined with the O-H and N-H distances restrained to 0.82 (2) and 0.86 (2) Å, respectively. In the final cycles of refinement they were constrained to ride on their parent atoms with U_{iso}(H) = 1.5U_{eq}(O) and 1.2U_{eq}(N), respectively. The C-bound H-atoms were included in calculated positions and treated as riding atoms: C—H = 0.93, 0.96 and 0.97 Å for CH, CH₃ and CH₂ H-atoms, respectively, with U_{iso}(H) = k x U_{eq}(C), where k = 1.5 for CH₃ H-atoms, and k = 1.2 for all other H-atoms. The crystal was refined as an inversion twin with a final refined BASF ratio of 0.54 (11)/0.46 (11) for 2232 Friedel pairs.

**Figure 1**

The molecular structure of the two independent molecules of the title compound, showing 40% probability displacement ellipsoids and the atomic numbering.

**Figure 2**

A partial crystal packing diagram of the title compound, viewed down the *a*-axis, showing a one-dimensional extended chain along the *c*-axis formed *via* intermolecular O—H \cdots S, N—H \cdots O, and N—H \cdots S hydrogen bonds [dashed lines; see Table 1 for details; only the H atoms involved in these interactions are shown].

(E)-1-(3-Ethoxy-2-hydroxybenzylidene)thiosemicarbazide*Crystal data*C₁₀H₁₃N₃O₂S $M_r = 239.29$ Monoclinic, $P2_1$

Hall symbol: P 2yb

 $a = 6.0728$ (3) Å $b = 16.1595$ (8) Å $c = 12.8490$ (6) Å $\beta = 90.238$ (3)° $V = 1260.91$ (11) Å³ $Z = 4$ $F(000) = 504$ $D_x = 1.261$ Mg m⁻³Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2525 reflections

 $\theta = 2.5$ – 29.5 ° $\mu = 0.25$ mm⁻¹ $T = 291$ K

Plate, colourless

 $0.24 \times 0.14 \times 0.08$ mm*Data collection*Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 φ and ω scansAbsorption correction: multi-scan
(*SADABS*; Bruker, 2005) $T_{\min} = 0.800$, $T_{\max} = 0.926$

12062 measured reflections

5428 independent reflections

2303 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.075$ $\theta_{\text{max}} = 28.3$ °, $\theta_{\text{min}} = 1.6$ ° $h = -8 \rightarrow 7$ $k = -21 \rightarrow 19$ $l = -17 \rightarrow 17$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.056$ $wR(F^2) = 0.119$ $S = 0.92$

5428 reflections

293 parameters

1 restraint

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0341P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} < 0.001$ $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0087 (9)

Absolute structure: Flack (1983), 2232 Friedel
pairs

Absolute structure parameter: 0.54 (11)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2843 (8)	0.3987 (3)	1.0328 (4)	0.0372 (13)

C2	0.1042 (9)	0.3625 (4)	1.0844 (4)	0.0424 (15)
C3	-0.0613 (8)	0.3257 (4)	1.0275 (4)	0.0485 (16)
H3	-0.1804	0.3020	1.0617	0.058*
C4	-0.0523 (9)	0.3236 (4)	0.9203 (4)	0.0562 (16)
H4	-0.1637	0.2980	0.8824	0.067*
C5	0.1241 (9)	0.3600 (3)	0.8695 (4)	0.0478 (15)
H5	0.1287	0.3594	0.7972	0.057*
C6	0.2951 (8)	0.3976 (3)	0.9255 (4)	0.0381 (13)
C7	-0.0470 (9)	0.3298 (4)	1.2529 (4)	0.062 (2)
H7A	-0.0526	0.2708	1.2390	0.074*
H7B	-0.1900	0.3534	1.2367	0.074*
C8	0.0081 (10)	0.3445 (4)	1.3637 (4)	0.079 (2)
H8A	0.1469	0.3190	1.3798	0.118*
H8B	-0.1046	0.3211	1.4069	0.118*
H8C	0.0177	0.4029	1.3763	0.118*
C9	0.4833 (8)	0.4345 (3)	0.8726 (4)	0.0420 (14)
H9	0.5945	0.4587	0.9121	0.050*
C10	0.7230 (8)	0.4779 (3)	0.6336 (4)	0.0528 (17)
N1	0.4996 (7)	0.4345 (3)	0.7738 (3)	0.0486 (13)
N2	0.6925 (7)	0.4696 (3)	0.7372 (3)	0.0530 (14)
H2N	0.7890	0.4848	0.7866	0.064*
N3	0.5579 (7)	0.4547 (3)	0.5733 (3)	0.0652 (15)
H3NA	0.4296	0.4346	0.5911	0.078*
H3NB	0.5698	0.4611	0.5043	0.078*
O1	0.4483 (5)	0.4339 (2)	1.0882 (2)	0.0512 (12)
H1O	0.4206	0.4284	1.1511	0.077*
O2	0.1182 (6)	0.3677 (2)	1.1904 (3)	0.0550 (10)
S1	0.9596 (2)	0.51784 (12)	0.58949 (9)	0.0666 (5)
C11	0.2184 (8)	0.6114 (3)	0.8747 (4)	0.0383 (13)
C12	0.3909 (9)	0.6490 (4)	0.8233 (4)	0.0451 (15)
C13	0.5576 (9)	0.6839 (4)	0.8776 (5)	0.0503 (17)
H13	0.6739	0.7091	0.8430	0.060*
C14	0.5540 (9)	0.6818 (3)	0.9861 (5)	0.0528 (17)
H14	0.6696	0.7051	1.0236	0.063*
C15	0.3844 (9)	0.6461 (4)	1.0377 (4)	0.0427 (15)
H15	0.3829	0.6460	1.1101	0.051*
C16	0.2128 (8)	0.6097 (3)	0.9825 (4)	0.0353 (13)
C17	0.5280 (10)	0.6936 (4)	0.6567 (4)	0.071 (2)
H17A	0.6758	0.6724	0.6673	0.086*
H17B	0.5247	0.7512	0.6779	0.086*
C18	0.4616 (12)	0.6855 (4)	0.5436 (4)	0.102 (3)
H18A	0.4782	0.6290	0.5219	0.152*
H18B	0.5537	0.7203	0.5018	0.152*
H18C	0.3106	0.7020	0.5353	0.152*
C19	-0.2344 (8)	0.5427 (3)	1.2735 (4)	0.0509 (16)
C20	0.0220 (8)	0.5732 (3)	1.0337 (4)	0.0389 (15)
H20	-0.0841	0.5461	0.9940	0.047*
N4	-0.0013 (6)	0.5781 (3)	1.1322 (3)	0.0391 (11)

N5	−0.1924 (6)	0.5445 (3)	1.1704 (3)	0.0479 (13)
H5N	−0.2845	0.5167	1.1315	0.058*
N6	−0.0761 (8)	0.5723 (3)	1.3345 (3)	0.0693 (16)
H6NA	0.0549	0.5759	1.3077	0.083*
H6NB	−0.0989	0.5638	1.4014	0.083*
O3	0.0473 (5)	0.5768 (2)	0.8190 (2)	0.0515 (11)
H3O	0.0650	0.5791	0.7552	0.077*
O4	0.3715 (6)	0.6459 (2)	0.7167 (3)	0.0589 (11)
S2	−0.4755 (2)	0.50462 (12)	1.31643 (9)	0.0625 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.034 (3)	0.040 (4)	0.038 (3)	−0.002 (3)	0.004 (3)	0.000 (3)
C2	0.040 (3)	0.048 (4)	0.039 (3)	0.001 (3)	0.009 (3)	0.005 (3)
C3	0.036 (3)	0.050 (4)	0.059 (4)	−0.006 (3)	0.012 (3)	0.001 (3)
C4	0.046 (4)	0.059 (5)	0.064 (4)	−0.011 (3)	−0.003 (3)	0.001 (3)
C5	0.054 (4)	0.044 (4)	0.045 (3)	−0.003 (3)	−0.003 (3)	0.004 (3)
C6	0.037 (3)	0.041 (4)	0.037 (3)	0.002 (3)	−0.002 (2)	0.000 (3)
C7	0.061 (4)	0.057 (5)	0.068 (5)	−0.002 (3)	0.032 (3)	0.020 (4)
C8	0.090 (5)	0.096 (7)	0.051 (4)	−0.002 (4)	0.028 (4)	0.014 (4)
C9	0.038 (3)	0.049 (4)	0.040 (3)	−0.002 (3)	−0.001 (2)	0.000 (3)
C10	0.047 (3)	0.077 (5)	0.035 (3)	−0.010 (3)	0.002 (3)	−0.005 (3)
N1	0.048 (3)	0.069 (4)	0.029 (3)	−0.006 (3)	0.007 (2)	0.003 (3)
N2	0.048 (3)	0.084 (4)	0.027 (2)	−0.011 (3)	0.001 (2)	0.002 (2)
N3	0.062 (3)	0.105 (5)	0.029 (3)	−0.030 (3)	0.000 (2)	0.002 (3)
O1	0.049 (2)	0.077 (3)	0.027 (2)	−0.016 (2)	0.0058 (18)	−0.005 (2)
O2	0.060 (3)	0.068 (3)	0.037 (2)	−0.010 (2)	0.0164 (19)	0.004 (2)
S1	0.0488 (9)	0.1187 (16)	0.0325 (8)	−0.0175 (10)	0.0068 (7)	0.0012 (10)
C11	0.038 (3)	0.039 (4)	0.038 (3)	−0.001 (3)	0.004 (3)	−0.008 (3)
C12	0.043 (3)	0.053 (4)	0.039 (3)	−0.007 (3)	0.014 (3)	−0.001 (3)
C13	0.037 (4)	0.054 (4)	0.060 (4)	−0.005 (3)	0.015 (3)	0.009 (3)
C14	0.043 (3)	0.047 (4)	0.068 (4)	−0.008 (3)	0.007 (3)	0.001 (3)
C15	0.045 (4)	0.041 (4)	0.042 (3)	0.003 (3)	−0.007 (3)	−0.005 (3)
C16	0.034 (3)	0.034 (4)	0.037 (3)	0.002 (3)	0.003 (2)	0.001 (3)
C17	0.081 (4)	0.077 (5)	0.056 (4)	−0.013 (4)	0.038 (4)	0.009 (4)
C18	0.131 (6)	0.123 (7)	0.050 (4)	0.001 (5)	0.039 (4)	0.008 (4)
C19	0.048 (3)	0.072 (5)	0.032 (3)	0.000 (3)	0.002 (3)	0.003 (3)
C20	0.038 (3)	0.050 (4)	0.030 (3)	0.006 (3)	−0.002 (2)	−0.002 (3)
N4	0.032 (2)	0.051 (3)	0.034 (3)	−0.006 (2)	−0.0008 (19)	0.005 (2)
N5	0.041 (3)	0.072 (4)	0.031 (2)	−0.010 (2)	−0.0018 (19)	0.000 (2)
N6	0.063 (3)	0.113 (5)	0.032 (3)	−0.025 (3)	−0.003 (3)	−0.004 (3)
O3	0.051 (2)	0.070 (3)	0.034 (2)	−0.017 (2)	0.0040 (18)	−0.0036 (19)
O4	0.063 (3)	0.067 (3)	0.047 (2)	−0.013 (2)	0.019 (2)	0.001 (2)
S2	0.0463 (9)	0.1101 (15)	0.0313 (8)	−0.0101 (10)	0.0066 (6)	0.0014 (9)

Geometric parameters (Å, °)

C1—O1	1.347 (5)	C11—O3	1.378 (5)
C1—C6	1.381 (6)	C11—C12	1.382 (6)
C1—C2	1.409 (6)	C11—C16	1.385 (6)
C2—O2	1.367 (6)	C12—C13	1.351 (7)
C2—C3	1.375 (7)	C12—O4	1.375 (6)
C3—C4	1.379 (7)	C13—C14	1.395 (8)
C3—H3	0.9300	C13—H13	0.9300
C4—C5	1.387 (6)	C14—C15	1.356 (6)
C4—H4	0.9300	C14—H14	0.9300
C5—C6	1.399 (6)	C15—C16	1.389 (7)
C5—H5	0.9300	C15—H15	0.9300
C6—C9	1.460 (6)	C16—C20	1.459 (6)
C7—O2	1.426 (5)	C17—O4	1.447 (5)
C7—C8	1.480 (8)	C17—C18	1.513 (8)
C7—H7A	0.9700	C17—H17A	0.9700
C7—H7B	0.9700	C17—H17B	0.9700
C8—H8A	0.9600	C18—H18A	0.9600
C8—H8B	0.9600	C18—H18B	0.9600
C8—H8C	0.9600	C18—H18C	0.9600
C9—N1	1.273 (6)	C19—N6	1.327 (6)
C9—H9	0.9300	C19—N5	1.350 (5)
C10—N3	1.319 (6)	C19—S2	1.684 (5)
C10—N2	1.351 (5)	C20—N4	1.277 (6)
C10—S1	1.676 (5)	C20—H20	0.9300
N1—N2	1.386 (5)	N4—N5	1.374 (5)
N2—H2N	0.8964	N5—H5N	0.8736
N3—H3NA	0.8753	N6—H6NA	0.8703
N3—H3NB	0.8958	N6—H6NB	0.8816
O1—H1O	0.8316	O3—H3O	0.8286
O1—C1—C6	119.7 (4)	O3—C11—C12	120.1 (5)
O1—C1—C2	120.0 (5)	O3—C11—C16	119.3 (4)
C6—C1—C2	120.3 (5)	C12—C11—C16	120.6 (5)
O2—C2—C3	126.7 (5)	C13—C12—O4	126.2 (5)
O2—C2—C1	113.5 (5)	C13—C12—C11	120.3 (5)
C3—C2—C1	119.8 (5)	O4—C12—C11	113.5 (5)
C2—C3—C4	120.6 (5)	C12—C13—C14	119.5 (5)
C2—C3—H3	119.7	C12—C13—H13	120.3
C4—C3—H3	119.7	C14—C13—H13	120.3
C3—C4—C5	119.6 (5)	C15—C14—C13	120.9 (5)
C3—C4—H4	120.2	C15—C14—H14	119.5
C5—C4—H4	120.2	C13—C14—H14	119.5
C4—C5—C6	121.0 (5)	C14—C15—C16	120.0 (5)
C4—C5—H5	119.5	C14—C15—H15	120.0
C6—C5—H5	119.5	C16—C15—H15	120.0
C1—C6—C5	118.7 (5)	C11—C16—C15	118.7 (5)

C1—C6—C9	120.0 (5)	C11—C16—C20	118.8 (5)
C5—C6—C9	121.3 (5)	C15—C16—C20	122.4 (5)
O2—C7—C8	108.4 (5)	O4—C17—C18	107.0 (5)
O2—C7—H7A	110.0	O4—C17—H17A	110.3
C8—C7—H7A	110.0	C18—C17—H17A	110.3
O2—C7—H7B	110.0	O4—C17—H17B	110.3
C8—C7—H7B	110.0	C18—C17—H17B	110.3
H7A—C7—H7B	108.4	H17A—C17—H17B	108.6
C7—C8—H8A	109.5	C17—C18—H18A	109.5
C7—C8—H8B	109.5	C17—C18—H18B	109.5
H8A—C8—H8B	109.5	H18A—C18—H18B	109.5
C7—C8—H8C	109.5	C17—C18—H18C	109.5
H8A—C8—H8C	109.5	H18A—C18—H18C	109.5
H8B—C8—H8C	109.5	H18B—C18—H18C	109.5
N1—C9—C6	121.9 (5)	N6—C19—N5	115.6 (4)
N1—C9—H9	119.0	N6—C19—S2	124.6 (4)
C6—C9—H9	119.0	N5—C19—S2	119.8 (4)
N3—C10—N2	116.4 (4)	N4—C20—C16	120.9 (5)
N3—C10—S1	124.1 (4)	N4—C20—H20	119.5
N2—C10—S1	119.5 (4)	C16—C20—H20	119.5
C9—N1—N2	114.1 (5)	C20—N4—N5	115.3 (4)
C10—N2—N1	119.6 (4)	C19—N5—N4	121.5 (4)
C10—N2—H2N	125.4	C19—N5—H5N	115.4
N1—N2—H2N	115.0	N4—N5—H5N	122.6
C10—N3—H3NA	128.8	C19—N6—H6NA	116.8
C10—N3—H3NB	119.0	C19—N6—H6NB	113.8
H3NA—N3—H3NB	112.1	H6NA—N6—H6NB	122.9
C1—O1—H1O	108.4	C11—O3—H3O	113.3
C2—O2—C7	119.6 (4)	C12—O4—C17	117.2 (4)
O1—C1—C2—O2	0.0 (7)	O3—C11—C12—C13	179.2 (5)
C6—C1—C2—O2	179.3 (5)	C16—C11—C12—C13	0.7 (8)
O1—C1—C2—C3	-179.1 (5)	O3—C11—C12—O4	-1.1 (7)
C6—C1—C2—C3	0.1 (8)	C16—C11—C12—O4	-179.6 (4)
O2—C2—C3—C4	-178.7 (5)	O4—C12—C13—C14	-179.8 (5)
C1—C2—C3—C4	0.3 (8)	C11—C12—C13—C14	-0.1 (8)
C2—C3—C4—C5	-0.9 (8)	C12—C13—C14—C15	-0.8 (8)
C3—C4—C5—C6	1.1 (8)	C13—C14—C15—C16	1.2 (8)
O1—C1—C6—C5	179.3 (4)	O3—C11—C16—C15	-178.9 (5)
C2—C1—C6—C5	0.1 (7)	C12—C11—C16—C15	-0.3 (7)
O1—C1—C6—C9	-0.1 (7)	O3—C11—C16—C20	-1.7 (7)
C2—C1—C6—C9	-179.4 (5)	C12—C11—C16—C20	176.8 (5)
C4—C5—C6—C1	-0.7 (8)	C14—C15—C16—C11	-0.6 (8)
C4—C5—C6—C9	178.7 (5)	C14—C15—C16—C20	-177.6 (5)
C1—C6—C9—N1	179.8 (5)	C11—C16—C20—N4	-172.5 (5)
C5—C6—C9—N1	0.4 (8)	C15—C16—C20—N4	4.5 (8)
C6—C9—N1—N2	-177.9 (4)	C16—C20—N4—N5	177.3 (4)
N3—C10—N2—N1	2.9 (8)	N6—C19—N5—N4	-2.6 (8)

S1—C10—N2—N1	-178.2 (4)	S2—C19—N5—N4	177.4 (4)
C9—N1—N2—C10	-175.9 (5)	C20—N4—N5—C19	177.6 (5)
C3—C2—O2—C7	2.3 (8)	C13—C12—O4—C17	-8.7 (8)
C1—C2—O2—C7	-176.8 (5)	C11—C12—O4—C17	171.6 (5)
C8—C7—O2—C2	179.8 (5)	C18—C17—O4—C12	-175.6 (5)

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1—H1O...S2 ⁱ	0.83	2.53	3.180 (4)	135
O3—H3O...S1 ⁱⁱ	0.83	2.43	3.143 (4)	145
N2—H2N...O3 ⁱ	0.90	2.20	2.954 (6)	142
N5—H5N...O1 ⁱⁱ	0.87	2.17	3.009 (5)	160
N3—H3NB...S2 ⁱⁱⁱ	0.90	2.53	3.403 (4)	166
N6—H6NB...S1 ^{iv}	0.88	2.55	3.398 (5)	161

Symmetry codes: (i) $x+1, y, z$; (ii) $x-1, y, z$; (iii) $x+1, y, z-1$; (iv) $x-1, y, z+1$.