

Peeling or not peeling – is that the question ?*

Helmut Friedrich
 Max-Planck-Institut für Gravitationsphysik
 Am Mühlenberg 1
 14476 Golm, Germany

September 25, 2017

Abstract

The concepts of isolated self-gravitating system, asymptotic flatness and asymptotic simplicity are reconsidered, various related results are discussed and put into perspective, basic open questions are posed.

1 Introduction

The direct measurement of gravitational radiation [1] must be seen a triumph for experimental as well as for theoretical physics. In view of the graphs showing the impressive coincidence of the measured and the calculated radiation signals, one might think that everything comes to a conclusion and is understood now. But new questions will come up (see [26] for an example) and may require more precise statements. It will thus still be worthwhile to reconsider questions that have been left open in the theory of gravitational radiation. After giving an outline of the basic ideas and results concerning the underlying mathematical structure I shall discuss some of the remaining unresolved problems.

Following a gestation period of several years, involving many workers, the basic setting for the analysis of gravitational waves was proposed in the early 1960's by H. Bondi et al. [11], R. Sachs [65], E.T. Newman and R. Penrose [59]. It requires: The idealization of an *isolated self-gravitating system*, the analysis of solutions to Einstein's field equations that are *asymptotically flat in null directions*, the control of the evolution by Einstein's field equations on large scales, the control of the geometry on large scales, precise asymptotics

*Extended version of a talk given at the symposium honoring Piotr T. Chruściel on the occasion of his 60th birthday, 17th - 18th August 2017.

at space-like and null infinity, and the definition of physical concepts related to physical observations ‘far away from the system’.

Carving out the role of null and conformal geometry in the analysis of space-time structures in the large, R. Penrose [60] combined the various ideas in the elegant geometric concept of *asymptotic simplicity*, which characterizes the expected asymptotic behavior by the requirement that *the conformal structure be smoothly extendable across null infinity*.

The basic model is provided by Minkowski space $\hat{M} = \mathbb{R}^4$, $\hat{g} = -dt^2 + dr^2 + r^2 d\sigma^2$, given here in spatial polar coordinates with $d\sigma^2$ denoting the standard line element on \mathbb{S}^2 and coordinates $t \in \mathbb{R}$ and $r \geq 0$. Performing the coordinate transformation

$$t(\tau, \chi) = \frac{\sin \tau}{\cos \tau + \cos \chi}, \quad r(\tau, \chi) = \frac{\sin \chi}{\cos \tau + \cos \chi},$$

and rescaling with the conformal factor $\Omega = \cos \tau + \cos \chi = \frac{2}{\sqrt{1+(t-r)^2}\sqrt{1+(t+r)^2}}$, the *conformal metric* and its domain of definition are obtained in the form

$$g = \Omega^2 \hat{g} = -d\tau^2 + d\chi^2 + \sin^2 \chi d\sigma^2, \quad \hat{M} = \{\chi \geq 0, |\tau \pm \chi| < \pi\}.$$

This metric, the conformal factor, and the underlying manifold smoothly extend to yield *conformally compactified Minkowski space* with manifold

$$M = \{\chi \geq 0, |\tau \pm \chi| \leq \pi\} = \hat{M} \cup \mathcal{J}^\pm \cup i^0 \cup i^\pm.$$

The two components $\mathcal{J}^\pm = \{|\tau \pm \chi| = \pi\}$ of the *conformal boundary*, on which $\Omega = 0$, $d\Omega \neq 0$, represent *future and past null infinity*, they are generated by the future and past endpoints respectively acquired by the null geodesics. They are null hypersurfaces with respect to the *conformal metric* g .

The two points $i^\pm = \{\tau = \pm\pi, \chi = 0\}$, where $\Omega = 0$, $d\Omega = 0$, and $Hess_g \Omega = -g$, represent the future and past endpoints of the time-like geodesics and thus *future and past time-like infinity*. The space-like geodesics run in both directions towards *space-like infinity*, represented by the point $i^0 = \{\tau = 0, \chi = \pi\}$, where $\Omega = 0$, $d\Omega = 0$, and $Hess_g \Omega = g$. By adding this point, the Cauchy hypersurface $\{t = 0\} = \{\tau = 0, 0 \leq \chi < \pi\}$ of Minkowski space with the metric induced by g extends to the sphere \mathbb{S}^3 endowed with its standard metric.

The process of extending the *differential structure* and the *conformal structure* of a given space-time (\hat{M}, \hat{g}) to obtain a smooth (resp. C^k with k sufficiently large) conformal extension (M, g, Ω) with boundary \mathcal{J} so that $M = \hat{M} \cup \mathcal{J}$, $g = \Omega^2 \hat{g}$ on \hat{M} and $\Omega = 0, d\Omega \neq 0$ on \mathcal{J} as observed above was largely generalized in [60]. It was suggested that it applies to many solutions of Einstein’s field equations. In the case of solutions which satisfy Einstein’s vacuum field equations near \mathcal{J} it turned out that \mathcal{J} (consisting then in general of two components \mathcal{J}^\pm) is in fact a null hypersurface for the conformal metric g that represents (future and past) null infinity. In the situations considered by the authors mentioned above it gives the precise fall-off behaviour required in the asymptotic analysis and it largely simplifies the latter by the possibility to use, if \mathcal{J}^\pm is sufficiently smooth, local differential geometry instead of taking complicated limits.

In the 1960's , 1970's a large number of articles analyzed the geometrical and physical implications of the new picture and various concepts related to \mathcal{J}^\pm were discussed: The Bondi mass, the radiation field, the BMS group, . . . , see e.g. [51], [53] [61], [62] and the literature given there. However, while various concepts seemed to find a natural home in the new picture, it was not universally accepted by all workers in the field. Many competing and conflicting aspects were to be considered:

- Questions of mathematical generality,
- the definability and properties of physical concepts,
- the capacity to model the various physical situations of interest,
- sharpness of results and avoidance of redundancies,
- existence of solutions to the field equations with the desired asymptotics,
- numerical or analytical calculability of observation related quantities.

Without stating it explicitly in each case, the following discussion will touch, in one way or other, on most of these points.

2 Asymptotic smoothness and peeling

R. Penrose [61] gives an argument that for an asymptotically simple vacuum solution (\hat{M}, \hat{g}) with sufficiently smooth conformal extension (M, g, Ω) and so that $\mathcal{J}^+ \sim \mathbb{R} \times \mathbb{S}^2$ the conformal Weyl curvature $C^\mu{}_{\nu\lambda\rho}[g]$, given in coordinates adapted to g , vanishes on \mathcal{J}^+ . Because it is based on implicit assumptions on the smoothness of the conformal extension, it is difficult to asses its precise range of validity. It certainly works if (M, g, Ω) is of class C^4 but weakening this assumption is a very delicate matter (the argument as reconsidered in [44] starts from smoothness assumptions stronger than those in [61]). It will be seen below that the vanishing of $C^\mu{}_{\nu\lambda\rho}[g]$ at null infinity is in fact necessary for the smoothness of the conformal extension.

The vanishing of the Weyl tensor of the conformal metric at \mathcal{J}^\pm is thus related to the smoothness of the conformal fields. But what does that mean in terms of the physical fields ? It turns out that it is directly related to the very specific fall-off behaviour of the Weyl tensor $C^\mu{}_{\nu\lambda\rho}[\hat{g}]$ at null infinity which is known as *Sachs peeling*. This has been suggested by R. Sachs [64] to be characteristic for the asymptotic behaviour of self-gravitating isolated systems.

Assume that the conformal extension (M, g, Ω) with $M = \hat{M} \cup \mathcal{J}^+$ and $g = \Omega^2 \hat{g}$ on \hat{M} is of class C^k , $k \geq 3$. Let the function u on \hat{M} with $du \neq 0$ define a family of null hypersurfaces $\{u = \text{const.}\}$ in \hat{M} that approach \mathcal{J}^+ in space-like surfaces $\sim \mathbb{S}^2$ and let \hat{r} be an affine parameter along the future directed null geodesics generating the hypersurfaces $\{u = \text{const.}\}$ so that $\hat{r} \rightarrow \infty$ at \mathcal{J}^+ . Denote by $\{\hat{\kappa}_A\}_{A=0,1}$ a spin frame so that $\hat{e}_{AA'} = \hat{\kappa}_A \hat{\kappa}_{A'}$ is a \hat{g} -pseudo-orthonormal frame with $\hat{e}_{00'} = \text{grad}_{\hat{g}} u$ and assume that $\langle \hat{e}_{00'}, d\hat{r} \rangle = 1$.

It can then be shown that the conformal Weyl tensor $C^\mu{}_{\nu\lambda\rho}[g]$ (in coordinates adapted to g) vanishes on \mathcal{J}^+ if and only if the components $\hat{\Psi}_k$ of the conformal Weyl spinor corresponding to \hat{g} in the spin-frame $\hat{\kappa}_A$ have the Sachs peeling property, i.e. they satisfy, with integer powers of \hat{r}

$$\hat{\Psi}_k = \hat{\psi}_k \hat{r}^{k-5} + o(\hat{r}^{k-5}), \quad k = 0, 1, \dots, 4, \quad \text{as } \hat{r} \rightarrow \infty,$$

where the $\hat{\psi}_k$ can be regarded as function of class C^{k-3} on \mathcal{I}^+ (see [62] for details). We note that the function ψ_4 is interpreted as the *radiation field on \mathcal{I}^+* .

Not everybody shared the opinion that asymptotic simplicity encodes the fall-off behaviour of self-gravitating isolated systems in an appropriate way. Workers who studied equations of motions and tried to calculate the radiation escaping from the system found it difficult to verify Sachs peeling in their settings. This provoked doubts, questions and gave rise to heated discussions (see [32] and the references in [21]). In the following years many colleagues who followed the subject only by hearsay seemed to consider peeling as a dubious concept. In 1993 D. Christodoulou and S. Klainerman [16] still write: ‘... it remains questionable whether there exists any non-trivial solution of the field equations that satisfies the Penrose requirements. Indeed, his regularity assumptions translate into fall-off conditions of the curvature that may be too stringent and thus may fail to be satisfied by any solution that would allow gravitational waves’

Regardless of its geometric elegance, at the time the concept of asymptotic simplicity was just a proposal based on (profound) guess work. It then appeared natural to demand that *the fall-off behaviour of gravitational fields at null infinity should not be a matter of guesses but should be derived by achieving precise control on the evolution process*. But this leaves the question: *Which are appropriate situations from which the fields should evolve and what precisely is to be achieved ?* Moreover, far into the 1970’s only local in time results had been obtained in the general analysis of the Cauchy problem for Einstein’s field equations [12], [55].

The first results aiming at the long term evolution of gravitational fields appeared in the early 1980’s:

Following Y. Choquet-Bruhat’s [34] treatment of the Einstein’s vacuum equations as non-linear system of wave equations for the coefficients of the ‘physical’ metric, D. Christodoulou and N. O’Murchadha [15] analysed the *boost problem*. They showed that any asymptotically flat initial data for Einstein’s vacuum field equations have a development which includes complete space-like surfaces boosted relative to the initial surface. Future or past complete null geodesics, however, were not under control yet.

H. Friedrich [37], [38] studied representations of the Einstein equations in terms of the conformal fields g , Ω and derived fields, referred to as *conformal Einstein equations*, and introduced new *hyperbolic reductions*, aiming at precise and general existence results on solutions admitting smooth conformal boundaries.

To avoid permanently switching in the following from one setting to the other, I shall give up any chronological order (which can roughly be reconstructed from the references) and sketch in outline the second and then the first line of this research and the results obtained. I shall try to put them into perspective and pose questions which I think are more relevant than the one in the title.

To keep this article at a reasonable length I clearly have to ignore many contributions, for which I apologize. For the many omitted details of this highly technical subject I refer to the original articles, in particular to [8], [16], [58] with regard to the first approach and to [49] and the references given there in the case of the second one.

3 The hyperboloidal Cauchy problem

The first results which show that the smoothness of a conformal boundary can be preserved as a consequence of the field equations was obtained by solving the *hyperboloidal Cauchy problem* [39].

A space-like hypersurface S in the conformal extension of an asymptotically simple space-time (\hat{M}, \hat{g}) is referred to as a *hyperboloidal hypersurface* if it extends smoothly to \mathcal{J}^+ and is also space-like there. The set $\partial S = S \cap \mathcal{J}^+ \sim \mathbb{S}^2$ then defines a boundary of S at which $\Omega = 0$ and $d\Omega \neq 0$ (there could be inner boundaries but we are not interested here in those). If $\hat{h}_{\alpha\beta}$ and $\hat{\chi}_{\alpha\beta}$ denote the first and the second fundamental form induced by \hat{g} on $\hat{S} = S \cap \hat{M}$, then, in marked contrast to the behavior of the mean intrinsic curvature of asymptotically flat standard Cauchy data, which must approach zero at space-like infinity, it holds $|\hat{h}^{\alpha\beta} \hat{\chi}_{\alpha\beta}| \geq c = \text{const.} > 0$ near the end at infinity represented by $S \cap \mathcal{J}^+$.

Examples of such hypersurfaces in the conformally compactified Minkowski space considered above are given by the sets $\{\tau = \text{const.} \neq 0\}$ which comprise, in particular, the extension of the unit hyperbola $\{\eta_{\mu\nu} x^\mu x^\nu = -1, x^0 > 0\} \subset \{\tau = \frac{\pi}{2}\}$ that motivated the name. There are, of course, many more general examples. For later use we discuss a particular class of hyperboloidal hypersurfaces in the Schwarzschild space-time with metric

$$\hat{g} = - \left(1 - \frac{2m}{r}\right) dt^2 + \left(1 - \frac{2m}{r}\right)^{-1} dr^2 + r^2 d\sigma^2,$$

where $d\sigma^2$ denotes the standard metric on \mathbb{S}^2 and we assume $r > 2m \geq 0$. Since all structures will be spherically symmetric, angular coordinates are suppressed. In terms of the coordinates $w = t - r - 2m \log(r - 2m)$ and $\rho = 1/r$ it follows that $\Omega^2 \hat{g} = g$ with

$$\Omega = \rho, \quad g = -(1 - 2m\rho) \rho^2 dw^2 + 2 dw d\rho + d\sigma^2.$$

These fields extend smoothly to the set $\{\rho = 0, w \in \mathbb{R}\}$ which describes the future conformal boundary \mathcal{J}^+ of the Schwarzschild solution. The Cauchy hypersurface $\{t = 0, r > 2m\}$ is given in this representation by

$$w = w(\rho) \equiv -\frac{1}{\rho} - 2m \log\left(\frac{1}{\rho} - 2m\right), \quad 0 < \rho < \frac{1}{2m}.$$

Choosing ρ_0 with $0 < \rho_0 < \min(\frac{1}{3m}, \frac{1}{1+2m})$ and replacing $w(\rho)$ by the C^1 function $w_*(\rho)$ with $w_*(\rho) = w(\rho)$ for $\rho_0 \leq \rho < \frac{1}{2m}$ and $w_*(\rho) = \frac{\partial w}{\partial \rho}|_{\rho_0} (\rho - \rho_0) + w(\rho_0)$ for $0 \leq \rho < \rho_0$, one obtains a spherically symmetric C^1 hypersurface S' which is hyperboloidal. It intersects \mathcal{J}^+ in the same sphere as the outgoing null hypersurface $\{w = -\frac{\partial w}{\partial \rho}|_{\rho_0} \rho_0 + w(\rho_0)\}$ and approaches a Minkowskian hyperboloidal hypersurface as $m \rightarrow 0$. Choosing ρ_0 small enough and smoothing S' near the sphere $\{w = w(\rho_0), \rho = \rho_0\}$ while preserving its space-like nature we find:

For given $R > 2m$ there exist smooth hyperboloidal hypersurfaces in the Schwarzschild solution which have intersections with the Cauchy hypersurface $\{t = 0, 2m < r\}$ that comprise the set $\{t = 0, 2m < r \leq R\}$ and approach Minkowskian hyperboloidal hypersurfaces as $m \rightarrow 0$.

If the asymptotically simple space-time (\hat{M}, \hat{g}) considered above solves Einstein's vacuum field equations, the data $\hat{h}_{\alpha\beta}$ and $\hat{\chi}_{\alpha\beta}$ induced on the hyperboloidal hypersurface S satisfy the vacuum constraints on space-like hypersurfaces and have a specific fall-off behaviour at the boundary $S \cap \mathcal{J}^+$ which allows them to be conformally transformed and smoothly extended to $S \cap \mathcal{J}^+$ so as to yield the 1st and 2nd fundamental form induced by the smooth conformal metric g on S . We refer to data with these properties as *smooth hyperboloidal Cauchy data*. For the associated *hyperboloidal Cauchy problem* holds the following:

H. Friedrich [39]: *Smooth hyperboloidal Cauchy data develop into a solution to the vacuum equations which admits a smooth conformal extension \mathcal{J}^+ in the future of S whose null generators have past end points on the boundary of S .*

There is no 'smallness condition' required here and no restriction on the topology of S besides orientability and the existence of a boundary representing the asymptotic end. The 'life time' of the solution depends, of course, on the nature of the data. In general there may be no conformal gauge in which the null generators of \mathcal{J}^+ are future complete (see R. Geroch, G. T. Horowitz [52] for a notion of completeness of null infinity).

There is an important difference here with the formal expansion type studies of asymptotically simple solutions. *In that case asymptotic smoothness and thus peeling is put in by hand all along \mathcal{J}^+ . In the present case it is imposed on the initial data but is then preserved along \mathcal{J}^+ as a consequence of the field equations.*

3.1 Strong non-linear future stability

More can be said with further assumptions on the data.

Friedrich [40]: *The future development of a smooth Minkowskian hyperboloidal initial data set $(S^*, \hat{h}_{ab}^*, \hat{\chi}_{ab}^*)$ is strongly stable: Any smooth hyperboloidal vacuum initial data set $(S, \hat{h}_{ab}, \hat{\chi}_{ab})$ sufficiently close (in suitable Sobolev norm) to $(S^*, \hat{h}_{ab}^*, \hat{\chi}_{ab}^*)$ develops into a solution to Einstein's vacuum equations whose causal geodesics are future complete. Moreover, it admits a smooth conformal extension at future null infinity with conformal boundary \mathcal{J}^+ . The extension can be chosen so that \mathcal{J}^+ is generated by the past directed null geodesics which emanate from a regular point i^+ in the conformal extension and have past end point on the boundary ∂S .*

It is a remarkable property of the field equations that they force the null generators to meet, under the given assumptions, in a regular point i^+ that represents future-time-like infinity. The result generalizes to the Einstein-Maxwell-Mills equations [41] and to other Einstein-matter systems with conformally covariant matter fields.

I considered the study of hyperboloidal problems in the beginning as a preparation for the standard Cauchy problem, but it turned out to be as well suited for the numerical calculation of gravitational radiation fields at null infinity as the standard Cauchy problem. Peter Hübner, who pioneered the numerical studies based on the conformal field equations, calculated future complete solutions as the ones considered above, including the set $\mathcal{J}^+ \cup \{i^+\}$ and the radiation field induced on it [54]. For further information on the numerics of hyperboloidal initial value problem we refer to [35] and [63].

3.2 Existence of smooth hyperboloidal data

A. Andersson, P. Chruściel, H. Friedrich [5] studied the construction of hyperboloidal data $(S, \hat{h}_{ab}, \hat{\chi}_{ab})$ on a 3-manifold with boundary with second fundamental forms $\hat{\chi}_{\alpha\beta}$ satisfying $\hat{\chi}_{\alpha\beta} = \frac{\hat{\chi}}{3} \hat{h}_{\alpha\beta}$ on $\hat{S} = S \setminus \partial S$, the analogue of assuming time reflection symmetric data in the standard Cauchy problem. The momentum constraint and the assumed geometry require then $\hat{\chi} = \text{const.} \neq 0$ so that the free datum is given by the conformal class of the physical 3-metric $\hat{h}_{\alpha\beta}$.

Let $\omega \in C^\infty(S)$ be a boundary defining function so that $\omega > 0$ on \hat{S} and $\omega = 0, d\omega \neq 0$ on ∂S and let $h_{\alpha\beta}$ be a smooth Riemannian metric on S . The ansatz $\hat{h}_{\alpha\beta} = \phi^4 \omega^{-2} h_{\alpha\beta}$ with an unknown scalar function ϕ reduces the Hamiltonian constraint to a singular elliptic problem for ϕ :

$$R[\phi^4 \omega^{-2} h] = -\frac{2}{3} \hat{\chi}^2, \quad \phi \geq \phi_0 = \text{const.} > 0 \quad \text{on } S.$$

There exists a unique solution ϕ to this problem. It is smooth on \hat{S} but admits in general only a polyhomogeneous asymptotic expansion at ∂S in terms of the functions $\omega^i (\log \omega)^j$. The logarithmic terms vanish and the obtained hyperboloidal data are smooth if and only if one of the following equivalent conditions is satisfied:

- (i) *The trace free part of the second fundamental form induced by $h_{\alpha\beta}$ on ∂S vanishes.*
- (ii) *The conformal Weyl tensor $C^\mu{}_{\nu\lambda\rho}$ calculated from the data on S vanishes on ∂S .*
- (ii) *The ‘asymptotic shear’ of the null geodesic congruence approaching ∂S which defines the Cauchy horizon of the past Cauchy development of the solution determined by the data vanishes at ∂S .*

If the fields \hat{h}_{ab} and $\hat{\chi}_{ab}$ satisfy the constraints, they allow us to calculate the conformal Weyl tensor $C^\mu{}_{\nu\lambda\rho}[\hat{g}]$ on \hat{S} , where \hat{g} denotes the physical solution metric determined from these data. On \hat{S} it is equal to the Weyl tensor $C^\mu{}_{\nu\lambda\rho}[g]$, where $g = \Omega^2 \hat{g}$ with a suitable conformal factor Ω . If g extended smoothly to the set $\{\Omega = 0\}$ the tensor $C^\mu{}_{\nu\lambda\rho}[g]$ would also extend smoothly. That it should satisfy in fact $C^\mu{}_{\nu\lambda\rho}[\hat{g}] = C^\mu{}_{\nu\lambda\rho}[g] \rightarrow 0$ at ∂S is a non-trivial condition.

That the asymptotic shear vanishes on the conformal boundary of asymptotically simple vacuum solutions has been observed already by Penrose [61]. That this condition is decisive in the smoothness discussion for hyperboloidal data is again non-trivial.

Most important is the first condition which is given directly in terms of the free data. It shows that the latter only need to satisfy asymptotic condition at the boundary ∂S for the data to evolve into a space-time that admits a smooth conformal boundary in its future. These conditions are easily satisfied.

This result opened the way to the construction more general smooth hyperboloidal data. A. Andersson and P. Chruściel generalized the result in two ways [4]: The second fundamental form was only subject to the requirement $\hat{\chi} = \hat{\chi}_{ab} \hat{h}^{ab} = \text{const.} \neq 0$ and the free data were admitted that have a polyhomogeneous expansions at ∂S . With these conditions they could show:

The solutions to the constraints again admit polyhomogeneous expansion at ∂S . The non-vanishing of the conformal Weyl tensor at ∂S again contributes to the occurrence of logarithmic terms in the solutions to the elliptically reduced constraints. Conditions on the

free data can be given under which the hyperboloidal data extend smoothly to ∂S .

While the solutions arising from smooth data admit a smooth conformal extension across null infinity, the much more complicated behaviour near null infinity of solution space-times arising from general Chruściel-Andersson hyperboloidal data has not been analysed yet.

P. T. Chruściel, M. A. H. MacCallum and D. B. Singleton [21] studied general formal Bondi expansion admitting again asymptotic polyhomogeneous expansions. While some Bondi expansions admitting some logarithmic terms had been discussed before (see the references in [21]) they had not been analysed in a systematic way.

4 Asymptotically simple vacuum solutions

C. Cutler and R. Wald [27] managed to construct a parameter dependent family of smooth asymptotically flat standard Cauchy data for the Einstein-Maxwell equations on \mathbb{R}^3 that are isometric to Schwarzschild data in a neighbourhood of space-like infinity. As we have seen above, the developments in time of such data contain smooth hyperboloidal hypersurfaces which carry smooth hyperboloidal initial data. Since the standard Cauchy data constructed by the authors approach Minkowskian standard Cauchy data for suitable values of the parameter, the hyperboloidal initial data approach Minkowskian hyperboloidal initial data. Invoking the strong stability result discussed above, they were able to conclude:

There exist non-trivial solutions to the Einstein–Maxwell equations whose causal geodesics are complete and which admit smooth conformal extensions with complete null infinity \mathcal{J}^\pm and regular points i^\pm that represent past and future time-like infinity.

This was the first demonstration of the existence of non-trivial solutions to Einstein’s field equations with smooth and complete asymptotics. At the time the data used here looked rather contrived but ten years later they turned out to be special examples of a much larger class of similar data.

4.1 The Corvino gluing construction

J. Corvino introduces in [23] *a general technique which allows him to deform time reflection symmetric, asymptotically flat vacuum Cauchy data on \mathbb{R}^3 (say) outside a prescribed compact set so that they become isometric to Schwarzschild data in a neighbourhood of space-like infinity and satisfy the constraints everywhere.*

This is a most remarkable result. It gives an unexpected freedom to construct solutions to the constraints which are not accessible by earlier methods [6]. It also sheds new light on the role of the asymptotic ends at space-like infinity (see the discussion below).

P. Chruściel and E. Delay [18] and J. Corvino and R. Schoen) [25] generalize this result, showing that *general asymptotically flat vacuum Cauchy data can be modified outside a prescribed compact sets to become in some neighbourhood of space-like infinity isometric to the Schwarzschild or an other static solution in the time reflection symmetric case and isometric to Kerr or other stationary solutions in the other cases.*

These data have developments in time that are static or stationary near space-like infinity and thus have smooth conformal asymptotics there. Generalizing the construction of the hyperboloidal hypersurfaces for the Schwarzschild solution considered above, we conclude that these results also provide means *to deform given asymptotically flat data, without changing them on a given compact set, so as to become smooth hyperboloidal at their asymptotic end.*

Instead of using this detour via the evolution in time, P. T. Chruściel and E. Delay [20] directly use gluing techniques to show *the existence of a class of non-trivial data which are diffeomorphic to Schwarzschild-anti-de Sitter data outside some compact set and thus provide non-trivial hyperboloidal data* (for the relation between anti-de Sitter type data and hyperboloidal data see [56]).

Obviously, Corvino’s method was crying for an application along the lines of the Cutler–Wald idea, but as it stood his method did not allow him to produce data with arbitrarily small masses. In the following years P. Chruściel and E. Delay [17] and J. Corvino [24] managed, however, to show the existence of continuous families of smooth, non-trivial standard vacuum Cauchy data which are exactly static or stationary near space-like infinity and approximate Minkowskian standard Cauchy vacuum data. Evolving these data they thus obtained *families of smooth hyperboloidal data approximating Minkowskian hyperboloidal data.* Invoking again the strong stability result on Minkowskian hyperboloidal developments, they conclude:

There exists large classes of non-trivial solutions to the Einstein vacuum field equations with complete and smooth conformal extension \mathcal{J}^\pm at null infinity and regular points i^\pm at past and future time-like infinity.

Concerning the ‘largeness’ of the class it should be observed that while the data need to be close to Minkowskian data, the deformation techniques discussed above leave the original data unchanged on prescribed compact sets.

Because the points i^\pm for any of the solutions above are regular, the vanishing of the radiation field on \mathcal{J}^- or \mathcal{J}^+ would imply that the solutions were flat. If they have non-vanishing ADM mass, however, they have a non-vanishing conformal Weyl tensor. It follows that they have non-trivial radiation content. *Any doubts about the existence of radiative solution with smooth Penrose asymptotics have been put to rest by these results.*

Of course, being exactly static or stationary in a neighbourhood of space-like infinity (any such neighbourhood is of infinite spatial extent) is a strong assumption on the data and having the conformal boundary C^∞ instead of C^k , with some $k \geq 4$, is a strong requirement. It should be possible to weaken the assumptions and strengthen the result. It will be seen below that vacuum data which are *asymptotically static or stationary (up to sufficiently high order)* represent good candidates for this task.

5 Space-like infinity touching null infinity

The hyperboloidal Cauchy problem makes a clear distinction between asymptotically smooth and non-smooth data and the smooth data develop into solutions that admit

smooth conformal extensions. In the standard Cauchy problem the situation at the asymptotic end at space-like infinity is not so clear. Compactified Minkowski space is smoothly foliated by the slices $\{s = s_* = \text{const.}\}$. These are hyperboloidal if $s_* \neq 0$ while the slice $\{s = 0\}$ is asymptotically Euclidean, extending to the regular point i^0 that represents space-like infinity. When $m_{ADM} > 0$, conformal extensions in which space-like infinity is represented by a regular point do not exist and the transition from an asymptotically Euclidean slice to hyperboloidal slices is in general more complicated.

To understand possible obstructions to asymptotic smoothness arising in standard Cauchy problems we need to analyse in detail the structure of solutions in *a domain where space-like and null infinity come close to each other*. In the physical standard representation of the metric, in terms of which the structures referred to are at infinity, it is not clear what should be meant by this and the analysis requires complicated limits. On the other hand, conformally compactified Minkowski space, in which space-like infinity is represented by the one point i^0 is not a good guide if $m_{ADM} \neq 0$. In such a picture the rich structure discussed below would be compressed into one point and it would be impossible to analyse the field equations.

The requirement above acquires a concrete meaning in a setting introduced by H. Friedrich [42]. Space-like infinity is represented there by a cylinder $I =]-1, 1[\times \mathbb{S}^2$ that intersects an extended Cauchy hypersurface in the sphere $I^0 = \{0\} \times \mathbb{S}^2$ and touches the sets $\mathcal{J}^\pm = \{\Omega = 0, d\Omega \neq 0\}_\pm \sim \mathbb{R} \times \mathbb{S}^2$ at the *critical sets* $I^\pm = \{\pm 1\} \times \mathbb{S}^2$. All these sets, which define boundaries and edges of the physical space-time manifold \hat{M} , are given at a finite location in a certain type of coordinate systems. The setting and the gauge, including the coordinates, a g -orthonormal frame, and the conformal factor, are determined, apart from some conditions on the initial slice, by the field equations and the conformal structure of its solutions. The conformal factor and thus the location of the prospective hypersurfaces \mathcal{J}^\pm are known explicitly (see [43] for illustrations and [45] for explicit formulas in the case of Minkowski space). Because the gauge is based on conformal geodesics, it may be possible to generalize the analysis to more general initial data than the ones considered below. This requires a careful analysis, however, because the cylinder I , which is not a part of the physical manifold, is generated by limits of these curves.

No smallness conditions are needed here but to analyse the resulting, somewhat special, initial boundary value problem, it is convenient to require the initial data to be *asymptotically clean* in the sense that they are smooth and admit expansions in terms of powers of a radial coordinate \hat{r} with $\hat{r} \rightarrow \infty$ at space-like infinity. Prescribing ‘free data’ which satisfy this condition, S. Dain and H. Friedrich [30] analyse the constraints by standard methods. Besides the solutions of the desired form there are also some with terms $\hat{r}^k \log \hat{r}$, $k \in \mathbb{Z}$, related to a non-vanishing linear ADM momentum. As in [16], these are omitted in the following discussion. *In a suitable conformal scaling and coordinates on the initial slice in which the boundary I^0 is a sphere at a finite location, the data for the conformal field equations then extend smoothly to the set I^0 .*

The reduced conformal field equations are in this setting hyperbolic on $\hat{M} \cup I$ and, if the frame admits a continuous extension, also on null infinity \mathcal{J}^\pm . *However, the hyperbolicity degenerates at the critical sets I^\pm .* That a standard Cauchy problem is underlying the construction is reflected by the fact that *the boundary I is a total characteristic*: The

system of reduced equations reduces on I to a system of interior equations. As a consequence, it allows us to calculate a formal expansion of the space-time solution at all orders along the cylinder I by integrating the interior equations on I with initial data provided on I^0 by the data for the conformal field equations on the initial slice and their derivatives at I^0 . The main observations are:

Even when the data on the initial slice are smooth near I^0 , the solutions on I will in general not extend smoothly to I^\pm but develop a polyhomogeneous behaviour.

The logarithmic terms can be made to vanish by suitable choices of the Cauchy data.

If the setting is linearized at Minkowski space, so that the equations reduce essentially to the spin-2 equation, the polyhomogeneous behaviour at I^\pm spreads along the characteristics represented by \mathcal{J}^\pm [45].

The occurrence of logarithmic terms is thus not a problem of the setting but a consequence of the evolution equations and the structure of the data. We cannot expect the situation to be any better in the non-linear case.

If the data are static or stationary near space-like infinity the whole setting is as smooth near $I \cup I^\pm \cup \mathcal{J}^\pm$ near I^\pm , as one could wish [2], [46].

If the data are only asymptotically static or stationary at space-like infinity the integration on I gives no logarithmic terms at I^\pm .

In the case of time reflection symmetric data there is a certain amount of evidence that asymptotic staticity of the Cauchy data at space-like infinity is also *necessary* for the non-occurrence of logarithmic terms at I^\pm [42], [66]. Less is known about necessary conditions for the smoothness at I^\pm for Cauchy data with non-vanishing second fundamental form. It can be expected that *asymptotic stationarity* of the data at space-like infinity is sufficient for the existence of a smooth conformal boundary at null infinity.

In cases in which sufficient smoothness of the boundary can be established, the setting above allows us to perform very detailed calculations which relate the data near i^0 to the fields near I^\pm where the cylinder at space-like meets null infinity [50].

All the results above on the existence of ‘general’ solutions admitting smooth conformal extensions were obtained by using the conformal field equations for certain conformal fields $\Omega, g, \dots W^\mu{}_{\nu\lambda\rho}$ derived from the physical metric \hat{g} and a conformal factor Ω subject to certain gauge conditions. An important subsystem of the equations is given by the conformally covariant Bianchi or spin-2 equation which is given in the vacuum case by

$$\nabla_\mu W^\mu{}_{\nu\lambda\rho} = 0,$$

where ∇ denotes the connection defined by the conformal metric g and $W^\mu{}_{\nu\lambda\rho}$ the rescaled version of the conformal Weyl tensor $\hat{C}^\mu{}_{\nu\lambda\rho} = C^\mu{}_{\nu\lambda\rho}[\hat{g}]$ with

$$W^\mu{}_{\nu\lambda\rho} = \Omega^{-1} \hat{C}^\mu{}_{\nu\lambda\rho} \quad \text{on } \hat{M}.$$

This shows that trying to exploit the conformal properties of the Einstein equations in the most direct way has its advantages and its risks. The conformal field equations lead to complete and sharp results in situations in which the solution to be constructed will admit a smooth conformal extension. If peeling will not hold, however, the unknowns develop a singular behaviour as exemplified above by the situation at space-like infinity. This is reflected by the symmetric hyperbolic equations of first order for $W^\mu{}_{\nu\lambda\rho}$ which is

implied by the overdetermined system above in the gauges employed. The integrand in the energy estimates is given by certain components of the rescaled Bel-Robinson tensor, which become singular if peeling fails.

6 Non-linear stability of Minkowski space

The results on the global non-linear stability by D. Christodoulou and S. Klainerman [16], L. Bieri and N. Zipser [8], H. Lindblad and I. Rodnianski [58] and H. Lindblad [57] are less detailed as far as the precise asymptotic behaviour is concerned but much more complete than the results referred to above in that they start from fairly general asymptotically flat standard vacuum Cauchy data on \mathbb{R}^3 and control the completeness of causal geodesics in the the past and in the future. All results are obtained by working in terms of the physical metric \hat{g} . In [16] and [8] the Bianchi equation $\hat{\nabla}_\mu \hat{C}^\mu{}_{\nu\lambda\rho} = 0$ for the conformal Weyl tensor also plays an important role, its conformal properties are exploited indirectly and the properties of the Bel-Robinson tensor are used extensively to derive the relevant estimates. The work in [58] and [57] is based again on the representation of the Einstein equations as a system of wave equation obtained by imposing a harmonic gauge. Only some observations of relevance for our discussion will be presented here.

All authors assume the first and second fundamental form \hat{h}_{ab} and $\hat{\chi}_{ab}$ to be smooth, to be close to Minkowskian data in a well-defined sense, and to satisfy certain fall-off conditions near space-like infinity. In the case of [16] these read

$$\hat{h}_{ab} = (1 + 2m|x|^{-1}) \delta_{ab} + o_4(|x|^{-3/2}), \quad \hat{\chi}_{ab} = o_3(|x|^{-5/2}),$$

which implies the vanishing of the linear ADM momentum. This is not the case for the generalization given by Bieri in [8] which requires

$$\hat{h}_{ab} = \delta_{ab} + o_3(|x|^{-1/2}), \quad \hat{\chi}_{ab} = o_2(|x|^{-3/2}) \quad \text{as } |x| \rightarrow \infty.$$

In both cases x denotes a \mathbb{R}^3 -valued coordinate near the asymptotic end. It is shown:

The causal geodesics of the maximal globally hyperbolic solutions determined by these data are complete and their curvature tensors $\hat{C}^\mu{}_{\nu\lambda\rho}$ approach zero asymptotically in all directions.

For our discussion the rate at which that happens along null geodesics going out to null infinity is important. The constructions are based on level surfaces H_t and C_u of a time function t and a retarded time function u . The function $\hat{r} = \hat{r}(t, u) > 0$ is chosen to satisfy $\text{Vol}_{\hat{g}}(H_t \cap C_u) = 4\pi \hat{r}^2$ on the spherical intersections of the level surfaces. Along the null geodesics generating the null hypersurfaces C_u it holds then $\hat{r} \rightarrow \infty$ as they run out to future null infinity. Adapting the notation for the components of the conformal Weyl tensor used in the discussion of Sachs peeling it follows

$$|\hat{\Psi}_k| = O(\hat{r}^{-7/2}), \quad k = 0, 1, \quad |\hat{\Psi}_k| = O(\hat{r}^{k-5}), \quad k = 2, 3, 4,$$

in the case of [16] and in the case of the generalization given by Bieri in [8]

$$|\hat{\Psi}_k| = o(\hat{r}^{-5/2}), \quad k = 0, 1, 2, \quad |\hat{\Psi}_k| = O(\hat{r}^{k-5}), \quad k = 3, 4,$$

as $\hat{r} \rightarrow \infty$ along a fixed null generator of C_u . For $k = 3, 4$ the behaviour is thus similar to Sachs peeling while it deviates from it, *in the case of solutions for which these estimates are sharp*, for $k = 0, 1$.

What is the origin of these deviations ? The logarithmic terms observed at the critical set in the analysis of space-like infinity outlined above should contribute to them and the logarithmic term observed on the initial slice in the construction of clean initial data [30] should contribute to the stronger deviation in the case of [8]. It would be interesting to know which of the logarithmic terms observed in [4] and [21] can in fact occur in the solutions considered in [16] and [8]. In any case, if the data are specified in terms of weighted Sobolev spaces or if they are just required to be smooth and to satisfy the fall-off at space-like infinity indicated above, there is a large freedom to have, besides the logarithmic terms considered above, all kinds of other terms in the data which may spoil the smoothness of any conformal extension at null infinity at higher orders. The question then is whether these contributions are of any physical relevance or just add ‘noise’.

7 Approximative solutions

Analytical and numerical approximations are of utmost importance, because they allow us to relate measured data to theoretical results. Nevertheless, I shall only make some sketchy remarks about certain aspects related to my topic.

The analytical approximation theory designed to produce quantitative results on the radiation generated e.g. by the merger of black holes should have a counterpart formulated in terms of the Cauchy problem. It certainly would be most useful if more were known about this. The relation between the abstract and the approximative analytical understanding is not easily extracted from the literature, however, because the latter usually immediately intertwines general considerations with the technical details of the approximation method.

L. Blanchet and T. Damour note in their extensive work on approximation methods begun in [10] a difficulty to verify the peeling behaviour (see also the remarks by D. Christodoulou [13]). On the other hand they impose near space-like infinity conditions to exclude radiation coming in from the infinite past. As stated more explicitly by L. Blanchet [9] and T. Damour and B. Schmidt [31], it amounts to requiring the solutions to be stationary near space-like infinity. In these articles are also given arguments that the solution will then admit a smooth conformal extension near space-like infinity (a gap in the argument in [31] has been filled in by S. Dain [29]).

This raises the question why the asymptotic smoothness should be lost later in (retarded) time. Is this due to the eruptive behaviour of some matter system ? But why should it happen in the pure vacuum case, e.g. in the merger of black holes ?

Corvino’s result and its generalizations were not available at the time. But even if they had been known already, it is hardly conceivable that something like the gluing procedure could be realized in the context of an approximation method. Is the loss of asymptotic smoothness possibly just an artifact of the approximation method ?

To keep their numerical grids finite, most relativists who develop numerical 3+1 codes for the standard Cauchy problem to calculate binary black hole merger (say) wave forms

use cut-off procedures and essentially ignore space-like and null infinity, thus also the finer details of the asymptotic behaviour there. The radiation field, originally defined at null infinity, is calculated only approximately at a finite, somewhat arbitrary location. This cut-off deletes a neighbourhood of $\mathcal{J}^\pm \cup i^0$ which is of infinite extent as measured in terms of affine parameters on the outgoing null geodesics (see, however, the work by F. Beyer et al [7] and J. Frauendiener and J. Hennig [36] which takes first steps towards calculating entire solutions determined by asymptotically flat Cauchy data). Nevertheless, for the time being the results seem to be satisfactory.

Characteristic or hyperboloidal Cauchy problems with data prescribed on null or space-like hypersurfaces that extend to null infinity have also been solved numerically. The freedom in the choice of data near \mathcal{J}^+ is used to extend the data smoothly to \mathcal{J}^+ . The difference with the standard Cauchy problem is that the wave form extraction can be done at the well and uniquely defined hypersurface \mathcal{J}^+ .

In both case there is a large arbitrariness, however, in choosing the data near the asymptotic end of the initial slice.

8 A different type of approximation

The following two results are of particular interest in our discussion. P. Allen and I. Stavrov Allen [3] show:

Polyhomogeneous, asymptotically hyperbolic, constant mean curvature data of Andersson-Chruściel type for the vacuum Einstein equations can be approximated arbitrarily closely in certain Hölder norms by smooth hyperboloidal constant mean curvature vacuum data.

J. Corvino and R. Schoen [25] state a density result by which:

Asymptotically flat initial data for the vacuum Einstein equations on a three-manifold \hat{S} can be approximated by data on \hat{S} which agree with the original data inside a given compact domain, and are in a given end identical to that of a suitable Kerr slice (or identical to a member of some other admissible family of solutions) outside a large ball.

It should be noted that these approximations are controlled in terms of Sobolev norms which are weighted so that $\hat{h}_{ab} - \delta_{ab}$ and $\hat{\chi}_{ab}$ and the corresponding approximating data are consistent with the fall-off behaviour required for these fields in [16]. We state the results here without further details, because the answer to the question they provoke may well require the use of different function spaces.

Are the asymptotically simple vacuum solutions in a some sense dense in a set of asymptotically flat vacuum solutions as considered in the non-linear stability results above ?

If a definite answer could be given to this question the ‘mathematical exercises’ discussed in the previous sections could be brought to a conclusion. The proof of any such density results should completely clarify the situation. ‘Peeling or not peeling’ would just become a matter of deciding between technically more or less convenient representations. It may also show that the questions raised in our discussion of analytic and numerical approximations may be essentially harmless. The situation would be reminiscent of the introduction of L^2 -Hilbert spaces in the analysis of hyperbolic equations as an intermediary step towards obtaining existence results about smooth solutions. A positive result

should provide interesting information about the precise way in which solutions which are ‘rough’ at null infinity are approached by solutions which have smooth \mathcal{J}^+ and in which way concepts which are easily defined on smooth \mathcal{J}^+ ’s can be transferred (if at all) to concepts on the ‘rough’ future null infinity.

On the other hand, showing that such a density property cannot hold should explain in which sense rough asymptotics can be superior to smooth asymptotics. It should give information about physical systems of interest which cannot be modeled in the class of asymptotically simple solutions and tell us what precisely is lost if we restrict to asymptotically simple solutions. It should further give answers to the following questions.

The logarithmic terms at null infinity mentioned so far (and of some possibly not considered yet) come with certain coefficients. Which information is encoded in these coefficients? What is the physical information in the coefficient in the ‘free data’ underlying the construction of hyperboloidal or Cauchy data which must be set to zero to get rid of logarithmic terms? Doing so, does it lead to a loss of essential physical information? What is the role of the logarithmic term on the initial slice which is related to the linear ADM momentum? What is its effect on the structure of the radiation field or other quantities of physical interest on null infinity? If it can be shown that the logarithmic terms found at the critical sets are indeed related to the deviation of the data from being asymptotically stationary, could this be interpreted as saying that radiation coming in from the infinite past has to be excluded (up to some order) to achieve asymptotic smoothness (of a prescribed order)? Are there similar interpretations related to the other logarithmic terms?

9 Isolated systems as part of our cosmos

Most of the considerations above are related to the structure of the *asymptotic end at space-like infinity* in the standard and to the *asymptotic end at null infinity* in the hyperboloidal Cauchy problem. But in the ‘real world’ of our cosmos a system which we would like to see as part of a *self-gravitating isolated system* does not possess an asymptotic end at space-like or null infinity.

The best we can do is to consider an open, relative compact subset S' of a time-slice S of our cosmos with the Cauchy data d' induced on it so that its domain of dependence $D(S')$ contains the essential part of the process of interest but no further comparable objects. As a next step we could try to *attach an asymptotically flat or hyperboloidal end* smoothly to (S', d') ¹.

If S is assumed to be compact and all the matter fields are ignored, space-time engineering as suggested by P.T. Chrusciel, J. Isenberg, D. Pollack [19] allows us in fact to glue an asymptotically flat or hyperboloidal end to S . The resulting standard Cauchy data will

¹G.F.R. Ellis [33] suggested to abandon the asymptotically flat model and to introduce instead in an ad hoc fashion a spatially compact time-like hypersurface \mathcal{T} to cut off ‘the system of interest’ from the ambient universe. We refer the reader to [46] for a discussion of the difficulties of this idea and to [47] for the unresolved (possibly unresolvable) difficulties with the underlying initial boundary value problem for Einstein’s field equations.

contain, however, a huge number of other systems, which we wish to exclude, and there may even be something like a minimal surface close to the location of the gluing process. What we want is closer to the results of S. Czimek [28], who constructs asymptotically flat extensions of solutions to the vacuum constraints on a compact manifold with boundary that have vanishing mean extrinsic curvature. Since this is shown so far only to work for data close to Minkowskian data, the result, as it stands, excludes systems containing e.g. black holes. Another possibility might be to extend the set (S', d') , possibly after some modification close to its boundary, along the lines of R. Bartnik's parabolic constructions of constrained data discussed in [6]. All these studies suggest that it is not too far-fetched to assume, as we shall do, that (S', d') can be embedded isometrically into some asymptotically flat standard Cauchy set. Our earlier discussions then show that it can equally well be embedded into smooth hyperboloidal initial data sets.

Whatever one does, while suggested by static or stationary exact solutions and consistent with the field equations also under much more general assumptions, the asymptotically flat or hyperboloidal end is a just figment. It allows the field equations themselves to construct in a marvelously effective way a null infinity and the radiation signal to unfold while approaching that null infinity along outgoing null rays.

As we have seen, however, there exists a huge freedom to choose or modify asymptotic ends. The transition from one choice to another one may come with some *spurious radiation* which we may not accept as being associated with the system we have in mind (an observation well known to numerical relativists). But who says that some such radiation had not been provided already by the first choice? Those of us who spent much of their time working on the Kerr family or other real analytic stationary solutions may find it strange, but if we consider ends of class C^∞ or C^k we have to face the fact that *there simply does not exist 'the' correct asymptotic end*. We can only hope to optimize the situation in some sense.

This raises the question: *How to make the best use of the freedom to choose the asymptotically Euclidean or hyperboloidal end?* And more specifically: *To what extent do radiation fields and other physically relevant quantities defined at null infinity depend on the precise structure of the initial data near space-like (or null) infinity?*

Analysing these questions should give deeper insight into some physics because the answers will depend very much on the nature of the system we wish to model and the conclusions we want to draw. The inspiral, merger, and ring down of black holes can be expected to be accompanied by *strong, well-pronounced signals in the domain of dependence of some interior domain*. Reasonable changes near the ends at space-like or null infinity are likely to have little effect on these and we may well choose the end to be stationary. In scattering problems involving weak fields, however, we can, in principle, only toy with the data on $\mathcal{J}^- \cup i^-$ (see [14] and, for a full treatment of a neighbourhood of i^- , also [22], [48]). Then we have to watch how things develop at space-like infinity and what comes out at \mathcal{J}^+ . There are data on \mathcal{J}^- which make the end at space-like infinity stationary but whether they will be useful in this context depends very much on the questions to be answered.

Acknowledgements: I would like to thank L. Bieri, J. Corvino, P. Chruściel, D. Garfinkel,

J. Isenberg, R. Schoen, and R. Wald for discussions. Part of this work was supported by the programme on Geometry and Relativity at the Erwin Schroedinger Institute, Vienna. I wish to thank the Institute for support, hospitality and the opportunity to talk about the subject with various colleagues.

References

- [1] B. P. Abbott et al. Observation of gravitational waves from binary black hole merger. *Phys. Rev. Lett.* 116 (2016) 061102.
- [2] A. E. Aceña, J. A. Valiente Kroon. Conformal extensions for stationary space-times. *Class. Quantum Grav.* 28 (2011) 225023
- [3] P. T. Allen, I. Stavrov Allen. Smoothly compactifiable shear-free hyperboloidal data is dense in the physical topology. *Ann. Henri Poincaré* 18 (2017) 2789 - 2814.
- [4] L. Andersson, P. Chruściel. Solutions of the constraint equations in general relativity satisfying ‘hyperboloidal boundary conditions’. *Dissertationes Mathematicae*, Polska Akademia Nauk, Inst. Matem., Warszawa, 1996.
- [5] L. Andersson, P. Chruściel, H. Friedrich. On the existence of solutions to the Yamabe equation and the existence of smooth hyperboloidal initial data for Einstein’s field equations. *Commun. Math. Phys.* 149 (1992) 587 - 612.
- [6] R. Bartnik, J. Isenberg. The constraint equations. In: P. T. Chruściel, H. Friedrich (eds.): *The Einstein equations and the large scale behaviour of gravitational fields*. Birkhäuser, Basel, 2004.
- [7] F. Beyer, G. Doulis, J. Frauendiener, B. Whale. Numerical space-times near space-like and null infinity: The spin-2-system on Minkowski space. *Class. Quantum Grav.* 29 (2012) 245013..
- [8] L. Bieri, N. Zipser. *Extensions of the stability theorem of the Minkowski space in general relativity*. AMS, International Press, 2009.
- [9] L. Blanchet. Radiative gravitational fields in general relativity II. Asymptotic behaviour at future null infinity. *Proc. R. Soc. Lond. A* 409 (1987) 383–399.
- [10] L. Blanchet, T. Damour. Radiative gravitational fields in general relativity I. General structure of the field outside the source. *Phil. Trans. R. Soc. Lond. A* 320 (1986) 379–430.
- [11] H. Bondi, M. G. J. van der Burg, A. W. K. Metzner. Gravitational waves in general relativity VII. Waves from axi-symmetric isolated systems. *Proc. Roy. Soc A* 269 (1962) 21–52.
- [12] Choquet-Bruhat, Y., York, J. (1980) The Cauchy problem. In: Held, A. (Ed.) *General Relativity and Gravitation*, Vol. I. Plenum, New York, 99–172.

- [13] D. Christodoulou. The global initial value problem in general relativity. In: V. G. Gurzadyan et al (eds.): *The Ninth Marcel Grossmann Meeting on recent developments in theoretical and experimental general relativity, gravitation and relativistic field theories*. World Scientific, Hong Kong, 2002
- [14] D. Christodoulou. *The formation of black holes in general relativity*. European Mathematical Society Publishing House, Zürich, 2009.
- [15] D. Christodoulou, N. O’Murchadha. The boost problem in general relativity. *Commun. Math. Phys.* 80 (1981) 271 - 300.
- [16] D. Christodoulou, S. Klainerman. *The global non-linear stability of Minkowski space*. Princeton University Press, Princeton, 1993.
- [17] P. T. Chruściel, E. Delay. Existence of non-trivial, vacuum, asymptotically simple spacetimes. *Class. Quantum Grav.*, 19 (2002) L 71 - L 79. Erratum *Class. Quantum Grav.*, 19 (2002) 3389.
- [18] P. T. Chruściel, E. Delay. On mapping properties of the general relativistic constraints operator in weighted function spaces, with application. *Mém. Soc. Math. Fr. (N.S.)* 94 (2003).
- [19] P.T. Chrusciel. J. Isenberg, D. Pollack. Initial data engineering. *Commun. Math. Phys.* (2005) 29 - 42.
- [20] P. T. Chruściel, E. Delay. Gluing constructions for asymptotically hyperbolic manifolds with constant scalar curvature. *Communications in Analysis and Geometry* 17 (2009) 343 - 381.
- [21] P. T. Chruściel, M. A. H. MacCallum, D. B. Singleton. Gravitational waves in general relativity XIV. Bondi expansion and the ‘polyhomogeneity’ of \mathcal{J} . *Phil. Trans. R. Soc. Lond. A* 350 (1995) 113 - 141.
- [22] P. T. Chruściel, T.-T. Paetz. Solutions of the vacuum Einstein equations with initial data on past null infinity. *Class. Quantum Grav.* 30 (2013) 235037.
- [23] J. Corvino. Scalar curvature deformation and a gluing construction for the Einstein constraint equations. *Comm. Math. Phys.* 214 (2000) 137–189.
- [24] J. Corvino. On the existence and stability of the Penrose compactification. *Ann. Henri Poincaré* 8 (2007) 597 - 620.
- [25] J. Corvino, R. Schoen. On the Asymptotics for the Vacuum Einstein Constraint Equations. *J. Differential Geometry* 73 (2006) 185 - 217.
- [26] J. Creswell, S.v. Hausegger, A.D. Jackson, H. Liu, P. Naselsky. On the time lags of the LIGO signals. arXiv:1706.04191.
- [27] C. Cutler, R. M. Wald. Existence of radiating Einstein-Maxwell solutions which are C^∞ on all of \mathcal{J}^+ and \mathcal{J}^- . *Class. Quantum Grav.* 6 (1989) 453 - 466.
- [28] S. Czimek. An extension procedure for the constraint equations. arXiv:1609.08814

- [29] S. Dain. Initial data for stationary spacetimes near spacelike infinity. *Class. Quantum Grav.*, 18 (2001) 4329–4338.
- [30] S. Dain, H. Friedrich. Asymptotically Flat Initial Data with Prescribed Regularity. *Comm. Math. Phys.* 222 (2001) 569–609.
- [31] T. Damour, B. Schmidt. Reliability of perturbation theory in general relativity. *J. Math. Phys.* 32 (1990) 2441–2453.
- [32] J. Ehlers (ed). *Isolated systems in general relativity*. Proceedings of the international school of physics ‘Enrico Fermi’, course LXVII, Varenna 1976. North-Holland, Amsterdam, 1979.
- [33] G. F. R. Ellis. Relativistic Cosmology: Its Nature, Aims, and Problems, In: B. Bertotti et. al (eds.) *General Relativity and Gravitation*. Reidel, Dordrecht, 1984.
- [34] Y. Fourès-Bruhat. Théorème d’existence pour certains systèmes d’équations aux dérivées partielles non linéaires. *Acta Mathematica* 88 (1952) 141 - 225.
- [35] J. Frauendiener. Conformal infinity. *Living Rev. Relativity* 7 (2004) <http://www.livingreviews.org/lrr-2004-1>
- [36] J. Frauendiener, J. Hennig. Fully pseudospectral solution of the conformally invariant wave equations near the cylinder at space-like infinity. *Class. Quantum Grav.*, 31 (2014) 085010.
- [37] H. Friedrich. On the regular and the asymptotic characteristic initial value problem for Einstein’s vacuum field equations. Proceedings of the 3rd Gregynog Relativity Workshop on Gravitational Radiation Theory MPI-PEA/Astro 204 (1979) 137-160 and *Proc. Roy. Soc.*, 375 (1981) 169-184.
- [38] H. Friedrich. The asymptotic characteristic initial value problem for Einstein’s vacuum field equations as an initial value problem for a first-order quasilinear symmetric hyperbolic system. *Proc. Roy. Soc. Lond. A* 378 (1981) 401-421.
- [39] H. Friedrich. Cauchy problems for the conformal vacuum field equations in General Relativity. *Commun. Math. Phys.* 91 (1983) 445-472.
- [40] H. Friedrich. On the existence of n-geodesically complete or future complete solutions of Einstein’s field equations with smooth asymptotic structure. *Commun. Math. Phys.* 107 (1986), 587 - 609.
- [41] H. Friedrich. On the global existence and the asymptotic behaviour of solutions to the Einstein-Maxwell-Yang-Mills equations. *J. Differential Geometry* 34 (1991) 275 - 345.
- [42] H. Friedrich. Gravitational fields near space-like and null infinity. *J. Geom. Phys.* 24 (1998) 83–163.
- [43] H. Friedrich. Einstein’s equation and geometric asymptotics. In: N. Dadhich, J. Narlikar (eds) *Gravitation and Relativity: At the turn of the Millennium*. Inter-University Centre for Astronomy and Astrophysics, Pune, India, 1998. arXiv:gr-qc/9804009

- [44] H. Friedrich. Conformal Einstein evolution. In: J. Frauendiener, H. Friedrich (eds.) *The Conformal Structure of Spacetime: Geometry, Analysis, Numerics*. Springer, Berlin, 2002.
- [45] H. Friedrich. Spin-2 fields on Minkowski space near spacelike and null infinity. *Class. Quantum Grav.* 20 (2003) 101 - 117.
- [46] H. Friedrich. Smoothness at null infinity and the structure of initial data. In: P. T. Chruściel, H. Friedrich (eds.): *The Einstein Equations and the Large Scale Behaviour of Gravitational Fields*. Birkhäuser, Basel, 2004.
- [47] H. Friedrich. Initial boundary value problems for Einstein's field equations and geometric uniqueness. *Gen. Relativ. Gravit.* 41 (2009) 1947 - 1966.
- [48] H. Friedrich. The Taylor expansion at past time-like infinity. *Commun. Math. Phys.* 324 (2013) 263 - 300.
- [49] H. Friedrich. Geometric asymptotics and beyond. In: L. Bieri, S.-T. Yau (eds), *Surveys in Differential Geometry*, Vol.20. International Press, Boston, 2015. arXiv:1411.3854
- [50] H. Friedrich, J. Kánnár. Bondi systems near space-like infinity and the calculation of the NP-constants. *J. Math. Phys.* 41, (2000), 2195 - 2232.
- [51] R. Geroch. Asymptotic structure of space-time. In: F. P. Esposito, L. Witten: *Asymptotic structure of space-time*. Plenum, New York, 1977
- [52] R. Geroch, G. T. Horowitz. Asymptotically simple does not imply asymptotically Minkowskian. *Phys. Rev. Lett.* 40 (1978) 203 - 206.
- [53] S. W. Hawking, G. F. R. Ellis. *The large scale structure of space-time*. Cambridge University Press, Cambridge, 1973.
- [54] P. Hübner. From now to timelike infinity on a finite grid. *Class. Quantum Grav.* 18 (2001) 1871 - 1884.
- [55] T. Hughes, T. Kato, J. Marsden. Well-posed quasi-linear second-order hyperbolic systems with applications to nonlinear elastodynamics and general relativity. *Arch. Ration. Mech. Anal.* 63 (1977) 273 -294.
- [56] J. Kánnár. Hyperboloidal initial data for the vacuum Einstein equations with cosmological constant. *Class. Quantum Grav.* 13 (1996) 3075 - 3084.
- [57] H. Lindblad. On the asymptotic behaviour of solutions to the Einstein vacuum equations in wave coordinates. *Commun. Math. Phys.* 353 (2017) 135 - 184.
- [58] H. Lindblad, I. Rodnianski. The global stability of Minkowski space-time in harmonic gauge. *Ann. Math.* 171 (2010) 1401 - 1477.
- [59] E. T. Newman, R. Penrose. An approach to gravitational radiation by a methods of spin coefficients. *J. Math. Phys.* 3 (1962) 566 - 578.
- [60] R. Penrose. Asymptotic properties of fields and space-time. *Phys. Rev. Lett.* 10 (1963) 66 - 68.

- [61] R. Penrose. Zero rest-mass fields including gravitation: asymptotic behaviour. *Proc. Roy. Soc. Lond. A* 284 (1965) 159 - 203.
- [62] R. Penrose. Relativistic Symmetry Groups. In: A. O. Barut (ed.): *Group theory in non-linear problems*, Reidel, Dordrecht, 1974
- [63] O. Rinne, V. Moncrief. Hyperboloidal Einstein-matter evolution and tails for scalar and Yang-Mills fields. *Class. Quantum Grav.* 30 (2013) 095009
- [64] R. K. Sachs. Gravitational waves in general relativity VI. The outgoing radiation condition. *Proc. Roy. Soc A* 264 (1961) 309–338.
- [65] R. K. Sachs. Gravitational waves in general relativity VIII. Waves in asymptotically flat space-time. *Proc. Roy. Soc A* 270 (1962) 103–126.
- [66] J. A. Valiente Kroon. Asymptotic properties of the development of conformally flat data near spatial infinity. *Class. Quantum Grav.* 24 (2007) 3037 - 3053