

Annual Review of Animal Biosciences

Bat Biology, Genomes, and the Bat1K Project: To Generate Chromosome-Level Genomes for All Living Bat Species

Emma C. Teeling,¹ Sonja C. Vernes,^{2,3}
Liliana M. Dávalos,⁴ David A. Ray,⁵
M. Thomas P. Gilbert,^{6,7} Eugene Myers,⁸
and Bat1K Consortium*

¹School of Biology and Environmental Science, University College Dublin, Belfield, Dublin 4, Ireland; email: emma.teeling@ucd.ie

²Neurogenetics of Vocal Communication Group, Max Planck Institute for Psycholinguistics, Nijmegen, 6500 AH, The Netherlands

³Donders Centre for Cognitive Neuroimaging, Nijmegen, 6525 EN, The Netherlands

⁴Department of Ecology and Evolution, Stony Brook University, Stony Brook, New York 11794-5245, USA

⁵Department of Biological Sciences, Texas Tech University, Lubbock, Texas 79409, USA

⁶Natural History Museum of Denmark, University of Copenhagen, 1350 Copenhagen, Denmark

⁷University Museum, Norwegian University of Science and Technology, 7491 Trondheim, Norway

⁸Max Planck Institute for Molecular Cell Biology and Genetics, 01307 Dresden, Germany

*Full list of Bat1K Consortium members in Supplemental Appendix



ANNUAL REVIEWS Further

Click here to view this article's online features:

- Download figures as PPT slides
- Navigate linked references
- Download citations
- Explore related articles
- Search keywords

Annu. Rev. Anim. Biosci. 2018. 6:23–46

First published as a Review in Advance on November 20, 2017

The *Annual Review of Animal Biosciences* is online at animal.annualreviews.org

<https://doi.org/10.1146/annurev-animal-022516-022811>

Copyright © 2018 by Annual Reviews.
All rights reserved

Keywords

echolocation, flight, longevity, immunity, ecosystem, mammals

Abstract

Bats are unique among mammals, possessing some of the rarest mammalian adaptations, including true self-powered flight, laryngeal echolocation, exceptional longevity, unique immunity, contracted genomes, and vocal learning. They provide key ecosystem services, pollinating tropical plants, dispersing seeds, and controlling insect pest populations, thus driving healthy ecosystems. They account for more than 20% of all living mammalian diversity, and their crown-group evolutionary history dates back to the Eocene.

Despite their great numbers and diversity, many species are threatened and endangered. Here we announce Bat1K, an initiative to sequence the genomes of all living bat species ($n \sim 1,300$) to chromosome-level assembly. The Bat1K genome consortium unites bat biologists (>148 members as of writing), computational scientists, conservation organizations, genome technologists, and any interested individuals committed to a better understanding of the genetic and evolutionary mechanisms that underlie the unique adaptations of bats. Our aim is to catalog the unique genetic diversity present in all living bats to better understand the molecular basis of their unique adaptations; uncover their evolutionary history; link genotype with phenotype; and ultimately better understand, promote, and conserve bats. Here we review the unique adaptations of bats and highlight how chromosome-level genome assemblies can uncover the molecular basis of these traits. We present a novel sequencing and assembly strategy and review the striking societal and scientific benefits that will result from the Bat1K initiative.

INTRODUCTION

Of all living mammals, bats are extraordinary given their unique and peculiar adaptations. From the largest golden-capped fruitbat (*Acerodon jubatus*), with a wingspan of ~ 1.5 m and a weight of ~ 1 kg, to the smallest, ~ 2 -g echolocating bumblebee bat (*Craseonycteris thonglongyai*), the huge diversity and extraordinary adaptive radiations in bat form and function have both fascinated and terrified people for centuries (1–3). Bats account for $\sim 20\%$ of all living mammals (>1,300 species) and are found throughout the globe, absent only from the extreme polar regions. They are the only mammals that can truly fly and likewise have uniquely evolved laryngeal echolocation, or biosonar, which enables them to orient in complete darkness using sound alone (4–5). Using these and other traits, they have evolved to thrive in diverse ecological niches and can feed on insects, small mammals, fish, blood, nectar, fruit, and pollen (5). They perform key ecosystem services, pollinating crop species in the tropics (e.g., bats pollinate the flowers of agave, making possible the distillation of tequila), dispersing seeds, and feeding on crop pests throughout their range (6–9). Without bats, it is estimated that the United States would spend more than \$3 billion a year on pesticides alone (10). Bats are suspected reservoirs for some of the deadliest viral diseases [e.g., Ebola, SARS (severe acute respiratory syndrome), rabies, and MERS (Middle East respiratory syndrome coronavirus); 11–14], but they appear to be asymptomatic and survive these infections. This suggests that bats have evolved unique immune systems, and potentially the solution to better tolerate these pathogens may lie in uncovering how bats limit their immunopathology upon infection (15, 16). Bats also exhibit extraordinary longevity—they can live up to 10 times longer than expected given their small body size and high metabolic rate (17). Only 19 mammal species live proportionately longer than humans given their body size, and 18 are bats (17), with Brandt’s bat (*Myotis brandtii*) holding the reported record for bat longevity [>41 years, ~ 7 g (18)]. Bats show few signs of senescence and low to negligible rates of cancer (11), suggesting they have also evolved unique mechanisms to extend their health spans, rendering them excellent models to study extended mammalian longevity and ageing (17). Bats face a variety of global threats that threaten populations with regional or global extinction. The IUCN (International Union for Conservation of Nature) Red List currently classifies 77 bat species as Critically Endangered or Endangered and a further 184 as Vulnerable or Near Threatened due to significant population declines from conservation threats. Lack of knowledge about bat species hampers our ability to assess population stability in many cases; 222 bat species are considered Data Deficient by the

IUCN, meaning that their status cannot yet be determined, and 105 newly identified species are not yet listed (<http://www.batcon.org>).

The evolutionary history of bats has stimulated some of the most passionate debates in science, some spanning decades, from the initial disbelief among the scientific community when Spallanzani discovered in 1794 that bats were using sound to orient in darkness (19–21); to heated debates regarding whether the order was monophyletic and hence questioning if flight had a single origin in mammals (e.g., the flying primate hypothesis; 22); to current debates over the convergent evolution of laryngeal echolocation in bats and potential loss in the Old World fruit bats (for review, see 23, 24). Molecular phylogenetic analyses have reclassified their position within the mammalian tree (25–27), refuting their position within Archonta (including Primates, Scandentia, and Dermoptera) and placing them within Laurasiatheria (including Carnivora, Perissodactyla, Eulipotyphla, Pholidota, and Cetartiodactyla); and reclassified familial and interordinal relationships (**Figure 1**) (5, 23). Despite these advances, however, the sister taxon to bats remains unresolved (for review, see 23). As a mammalian order, bats have an impoverished fossil record, with an estimated >70% of fossil data missing (5, 28). There are astounding Lagerstätten Eocene fossils, but there is limited fossil representation thereafter until the Pliocene (29), making it difficult to reconstruct bat evolutionary history from fossils alone. Solving these outstanding evolutionary questions is difficult, as convergent homoplastic characters, incomplete lineage sorting, and a fragmented fossil record have limited phylogenetic reconstruction and obscured current understanding of the evolution and basis of the unique adaptations in bats.

Animal genomes are increasingly revealing the genetic basis of environmental niche specialization and adaptation (30–33), and studying the molecular mechanisms responsible for this diversity has allowed some of the greatest insights into the functioning and evolution of our own genome (32, 34, 35). Some of the most important challenges facing humanity into the next century are biological. These include improving the well-being of our large and rapidly ageing human populations (36), preventing the spread of emergent infectious diseases (37), maintaining agricultural productivity (10), and restoring natural ecosystems worldwide (38). These challenges will require a range of approaches to overcome them, starting with understanding the intrinsic mechanisms that make us vulnerable to disease and the ecological relationships underlying ecosystem maintenance and resilience. To date, insights into these various health-related challenges have primarily come from model organisms such as the fruit fly (*Drosophila melanogaster*) or the house mouse (*Mus musculus*). Having been optimized to be reared and studied in labs, these organisms reproduce in great numbers and live short lives, and thus are ideal experimental subjects. However, insights gained from the behavior and responses of short-lived organisms do not always translate to those that live longer, such as humans (17, 39, 40). Studying bats will enable us to address all of these challenges, as many of their biological features mirror humans and their ecological roles both contribute to and prevent the spread of infectious diseases, and structure functional ecosystems today and into the future.

We announce Bat1K, a global effort to sequence and annotate chromosome-level genome assemblies of all living bat species. Prioritization of bat genomes is not just desirable but indispensable to confront the many challenges to human well-being, ecosystem function, and biodiversity conservation we now face.

Central to the success of Bat1K is wide involvement from bat researchers across diverse research areas. This article aims to provide information to the scientific community about the Bat1K effort; to encourage participation; to set standards for tissue collection and vouchering, assembly quality, and data release; and to outline the major research endeavors that we anticipate will benefit from Bat1K.




Figure 1

Molecular consensus on bat familial relationships and divergence times (23). Asterisks indicate bat families with genomes available in the National Center for Biotechnology Information. Those species are listed in **Table 2**. Bat artwork created by Fiona Reid.

KEY AREAS OF IMPACT

The study of bats and their genomes is likely to have widespread impacts on a range of diverse scientific fields. Below we outline some key areas of study.

Model for Healthy Ageing

Unlike most lab animals, bats are excellent models for understanding human senescence and ageing and to discover the means to improve health into old age. Although there are some merits to mechanistic hypotheses of ageing (e.g., that life span should be inversely related to metabolic rates, as the latter contribute to the accumulation of intracellular debris leading to ageing), the best-supported theory of ageing is evolutionary and linked to life history (41). According to this theory, if adults in a population experience high extrinsic mortality (e.g., from predation), natural selection will favor short-term reproductive success over long-term survival and maintenance, resulting in rapid senescence after reproduction, past the age at which most individuals would have died in a natural population (42). The survivorship and life history of house mice reflect this pattern, with intrinsic mortality and morbidity rising sharply past one year of age, even under highly favorable lab conditions. Conversely, populations with low extrinsic mortality will experience continued selection throughout longer lifetimes, favoring slow senescence and resulting in longer, healthier lifetimes. Because bats are both nocturnal and capable of active flight, they have escaped the attention of most predators. This in turn has led them to evolve the relatively unusual vertebrate combination of long life spans with small bodies (17, 40, 43). As long-lived mammals, in some cases living >41 years, bats offer clues regarding the mechanisms for maintaining high function across internal systems over a long life span, longer than any similar-sized mouse can live (17, 44, 45).

Natural selection over millions of generations for continued health and reproduction throughout a long lifetime has equipped bats with excellent cellular and system-wide mechanisms of maintenance (45). This is particularly impressive considering that the high metabolic rates characteristic of bats are expected to produce reactive oxygen species, typically causing chronic inflammation and hastening senescence (46). However, the maintenance of function alone is not enough, as cells and tissues need constant repair over the course of a multiyear life span. Studies focused on bats have identified suites of cellular repair mechanisms that potentially evolved to support the unusual longevity of bats (11, 40, 44, 45). These genes and variants can be readily compared with human genes to discover specific features that would enable a healthy old age (11, 44, 45).

Model for Enhanced Disease Resistance

Bats have enhanced immune function, coupled with a potentially modulated inflammatory response (11, 16, 47). This holds considerable potential for addressing some of the worst consequences of senescence of humans. Inflammatory disorders associated with autoimmune diseases are among the fastest growing causes of disease worldwide, particularly in ageing populations (48). The ability to modulate inappropriate inflammation in response to stressors without impairing immune function could improve the lives of millions. Hence, detailed exploration of the genomic mechanisms of gene expression in wild bats could hold the key to improving health conditions worldwide (16). High-quality bat genomes will drive a better understanding of molecular bases underlying the resistance/tolerance of European bats to white-nose syndrome (49), which could ultimately be used to inform future bat conservation and management efforts within the United States. White-nose syndrome is a deadly fungal disease recently introduced to North America from Europe (50) that has decimated US bat populations, particularly of little brown bats (*Myotis lucifugus*) and tricolored bats (*Perimyotis subflavus*), and is responsible for an estimated 5–6 million

Table 1 Selected emerging viral pathogens, bat species in which they have been found, geographic locales of infected bats, and bibliographic sources

Emerging viral pathogen	Bat species	Geographic locales	Source
Hendra virus	<i>Pteropus poliocephalus</i>	Queensland, Australia	170
Nipah virus	<i>Pteropus hypomelanus</i> , <i>Pteropus lylei</i> , <i>Pteropus vampyrus</i>	Cambodia, Malaysia	171
Australian bat lyssavirus	<i>Pteropus alecto</i> , <i>Pteropus poliocephalus</i> , <i>Pteropus conspicillatus</i> , <i>Pteropus</i> <i>scapulatus</i> , <i>Saccolaimus flaviventris</i>	Queensland, Australia	172
European bat lyssavirus 1	<i>Eptesicus serotinus</i> , <i>Eptesicus isabellinus</i> , <i>Plecotus auritus</i> , <i>Pipistrellus pipistrellus</i> , <i>Pipistrellus nathusii</i>	Germany, the Netherlands, Denmark, Poland, France, Spain	173
Marburg virus	<i>Rousettus aegyptiacus</i>	Gabon	174
SARS-like coronaviruses	<i>Rhinolophus sinicus</i>	Kunming, Yunnan Province, China	175
Relative of Middle East respiratory syndrome coronavirus	<i>Neoromicia</i> cf. <i>zuluensis</i>	South Africa	176

bat deaths (51). Closely related European bats, such as greater mouse-eared bats (*Myotis myotis*) and Brandt's bat (*Myotis brandtii*), can be colonized by the responsible fungus *Pseudogymnoascus destructans* but do not suffer the same mass mortality (52), suggesting that European bats have evolved the required immune/behavioral response to survive this infection.

As wide-ranging individuals that constantly encounter new environments, bats come into contact with diverse viral pathogens over the course of a life span (15, 53). Again, over eons, bats have evolved to cope with these challenges, often leading to survival despite viral exposure (for a comprehensive list of viruses detected in bats, see 54). Epidemiological studies and field surveys suggest viruses circulate in wild bat populations (12, 55) without causing the great morbidity or mortality observed as a result of viral spillovers into humans (Table 1) (14, 56). Although the processes through which bats manage to clear viral infections remain poorly understood, possible explanations for the resilience of individual bats to these infections include not only heightened immune function but modulation of inflammation and mechanisms of repair (16). Genomic analyses have already revealed coevolution with viruses as a long-term consequence of bat evolution (57). High-quality bat genomes (e.g., including regulatory regions) from diverse species are needed to determine both how extensive these interactions between bats and viruses have been and what innate genetic mechanisms bats use to clear these viruses. Coupled with viral monitoring in the wild and viral genomics, bat genomic analyses have the potential to predict and manage future spillover events (56).

Key Species for Ecosystem Functioning

The unique biology of bats is important not only for understanding human health but also for maintaining and improving ecosystem health (7, 58). Bats contribute disproportionately to ecosystem functions, for example, in the regeneration of tropical forests (59–61). The convergent evolution of mutualism with plants in both Old World and New World tropics has led to the dependence of many flowering plant species on bats for their reproduction and dispersal. Pollinating and seed-dispersing bats are keystone species across the largest stands of tropical forests in biodiversity hotspots from Amazonia to the Congo Basin to Southeast Asia (7, 60). As natural forests are increasingly fragmented for human uses, bats play a unique role in dispersing pollen among plants

and seeds across large distances, maintaining plant genetic diversity and aiding the regeneration of forests after clearing. Breaching ocean barriers through active flight, bats are often the only mammals native to oceanic islands and are important to the survival and maintenance of those isolated ecosystems. As people have brought in predators and commensal species, island bats have become increasingly imperiled, threatening to collapse local trophic chains and disrupt pollination and seed dispersal (61). Bats also feed on many pest insect species and act as a natural and effective pest control (6, 8–10, 58). Insect-eating bats save maize farmers globally an estimated ~US\$1 billion a year from crop damage (58). Bat genomes will help elucidate the genetic adaptations enabling bats to both detect flowering plants and act as pollinators and will provide the basic genomic resources and tools for future population-level analyses to investigate and understand the relationships between their dwindling populations and the ecosystems on which they rely and that rely on them.

Model for the Evolution of Sensory Perception

Sensory perception plays one of the most important roles in the survival of an individual and is responsible for many key behaviors (e.g., foraging, predator avoidance, mate recognition, and communication) that drive evolution. Therefore, genes involved in sensory perception should show signs of adaptation and selection, as these genes are at the frontline of evolutionary pressures. Indeed, spectacular evidence of molecular convergence has been uncovered in hearing genes in echolocating whales and certain echolocating bats (62–65). Olfactory receptor genes that are directly involved in olfaction show evidence of environmental niche specialization in aquatic, terrestrial, and flying mammals, even after controlling for phylogeny (31, 66, 67). Similar loss of function in short-wave opsin visual genes has been found in bats with advanced echolocation capabilities (30). Bats have undergone the most extensive loss of function of the pheromone-detecting vomeronasal system compared with any other mammalian group (68–70), offering an opportunity to better understand the process of sensory vestigialization and the vomeronasal loss seen throughout mammals, including humans. Therefore, studying the evolution of these genes and genomic regions in bats—the sensory specialists—will elucidate how the mammalian genome has responded to past evolutionary pressures driven by changing environmental conditions. Comparative genomic analyses will also help illuminate the evolution and inheritance of blindness and deafness in humans by identifying regions of these genes that are most likely to cause disease and thus identifying putative targets for downstream gene therapeutics (71–73). This is an important advance, as the World Health Organization has estimated that 285 million people worldwide are visually impaired, 39 million of them are blind (74, 75), and over 360 million people have severe hearing loss. Bats have also been shown to be an excellent model for sensory-driven speciation [e.g., *Rhinolophus philippinensis* (76), *Craseonycteris thonglongyai* (77)]. High-quality bat genomes will enable further elucidation of the molecular basis of sensory adaptation and finally untangle the evolutionary mechanisms driving speciation (77, 78).

Model of Human Communication

Humans are unique in their capacity for language, but a core component of spoken language is the ability to learn new vocalizations, and this is shared with only a few other species, including some bats, birds, cetaceans, pinnipeds, and elephants (79–82). Comparative analyses of humans with other vocal learning species will be key for revealing its biological and evolutionary underpinnings. To date, vocal learning has been studied extensively in birds. For example, comparative analyses facilitated by the release of avian genomes have revealed genes with unique signatures in vocal

learning birds (83, 84). Genome-wide expression analyses have also found that many of these genes are differentially expressed in vocal learning brain regions, pointing to a link between these genes and the distinctive vocal abilities of these birds (85). Mammalian vocal learning is, by comparison, understudied, particularly at a neurogenetic level. This is largely because of the scarcity of the trait and the large size and intractability of most vocal learning mammals (e.g., elephants, whales, dolphins). Their diversity, strong reliance on vocal social communication, and small size make bats an attractive and experimentally tractable model for studying vocal learning (85a). Avian research has highlighted shared neurogenetic mechanisms that may underlie vocal learning across divergent species. Adding bats to this field will bridge the wide evolutionary gap between birds and humans and will bring us closer to understanding how this complex behavioral trait can be genetically encoded in the brain. The availability of high-quality bat genomes will provide a mammalian system in which the biological basis of vocal learning can be explored and shed light on how this language-relevant trait has evolved across Aves, Mammalia, and humans. In addition, understanding the genetic bases of language-relevant traits like vocal learning in bats will help us to understand the genetic causes of disorders involving speech and/or language (e.g., language disorder, autism, and speech apraxia) in humans. These disorders are highly prevalent (up to 25% of children; 86) and have a strong genetic component, and understanding their causes will lead to better genetic diagnoses and new potential therapeutics.

Model for Limb Development

The bat wing is a striking example of morphological adaptation and variation in mammals, characterized by dramatically elongated fingers and retained interdigital webbing. Several genomic studies have recently been carried out on developing bat embryonic limbs that identified numerous candidate wing development-associated genes and regulatory elements (87–89). Importantly, complete genomes are critical to this task, which requires tracking changes in expression encoded by regulatory regions, and not just protein-coding changes. The Bat1K project would enable comparative genomic analyses that can highlight specific genes, regulatory elements, and bat-specific nucleotide changes that are associated with wing development. Characterizing bat wing development will also improve our understanding of how changes in limb developmental building blocks can lead to human limb malformations, such as arachnodactyly (long fingers), brachydactyly (short fingers), and syndactyly (webbed fingers).

Evolution of Mammalian Genome Architecture

Bats exhibit the smallest genome size of all extant mammals, ranging in size from ~1.6 to 3.54 Gb based on 266 records, with the majority of genomes being ~2 Gb (90; data from <http://www.genomesize.com>). Birds, the other class of flying vertebrates, show similarly reduced genome sizes, an observation that has been used to suggest that the acquisition of flight requires a loss of genomic redundancy and a streamlining of genomic structure within vertebrates (90–92). Bats represent the lower limit of mammalian genome content and therefore can show the essential genomic information required for being mammalian. In addition, vespertilionid bats are unique among studied mammals in harboring multiple lineages of active transposable elements that are not found in other members of the clade (93). Despite this unique accumulation, they continue to maintain small genomes, suggesting an ability to balance genomic gain from transposable repeats via the loss of genomic mass (90). Chromosome-level, high-coverage bat genomes across the order will drive a unique understanding of the limits of mammalian genome structure in general and the genomic consequences of flight adaptation in particular.

Although many other species have some of these attributes to varying degrees, bats are unique because their genomes harbor secrets related to all of them. Thus, obtaining genome assembly data across the bat clade will drive the advancement of these and numerous other fields given the unique biology and evolutionary history of bats.

BAT GENOMES SEQUENCED TO DATE

The first complex eukaryotic genomes were generated using Sanger (dideoxynucleotide) chemistry, which, given the output limitations, rendered them both complex and expensive endeavors. For example, the first human genome draft took well over a decade to produce and cost over US\$1 billion (94). Sanger sequencing was the state of the art for more than two decades, with few eukaryotic genomes being released owing to the cost and time commitments. However, since the introduction of new, high-throughput chemistries in the mid-2000s (95), those costs and time commitments have been substantially reduced.

Currently, two major technologies dominate the field. Illumina sequencing is the highest-throughput option available at present, with some equipment able to generate the equivalent of a human genome in just a few hours, at a cost of only a few hundred US dollars. Briefly, Illumina sequencing relies on the sequential addition of fluorescently labeled nucleotides and their subsequent detection. The technology is very accurate, with error rates of only $\geq 0.1\%$ with variation depending on genomic region, read lengths, and sequencing chemistry (96, 97). The trade-off for accuracy and throughput, however, is read length. Illumina sequencing chemistry has yet to break the 300-bp limit, with most users limiting themselves to reads of 150 bp or less, which can make genome assembly across complex repetitive regions difficult if not impossible (33).

Pacific Biosciences (PacBio) has adopted a different sequencing strategy based on single-molecule real-time (SMRT) sequencing. Briefly, this method relies on a tethered polymerase that incorporates labeled nucleotides using the template to be read. The fluorescent labels are cleaved on addition of the nucleotide, and a detector identifies the base added. Current throughput averages approximately 8 Gb/day (98, 99), with reads averaging 10–15 kb and some ranging up to 100,000 bp. Unfortunately, PacBio reads exhibit a substantially higher error rate of 10–15%, mostly in the form of insertions and deletions. Methods exist to improve this to 99.99% accuracy, but this requires at least $50 \times$ SMRT-based coverage or the addition of Illumina's shorter reads (100–106), making this option more expensive than Illumina-only approaches.

As of writing, 14 bat genome assemblies are readily available from the National Center for Biotechnology Information (NCBI) database (**Table 2**). The first bat to have its genome released, the little brown bat [*Myotis lucifugus* (107)], was sequenced by the Broad Institute using Sanger chemistry. This member of the clade Yangochiroptera (**Figure 1**) (5) has been the subject of most comparative analyses among mammals and has served as the chiropteran representative.

The remaining assemblies vary in quality and completeness. For example, Parker et al. (65) took a low-coverage approach, which was valid given that they were interested primarily in coding sequences. Unfortunately, such low coverage makes overall contiguity of the assembly very low, as evidenced by the low-contig N50s. In comparison, the Egyptian fruit bat (*Rousettus aegyptiacus*) assembly, made possible by the combination of long- and short-read technologies, is of higher quality (**Table 2**). The search for genomic signatures for complex traits underlying the unique biology of bats, however, requires a common standard for greater comparability.

Of course, other efforts are under way or completed that are not represented in the NCBI database. Members of Bat1K are, for example, currently involved in efforts to sequence and assemble the genomes of the Northern long-eared bat (*Myotis septentrionalis*), the greater mouse-eared

Table 2 Summary statistics of bat genomes currently available via National Center for Biotechnology Information databases^a

Suborder	Species/family	Common name	Estimated genome size (C) ^b	Assembly size (Gb)	GenBank accession	Publication	Sequencing technology	Coverage	Contigs	Contig N50	Scaffolds	Scaffold N50	Assembly score
Ynpiterochiroptera	<i>Eidolon helvum</i> / Pteropodidae	African straw-colored fruit bat	2.03	1.44	AH60+-1	65	IL	18x	288,446	12,668	6,789	27,684	0.1.1Q35
	<i>Hipposideros armiger</i> / Hipposideridae	Great leaf-nosed bat	ND	2.24	JXIKO+-1	168	IL	218x	128,956	39,863	7,571	2,328,177	0.3.1Q40
	<i>Megaderma lyra</i> / Megadermatidae	Greater false vampire bat	ND	1.44	AH60+-1	65	IL	18x	518,327	7,043	192,872	16,881	0.1.1Q35
	<i>Pteropus alerno</i> / Pteropodidae	Black flying fox	ND	1.99	ALMSO+-1	47	IL	110x	170,164	18,318	65,598	15,954,802	1.4.1Q40
	<i>Pteropus vampyrus</i> / Pteropodidae	Large flying fox	2.37	2.14	ABRP0+-2		IL	188x	225,433	21,866	36,094	5,954,017	2.3.1Q40
	<i>Rhinolophus ferrumequinum</i> / Rhinolophidae	Greater horseshoe bat	1.92–2.68	1.31	AH60+-1	65	IL	17x	290,685	11,695	160,500	21,151	1.1.1Q35
	<i>Rhinolophus sinicus</i> / Rhinolophidae	Chinese rufous horseshoe bat	ND	2.07	LVEHO+-1	168	IL	146x	182,995	37,803	63,439	3,754,400	1.3.1Q40
	<i>Rousettus aegyptiacus</i> / Pteropodidae	Egyptian rousette	2.11	1.91	LOCFO+-2		IL & PB	169x	3,049	1,488,988	2,490	2,007,187	3.3.1Q40
	<i>Eptesicus fuscus</i> / Vespertilionidae	Big brown bat	2.25–2.70	2.02	AL6HO+-1		IL	84x	137,058	21,392	133,538	13,454,942	1.4.1Q40
	<i>Minioterus natalensis</i> / Miniotertidae	Natal long-fingered bat	ND	1.80	LDJU0+-1	89	IL	77x	114,632	29,777	1,269	4,315,193	1.3.1Q40
	<i>Myotis brandtii</i> / Vespertilionidae	Brandt's bat	ND	2.11	AMKRO+-1	169	IL	120x	325,414	23,289	169,750	3,225,832	1.3.1Q40
	<i>Myotis dasidii</i> / Vespertilionidae	David's myotis	ND	2.06	ALMTO+-1	47	IL	110x	325,280	15,182	101,769	3,454,484	1.3.1Q40
	<i>Myotis lucifugus</i> / Vespertilionidae	Little brown myotis	2.26	1.96	AAPEO+-2	107	SG	7x	72,785	64,330	11,654	4,293,315	1.3.1Q35
<i>Pteronotus parnellii</i> / Mormoopidae	Common mustached bat	2.35–2.71	1.56	AM6ZO+-1	65	IL	17x	443,121	9,502	177,401	22,675	0.1.1Q35	

^aPhylogenetic relationships of bat families are depicted in Figure 1.

^bData taken from <https://genomesize.com>, the animal genome size database.

Abbreviations: 0+, no data; ND, no data; PB, Pacbio; SG, Sanger.


bat (*Myotis myotis*), the common vampire bat (*Desmodus rotundus*), and the pale spear-nosed bat (*Phyllostomus discolor*). These assemblies use multiple strategies with results that vary in ways similar to those observed in **Table 2**. Indeed, one primary goal of Bat1K is to standardize assembly strategies to provide assemblies of uniform optimal quality for the bat genomics community through combining multiple sequencing and scaffolding technologies (e.g., PacBio+Illumina+HiC+10X; see below for details).

Given the more than 1,300 species of bat currently recognized, there is still a long way to go to generate genome sequence data covering Chiroptera (108, 109). However, as we outline in the next section, we believe it is important not just to generate genome-level data but to produce high-quality genome sequences that maximize the usefulness and accessibility of the data for all research fields. Below, we outline the specific goals of the Bat1K Consortium and how we aim to achieve these goals.

THE BAT1K PROPOSAL

The ultimate goal of Bat1K is to sequence, assemble to chromosome-level contiguity, and make publicly available the full genome of every living bat species. As one might imagine, given the specialized needs of bats in captivity, determining species boundaries is difficult in chiropterans. Indeed, applying a particular species concept to bats is difficult. The Biological Species concept (110) is commonly applied to mammals, but investigations of gene flow among bat species have been attempted but are not commonly performed and often rely on indirect evidence (76–78, 111–115). The Genetic Species concept was tested for multiple species and suggested hybridization in several cases (116). Some clades, in particular the vesper bats, likely harbor significant numbers of cryptic species (117). Thus, the actual number of species that we could conceivably target is unknown. However, our best estimate is ~1,300 (108, 109; <http://www.bat1k.com>). The Bat1K genome consortium unites bat biologists (>148 members as of writing; see full list of consortium members in the Supplemental Appendix), computational scientists, conservation organizations, genome technologists, and any interested individuals committed to a better understanding of the genetic and evolutionary mechanisms that underlie the unique adaptations of bats. Our aim is to catalog the unique genetic endowment and diversity present in all living bats to better understand the molecular basis of their unique adaptations; uncover their evolutionary history; link genotype with phenotype; and ultimately better understand, promote, and conserve bats. To achieve this, we must meet the following objectives.

Acquire high-molecular weight DNA from all living bat species. Highly contiguous assemblies require high-molecular weight DNA as starting material. We are coordinating members of Bat1K to obtain such material. Many members have already contributed or pledged samples, and others have committed to acquiring other samples. Our preference is that fresh tissue should be flash frozen in liquid nitrogen immediately after death (see <http://www.bat1k.com> website for methodology) and remain frozen (at <80°C) until DNA extraction and immediate library preparation on site at the sequencing location. We understand this may not always be possible. In cases where it is not possible to lethally sample a particular species given its conservation/protected status, nonlethal wing-punch biopsies should be taken, placed in viable media, and sent to appropriate laboratories for cell culture, as described by Kacprzyk et al. (118). In these cases, high-molecular weight DNA will be harvested after minimal passage of cells and immediately flash frozen or subjected to DNA extraction. All samples will be taken and transported with full capture, licensing, and legal permits required. Additionally, each individual selected for genomic sequencing must be associated with an electronic voucher or, even better, a specimen voucher deposited in a collection.

 Supplemental Material

The minimal requirements for an adequate specimen voucher are (a) species identification based on expert assessment; (b) standard mammalian measurements, including forearm and body mass; (c) geographical coordinates of capture; and (d) photographs documenting the individual, its genitalia, and its dentition. Because male mammals are heterogametic, we recommend sampling both males and females where possible; if only one is available, males should be chosen to allow for Y-chromosome sequencing. However, given that many bat species are typically accessible only at maternity roosts, we will sequence a female representative when no male samples are available. Geographic provenance and morphometric data will be recorded, and when possible, additional tissues harvested from the same animal will be frozen and biobanked by consortium members.

Produce reference quality-based genomes. Informally, a chromosome-level assembly refers to a genome reconstruction for which the sequence is in megabase contigs that are ordered and oriented into scaffolds that effectively cover each chromosome arm. Very few vertebrate genomes today approach such a level of contiguity and completeness. However, the new single-molecule, long-read technologies (119, 120) are changing this situation. Our goal is to produce reconstructions with a contig N50 of 1 Mbp or greater, a scaffold N50 of 10 Mbp or greater, at least 90% of the contigs mapped to chromosomes, and a consensus accuracy of Q40 or better, or what we call a 3.4.2Q40 assembly, as we will now rigorously define.

Formally, a *c.s.m.Qx* (contig/scaffold/map-chromosome/Qscore) genome reconstruction is one that has a contig N50 of 10^m Kbp, a scaffold N50 of 10^m Kbp, and a consensus accuracy of Qx; $m = 1, 2, 3$, or 4 is a descriptive designator of how well the scaffolds are mapped to chromosomes, as follows:

1. Not assigned to chromosomes, or under 90% assigned.
2. >90% assigned to chromosomes, either directly to species-specific chromosomal maps, or by in silico assignment to closest reference or ancestral reconstructed chromosomes.
3. >90% assigned to chromosomes as for 2, but also requiring that all interchromosomal breakpoints with respect to the reference are confirmed by at least two independent data sources.
4. >90% assigned to species-specific chromosomal maps via direct within-species map data, with all interchromosomal and a representative sample of intrachromosomal breakpoints with respect to the reference confirmed.

This previously unpublished characterization (Richard Durbin & Harris Lewin, private communication) is the standard that has been adopted by the broader Vertebrate Genome Project (<http://www.genomeark.org>) as part of the Genome 10K initiative (<https://genome10k.soe.ucsc.edu>). By way of reference, at the end of 2016, 3 mammals of the 290 sequenced vertebrates met the 3.4.2Q40 standard that we will target, and the human genome is currently 4.4.2.Q40. Estimated scores are given for the bat genomes in **Table 2**.

Our proposed strategy, to generate highly contiguous and complete assemblies, will be to collect sequence data structured as follows: (a) $60\times$ coverage in PacBio long reads, (b) $50\times$ coverage in Illumina short reads collected as 10X Genomics read clouds, and (c) $10\times$ coverage in Illumina short reads collected as long-range Hi-C read pairs. Initial assemblies will be accomplished by using the most accurate current assemblers, such as Falcon and Canu (121, 122), with long reads of an average length of 10–15 Kb. Assemblies of other mammals (101, 123) suggest that we will obtain initial contigs with an N50 length of over 1 Mb and often as high as 5 Mb. Genomes that are highly repetitive and large, such as those of amphibians (e.g., the axolotl at 32 Gb, with transposable elements comprising $\sim 60\%$), do not assemble to this level. However, we have no

reason to expect any bat genomes to prove problematic, as the bat genomes sequenced to date are mostly ~ 2 Gb (90; <http://www.genomesize.com>), and although some bats contain unique transposable element content by type, they are no more repetitive on average than those of other mammals (90).

Scaffolding of the contigs will then be performed using the 10X Genomics read clouds and Hi-C data. The 10X Genomics device and protocol generate short-read templates that lightly cover (i.e., $0.2\text{--}3\times$) a handful (i.e., $5\text{--}15$) of $50\text{--}200\text{-Kbp}$ molecules in a droplet microchamber (124). Conceptually, sequencing the templates produced in each droplet yields clouds of reads that gather over a handful of regions of the underlying genome. Directly assembling these data with, for example, Supernova (125) has to date delivered results with large scaffolds but unfortunately small contigs. We will instead use the clouds for short-range scaffolding to link the PacBio contigs that are less than $100\text{--}200$ Kb apart, thereby bridging many of the gaps between contigs in the long-read assembly. We will use the first-generation software available to do this (126; <https://sourceforge.net/projects/phusion2/files/scaff10x>) and use newer, more optimal methods as they develop. Early anecdotal experience shows that this results in scaffolds well over 10 Mbp. Some large gaps will remain unspanned, and therefore we will use the long-range Hi-C paired-end data to link the 10X scaffolds into large super-scaffolds and separate these into chromosomal units (127, 128) with the aid of syntenic mapping to a phylogenetically close relative.

Finally, to achieve a Q40 consensus accuracy, we will use the $50\times$ coverage in Illumina reads provided by data sets *b* and *c* to polish the consensus (see below). This last step is necessary given that the high error rate of the long reads ($10\text{--}12\%$) typically results in a Q30 consensus when only the long-read data are used.

In summary, for Bat1K, we propose to collect (*a*) $60\times$ Pacbio long reads, (*b*) $50\times$ Illumina reads in 10X read clouds, and (*c*) $10\times$ Illumina reads in long-range Hi-C read pairs (4-cutter protocol) and follow the computational sequence below:

1. Assemble the $60\times$ PacBio data (*a*) into contigs.
2. Use the $50\times$ read clouds (*b*) to link contigs into scaffolds.
3. Find missed spanning reads and close the spanned gaps.
4. Use the Hi-C pairs (*c*) to order and orient the scaffolds into superscaffolds covering chromosome arms.
5. Use the $60\times$ Illumina read data from *b* and *c* to polish the PacBio consensus contigs to Q40.

A virtue of the proposed strategy is that only three commonly used library preparations and two sequencing technologies are required. Of course, should the protocol need amendment or augmentation, then such will be undertaken. This is especially true given the rapid advancements in sequencing technology observed today. Indeed, we see Bat1K as an opportunity to benchmark new genomic technologies that will enable a fast and cost-effective 3.4.2.Q40 genome.

Project Phases

This large sequencing project will be accomplished in three phylogenetically delimited phases (Figure 2). Phase 1 will target representatives of each bat family ($N=21$). In Phase 2, we will expand to include representatives for every genus of bat ($N=220$). Phase 3 will expand further to target all remaining species ($N=1,288$). As funding and samples becomes available from other parties, those efforts will be incorporated and lead the sequencing priority. At each phase, the community will publish key synthesis papers describing progress and exploring the unique adaptations within bats that drive this initiative.




Figure 2
Sequencing phases of Bat1K.

EXPECTED BENEFITS

The resources of Bat1K are of clear value to any researchers interested in linking genes to phenotype and understanding how genomes produce complex organisms, a pursuit that has been identified as one of the Grand Challenges of the Twenty-First Century (129, 130). A set of open-access, high-quality bat genomes that are sequenced, assembled, and annotated using uniform laboratory and bioinformatic pipelines will confer immense benefits to researchers from across the life sciences, ultimately yielding an unparalleled research tool that would form the basis for an uncountable number of research projects. Naturally, many of these researchers will be those whose interests focus principally on bats, and who wish to access the genomes of single, or a few closely related, species. However, as demonstrated in several other recently published phylogenomic-scale projects [for example, those on birds (131, 132), mammals (107), insects (133), and land plants (134)], as both the number and pan-clade distribution of genomes increase, so do the research possibilities. We foresee multiple avenues to pursue but happily acknowledge our inability to anticipate all of the research efforts that will be enabled.

Phylogeny and Population Genomics

In light of the rapidly decreasing costs of resequencing genomes, phylogenetic and population analyses are now increasingly being undertaken at the genome level, as it is widely understood that analyses based upon whole-genome data provide the most accurate reconstructions of species' evolutionary histories. This is particularly true for bats, for which many species are likely to have undergone rapid radiation, diversification, and potentially hybridization and/or incomplete lineage sorting (135–138). In particular, genome-level data sets should at best be able to account for the phylogenetic noise inherent in analyses of single genes or at worst at least clearly indicate the degree of noise that they impart (139). Obviously, the higher the quality of the genomic data, the greater the expected improvement. Given that different parts of the genomes can show strikingly different histories (e.g., 132, 140), it is important that the correct homology is inferred, which requires well-assembled genomes to enable alignment across divergent evolutionary time frames.

The Bat1K genome resource should enable a full reconstruction of the evolutionary history of this phylogenetically controversial clade for the first time and resolve long-standing debates by producing high-quality genomes that should overcome most previously encountered problems.

At the population scale, not only do a suite of genome-scale interpretations of standard tools now exist (e.g., nucleotide diversity, heterozygosity), but this can be complemented by other tools specifically developed to exploit the genome in new ways. Such tools include reconstruction of population size using pairwise sequentially Markovian coalescent (141), multiple sequentially Markovian coalescent (142), and the identification and quantification of admixture between genetically distinct groups using D-statistics (143). Although many of these analyses can be performed through mapping to the genomes of related species, in general the closer the reference genome, the more refined the observations made (144). With such genomes and tools in place, a wide range of questions that are central to current bat population biology and ecology can subsequently be addressed. These include what determines evolutionarily significant units (ESUs), what the degree of gene flow is among them, and even the dynamics of the speciation process itself (77, 78, 112), a critical question in bats given their dispersal ability and the high taxonomic diversity of some clades [e.g., genus *Myotis*, with 100+ species (145)]. These are, in turn, critical to more applied efforts, ranging from optimizing conservation strategies, for example, by focusing on ESUs at most risk or most likely to respond to efforts, to defining bat clades that represent possible high risks of cross-species pathogen outbreaks, for example, those exhibiting the greatest degree of inter-ESU, population, or even species admixture, and thus possible cross-host pathogen transmissions.

Uncover Genotype-To-Phenotype

High-quality genomic reference data also lie at the heart of studies aiming to reveal links between genotype and phenotype. Although they have historically been the domain of model organisms, in the era of genomics we are now able to expand our understanding of genotype-to-phenotype links beyond the typical model species. Thus, genomics allows us to choose the most appropriate model for the question, rather than just relying on the usual (and often less-well phenotypically suited) mouse, *Drosophila*, or nematode models. Although it was once acceptable to provide candidate loci identified through genome-wide association studies scans for loci under selection as evidence of genetic mechanisms in nonmodel systems, transcriptional analyses, and even knock-out/knock-in experiments, have become feasible and are increasingly required by journals (and the research community) today. Only with such evidence can we develop an understanding of how genes (or other genomic elements) function at a molecular level and thus truly bridge the gap between gene and phenotype. Animal systems with complete genomic information present an ideal opportunity, and indeed are essential to accomplish this. One of the clearest ways to prove genotype-to-phenotype links is to create transgenic animals or cell lines in which the candidate gene has been knocked down (e.g., via shRNAs) or knocked out (e.g., via CRISPR/Cas9 genome editing). But these approaches require detailed knowledge of the target organism's genome, as shRNA and CRISPR/Cas9 planning is based on sequence complementarity, and targeting approaches must be designed such that they are complementary only to the gene in question and to no other sequence (146, 147). Such progress has been exemplified by studies in birds in which, for example, genome sequence availability made it possible to knock down a key human language-related gene (*FoxP2*) in zebra finches to determine its role in circuitry underlying vocal learning (148–153).

Genomes also facilitate an understanding of the molecular basis of the gene-phenotype relationship through analysis of gene regulation. Techniques to understand regulatory networks, including small RNAs, transcription factor cascades (e.g., ChIP-Seq, enhancer mapping), epigenetic marks, and chromatin structure (e.g., histone mapping, chromatin conformation capture,

topological domain mapping), rely on a comprehensive understanding of the identity and position of both coding and noncoding DNA sequences (154–156). The perturbation of regulatory networks represents a particularly powerful way to invoke evolutionary change without the need to evolve novel genes (45). Such perturbations have been suggested as influential in bat diversification (157). This represents a potentially rich area of study, especially considering the rapid and massive changes to gene regulation that must have occurred when, over the course of only a few million years, the phyllostomid (Phyllostomidae) bats evolved from an insectivorous ancestor into the clade of carnivores, sanguivores, nectarivores, frugivores, and palynivores we see today (29, 158–160).

Evolution of Genome Architecture

As we move beyond questions of function, we reach those relating to more general questions, such as how genomes evolve. With high-quality reference genomes, this can be studied at levels ranging from the nucleotide up to the chromosome. At the nucleotide level, one series of questions of growing interest relates to constrained sequence evolution, for example, whether specific nucleotides or amino acids are under convergent evolution in phylogenetically disparate taxa, as has been showcased with regard to the evolution of echolocation in bats and marine mammals (e.g., 65, 161). At the larger scale, there is considerable focus today on evolutionary trajectories of autosomal and sex chromosomes (162), and in particular whether major evolutionary transitions accompany significant chromosome-scale rearrangements. The phyllostomids again represent an exciting test case. This clade exhibits a wide range of chromosomal variation, and chromosome-level genome assemblies across the group will allow us, Bat1K members, and other researchers to investigate this phenomenon from nucleotide, syntenic, and phylogenomic perspectives.

A third area that will benefit significantly from high-quality reference genomes is our understanding of the dark matter that large fractions of our genomes are built of—specifically the abundance of seemingly noncoding elements. Estimates from the human-focused ENCODE (163) project predict that between 10% and 80% of these noncoding components are functional (164)—clearly an unsatisfactory answer. Thus, one of the key aims of the ENCODE and other related projects is the combined use of biochemical assays and associated computational analyses to query the underlying DNA sequence, including the application of methods such as Hi-C to refine tertiary structure interactions (127, 165). However, a major challenge is simply the sheer size of genomic data sets—how can one effectively parse the data? In this regard, organisms such as bats, with their naturally reduced genome sizes, represent a unique opportunity to study a naturally filtered subset of this noncoding data. In short, one might hypothesize that elements retained in the reduced genomes of bats and birds (and in particular those with significant cross-species sequence conservation) may play important functions, whereas those purged do not. Indeed, preliminary analyses of bird genomes for both transposable elements and avian-specific highly conserved element content have already been promising in this regard (131, 166), indicating that the secrets within bat genomes may be of considerable use.

Wider Community Benefits

There are potential benefits on an even broader scale. The data set will represent a critical component of wider community efforts to catalog life's diversity, such as the Genome10K (129) and Earth Biogenome (167) projects—the latter of which aims to coordinate the sequencing and eventual public release of a full catalog of life's genomes. As such, these wider efforts will almost certainly contribute to the ongoing developments of not only the technologies underlying DNA


sequencing itself but the associated bioinformatics for both assembling and annotating the genomes and analyzing the data within them. Although many of these efforts will be directly linked to the sequencing industry, we highlight that because the resulting Bat1K data will be accessible by any scientists connected to the internet, it will represent an unparalleled, free opportunity for researchers around the world, including those in tropical countries where bat diversity concentrates, to both be trained in genomics and undertake genomic research. Critically, therefore, the training that this would confer on countless scientists would both strengthen local research across the planet and be taken into other industries, thus radically accelerating developments and discoveries linked to genomics in general—a topic of growing importance in the modern biomedicine and agriscience industries. Bat1K will develop a genomic record of all living bat species, ultimately a genomic ark that can be used to benchmark the genomic health of different bat species to uncover populations in need of immediate conservation efforts. Considering the challenges to well-being facing humanity, from ageing populations to emerging infectious diseases, and from conserving fragmented habitats to maintaining the ecosystems that preserve productivity, undertaking Bat1K is not only opportune but indispensable.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

A full list of Bat1K consortium members is presented in the **Supplemental Appendix**. We thank Fiona Reid for her contributed artwork.

 Supplemental Material

LITERATURE CITED

1. Allen GM. 1939. *Bats*. Cambridge, MA: Harvard Univ. Press
2. Nagel T. 1974. What is it like to be a bat? *Philos. Rev.* 83(4):435–50
3. Fenton MB, Grinnell AD, Popper AN, Fay RR, eds. 2016. *Bat Bioacoustics*. New York: Springer
4. Teeling EC, Dool S, Springer MS. 2012. Phylogenies, fossils and functional genes: the evolution of echolocation in bats. See Reference 29, pp. 1–22
5. Teeling EC, Springer MS, Madsen O, Bates P, O'Brien SJ, Murphy WJ. 2005. A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science* 307(5709):580–84
6. McCracken GF, Westbrook JK, Brown VA, Eldridge M, Federico P, et al. 2012. Bats track and exploit changes in insect pest populations. *PLOS ONE* 7(8):e43839
7. Bumrungsri S, Lang D, Harrower C, Sripaoraya E, Kitpipit K, et al. 2013. The dawn bat, *Eonycteris spelaea* Dobson (Chiroptera: Pteropodidae) feeds mainly on pollen of economically important food plants in Thailand. *Acta Chiropterologica* 15(1):95–104
8. Riccucci M, Lanza B. 2014. Bats and insect pest control: a review. *Vespertilio* 17:161–69
9. Puig-Montserrat X, Torre I, López-Baucells A, Guerrieri E, Monti MM, et al. 2015. Pest control service provided by bats in Mediterranean rice paddies: linking agroecosystems structure to ecological functions. *Mamm. Biol.* 80(3):237–45
10. Boyles JG, Cryan PM, McCracken GF, Kunz TH. 2011. Economic importance of bats in agriculture. *Science* 332(6025):41–42
11. Wang LF, Walker PJ, Poon LL. 2011. Mass extinctions, biodiversity and mitochondrial function: Are bats “special” as reservoirs for emerging viruses? *Curr. Opin. Virol.* 1(6):649–57
12. Drexler JF, Corman VM, Müller MA, Maganga GD, Vallo P, et al. 2012. Bats host major mammalian paramyxoviruses. *Nat. Commun.* 3:796

13. Brook CE, Dobson AP. 2015. Bats as “special” reservoirs for emerging zoonotic pathogens. *Curr. Trends Microbiol.* 23(3):172–80
14. Anthony SJ, Johnson CK, Greig DJ, Kramer S, Che X, et al. 2017. Global patterns in coronavirus diversity. *Virus Evol.* 3(1). <https://doi.org/10.1093/ve/vex012>
15. O’Shea TJ, Cryan PM, Cunningham AA, Fooks AR, Hayman DT, et al. 2014. Bat flight and zoonotic viruses. *Emerg. Infect. Dis.* 20(5):741–45
16. Kacprzyk J, Hughes GM, Palsson-McDermott EM, Quinn SR, Puechmaille SJ, et al. 2017. A potent anti-inflammatory response in bat macrophages may be linked to extended longevity and viral tolerance. *Acta Chiropterologica* 19:219–28
17. Austad SN. 2010. Methusaleh’s zoo: how nature provides us with clues for extending human health span. *J. Comp. Pathol.* 142:S10–S21
18. Podlutzky AJ, Khritankov AM, Ovodov ND, Austad SN. 2005. A new field record for bat longevity. *J. Gerontol. A Biol. Sci. Med. Sci.* 60(11):1366–68
19. Galambos R. 1942. The avoidance of obstacles by flying bats: Spallanzani’s ideas (1794) and later theories. *Isis* 34(2):132–40
20. Griffin DR. 1958. *Listening in the Dark: The Acoustic Orientation of Bats and Men*. New Haven, CT: Yale Univ. Press
21. Dijkgraaf S. 1960. Spallanzani’s unpublished experiments on the sensory basis of object perception in bats. *Isis* 51(1):9–20
22. Pettigrew JD. 1986. Flying primates? Megabats have the advanced pathway from eye to midbrain. *Science* 231(4743):1304–6
23. Teeling EC, Jones G, Rossiter SJ. 2016. Phylogeny, genes, and hearing: implications for the evolution of echolocation in bats. In *Bat Bioacoustics* ed. MB Fenton, AD Grinnell, AN Popper, RR Fay, pp. 25–54. New York: Springer
24. Wang Z, Zhu T, Xue H, Fang N, Zhang J, et al. 2017. Prenatal development supports a single origin of laryngeal echolocation in bats. *Nat. Ecol. Evol.* 1:0021
25. Murphy WJ, Eizirik E, Johnson WE, Zhang YP, Ryder OA, et al. 2001. Molecular phylogenetics and the origins of placental mammals. *Nature* 409(6820):614–18
26. Meredith RW, Janečka JE, Gatesy J, Ryder OA, Fisher CA, et al. 2011. Impacts of the Cretaceous Terrestrial Revolution and KPg extinction on mammal diversification. *Science* 334(6055):521–24
27. Tsagkogeorga G, Parker J, Stupka E, Cotton JA, Rossiter SJ. 2013. Phylogenomic analyses elucidate the evolutionary relationships of bats. *Curr. Biol.* 23(22):2262–67
28. Eiting TP, Gunnell GF. 2009. Global completeness of the bat fossil record. *J. Mamm. Evol.* 16(3):151–73
29. Gunnell GF, Simmons NB, eds. 2012. *Evolutionary History of Bats: Fossils, Molecules and Morphology*. Cambridge, UK: Cambridge Univ. Press
30. Zhao H, Rossiter SJ, Teeling EC, Li C, Cotton JA, et al. 2009. The evolution of color vision in nocturnal mammals. *PNAS* 106(22):8980–85
31. Hayden S, Bekaert M, Crider TA, Mariani S, Murphy WJ, et al. 2010. Ecological adaptation determines functional mammalian olfactory subgenomes. *Genome Res.* 20(1):1–9
32. Shapiro MD, Kronenberg Z, Li C, Domyan ET, Pan H, et al. 2013. Genomic diversity and evolution of the head crest in the rock pigeon. *Science* 339(6123):1063–67
33. Jarvis ED. 2016. Perspectives from the avian phylogenomics project: questions that can be answered with sequencing all genomes of a vertebrate class. *Annu. Rev. Anim. Biosci.* 4:45–59
34. Burk-Herrick A, Scally M, Amrine-Madsen H, Stanhope MJ, Springer MS. 2006 Natural selection and mammalian BRCA1 sequences: elucidating functionally important sites relevant to breast cancer susceptibility in humans. *Mamm. Genome* 17(3):257–70
35. Haussler D, O’Brien SJ, Ryder OA, Barker FK, Clamp M, et al. 2009 Genome 10K: a proposal to obtain whole-genome sequence for 10,000 vertebrate species. *J. Hered.* 100(6):659–74
36. De Magalhães JP. 2014. The scientific quest for lasting youth: prospects for curing aging. *Rejuvenation Res.* 17(5):458–67
37. Morens DM, Fauci AS. 2013. Emerging infectious diseases: threats to human health and global stability. *PLOS Pathog.* 9(7):p.e1003467

38. Barnosky AD, Hadly EA, Bascompte J, Berlow EL, Brown JH, et al. 2012. Approaching a state shift in Earth's biosphere. *Nature* 486(7401):52–58
39. Holmes DJ, Kristan DM. 2008. Comparative and alternative approaches and novel animal models for aging research. *Age* 30(2–3):63–73
40. Munshi-South J, Wilkinson GS. 2010. Bats and birds: exceptional longevity despite high metabolic rates. *Ageing Res. Rev.* 9(1):12–19
41. Austad SN, Fischer KE. 1991. Mammalian aging, metabolism, and ecology: evidence from the bats and marsupials. *J. Gerontol.* 46(2):B47–B53
42. Williams G. 1957. Pleiotropy, natural selection, and the evolution of senescence. *Evolution* 11:398–411
43. Healy K, Guillerme T, Finlay S, Kane A, Kelly SB, et al. 2014. Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc. R. Soc. Lond. B Biol. Sci.* 281(1784):20140298
44. Huang Z, Gallot A, Lao NT, Puechmaile SJ, Foley NM, et al. 2016. A nonlethal sampling method to obtain, generate and assemble whole blood transcriptomes from small, wild mammals. *Mol. Ecol. Resour.* 16(1):150–62
45. Huang Z, Jebb D, Teeling EC. 2016. Blood miRNomes and transcriptomes reveal novel longevity mechanisms in the long-lived bat, *Myotis myotis*. *BMC Genom.* 17(1):906
46. Salminen A, Kaarniranta K, Kauppinen A. 2012. Inflammaging: disturbed interplay between autophagy and inflammasomes. *Ageing* 4(3):166–75
47. Zhang G, Cowled C, Shi Z, Huang Z, Bishop-Lilly KA, et al. 2013. Comparative analysis of bat genomes provides insight into the evolution of flight and immunity. *Science* 339(6118):456–60
48. Franceschi C, Campisi J. 2014. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J. Gerontol. A Biol. Sci. Med. Sci.* 69(Suppl. 1):S4–S9
49. Puechmaile SJ, Frick WF, Kunz TH, Racey PA, Voigt CC, et al. 2011a. White-nose syndrome: Is this emerging disease a threat to European bats? *Trends Ecol. Evol.* 26(11):570–76
50. Leopardi S, Blake D, Puechmaile SJ. 2015. White-nose syndrome fungus introduced from Europe to North America. *Curr. Biol.* 25(6):R217–R219
51. Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, et al. 2010. An emerging disease causes regional population collapse of a common North American bat species. *Science* 329(5992):679–82
52. Puechmaile SJ, Wibbelt G, Korn V, Fuller H, Forget F, et al. 2011b. Pan-European distribution of white-nose syndrome fungus (*Geomyces destructans*) not associated with mass mortality. *PLOS ONE* 6(4):e19167
53. Anthony SJ, Gilardi K, Menachery VD, Goldstein T, Ssebide B, et al. 2017. Further evidence for bats as the evolutionary source of Middle East respiratory syndrome coronavirus. *mBio* 8(2):e00373–17
54. Chen L, Liu B, Yang J, Jin Q. 2014. *DBatVir: The Database of Bat-Associated Viruses*. <http://www.mgc.ac.cn/DBatVir/>
55. Menachery VD, Yount BL Jr., Debink K, Agnihothram S, Gralinski LE, et al. 2015. A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence. *Nat. Med.* 21:1508–13
56. Olival KJ, Hosseini PR, Zambrana-Torrel C, Ross N, Bogich TL, et al. 2017. Host and viral traits predict zoonotic spillover from mammals. *Nature* 546:646–50
57. Taylor DJ, Dittmar K, Ballinger MJ, Bruenn JA. 2011. Evolutionary maintenance of filovirus-like genes in bat genomes. *BMC Evol. Biol.* 11(1):336
58. Maine JJ, Boyles JG. 2015. Bats initiate vital agroecological interactions in corn. *PNAS* 112(40):12438–43
59. Kunz TH, Braun de Torrez E, Bauer D, Lobova T, Fleming TH. 2011. Ecosystem services provided by bats. *Ann. N.Y. Acad. Sci.* 1223(1):1–38
60. Maas B, Clough Y, Tschardt T. 2013. Bats and birds increase crop yield in tropical agroforestry landscapes. *Ecol. Lett.* 16(12):1480–87
61. Vincenot CE, Florens FV, Kingston T. 2017. Can we protect island flying foxes? *Science* 355(6332):1368–70
62. Liu Y, Rossiter SJ, Han X, Cotton JA, Zhang S. 2010. Cetaceans on a molecular fast track to ultrasonic hearing. *Curr. Biol.* 20(20):1834–39
63. Liu Y, Cotton JA, Shen B, Han X, Rossiter SJ, et al. 2010. Convergent sequence evolution between echolocating bats and dolphins. *Curr. Biol.* 20(2):R53–R54


64. Li Y, Liu Z, Shi P, Zhang J. 2010. The hearing gene *Prestin* unites echolocating bats and whales. *Curr. Biol.* 20(2):R55–R56
65. Parker J, Tsagkogeorga G, Cotton JA, Liu Y, Provero P, et al. 2013. Genome-wide signatures of convergent evolution in echolocating mammals. *Nature* 502(7470):228–31
66. Hayden S, Bekaert M, Goodbla A, Murphy WJ, Dávalos LM, et al. 2014. A cluster of olfactory receptor genes linked to frugivory in bats. *Mol. Biol. Evol.* 31(4):917–27
67. Hughes GM, Boston EM, Finarelli JA, Murphy WJ, Higgins DG, et al. 2017. The birth and death of olfaction: the role of olfactory receptor evolution in mammalian niche specialization. *Mol. Biol. Evol.* Manuscript in review
68. Zhao H, Xu D, Zhang S, Zhang J. 2010. Widespread losses of vomeronasal signal transduction in bats. *Mol. Biol. Evol.* 28(1):7–12
69. Hayden S, Teeling EC. 2014. The molecular biology of vertebrate olfaction. *Anat. Rec.* 297(11):2216–26
70. Yohe LR, Abubakar R, Giordano C, Dumont E, Sears KE, et al. 2017. *Trpc2* pseudogenization dynamics in bats reveal ancestral vomeronasal signaling, then pervasive loss. *Evolution* 71(4):923–35
71. Kirwan JD, Bekaert M, Commins JM, Davies KT, Rossiter SJ, et al. 2013. A phylomedicine approach to understanding the evolution of auditory sensory perception and disease in mammals. *Evol. Appl.* 6(3):412–22
72. Davies KT, Maryanto I, Rossiter SJ. 2013. Evolutionary origins of ultrasonic hearing and laryngeal echolocation in bats inferred from morphological analyses of the inner ear. *Front. Zool.* 10(1):2
73. Cremers FP, Collin RW. 2009. Promises and challenges of genetic therapy for blindness. *Lancet* 374(9701):1569–70
74. World Health Organ. 2014. *Visual impairment and blindness*. Fact Sheet 282, World Health Organ., Geneva. <http://www.who.int/mediacentre/factsheets/fs282/en/>
75. World Health Organ. 2017. *Deafness and hearing loss*. Fact Sheet, World Health Organ., Geneva. <http://www.who.int/mediacentre/factsheets/fs300/en/>
76. Kingston T, Rossiter SJ. 2004. Harmonic-hopping in Wallacea's bats. *Nature* 429(6992):654–57
77. Puechmaile SJ, Gouilh MA, Piyapan P, Yokubol M, Mie KM, et al. 2011. The evolution of sensory divergence in the context of limited gene flow in the bumblebee bat. *Nat. Commun.* 2:573
78. Puechmaile SJ, Borissov IM, Zsebok S, Allegrini B, Hizem M, et al. 2014. Female mate choice can drive the evolution of high frequency echolocation in bats: a case study with *Rhinolophus mehelyi*. *PLOS ONE* 9(7):e103452
79. Janik VM, Slater PJ. 1997. Vocal learning in mammals. *Adv. Study. Behav.* 26:59–100
80. Doupe AJ, Kuhl PK. 1999. Birdsong and human speech: common themes and mechanisms. *Annu. Rev. Neurosci.* 22(1):567–631
81. Petkov CI, Jarvis E. 2012. Birds, primates, and spoken language origins: behavioral phenotypes and neurobiological substrates. *Front. Evol. Neurosci.* 4:12
82. Knörnschild M. 2014. Vocal production learning in bats. *Curr. Opin. Neurobiol.* 28:80–85
83. Pfenning AR, Hara E, Whitney O, Rivas MV, Wang R, et al. 2014. Convergent transcriptional specializations in the brains of humans and song-learning birds. *Science* 346(6215):1256846
84. Lovell PV, Wirthlin M, Wilhelm L, Minx P, Lazar NH, et al. 2014. Conserved syntenic clusters of protein coding genes are missing in birds. *Genome Biol.* 15(12):565
85. Whitney O, Pfenning AR, Howard JT, Blatti CA, Liu F, et al. 2014. Core and region-enriched networks of behaviorally regulated genes and the singing genome. *Science* 346(6215):1256780
- 85a. Vernes SC. 2017. What bats have to say about speech and language. *Psychon. Bull. Rev.* 24:111–17
86. Law J, Boyle J, Harris F, Harkness A, Nye C. 2000. Prevalence and natural history of primary speech and language delay: findings from a systematic review of the literature. *Int. J. Lang. Commun. Disord.* 35:165–88
87. Booker BM, Friedrich T, Mason MK, VanderMeer JE, Zhao J, et al. 2016. Bat accelerated regions identify a bat forelimb specific enhancer in the *HoxD* locus. *PLOS Genet.* 12(3):e1005738
88. Kinsella E, Dora N, Mellis D, Lettice L, Deveney P, et al. 2016. Use of a conditional *Ubr5* mutant allele to investigate the role of an N-end rule ubiquitin-protein ligase in hedgehog signalling and embryonic limb development. *PLOS ONE* 11(6):e0157079

89. Eckalbar WL, Schleich SA, Mason MK, Gill Z, Parker AV, et al. 2016. Transcriptomic and epigenomic characterization of the developing bat wing. *Nat. Genet.* 48(5):528
90. Kapusta A, Suh A, Feschotte C. 2017. Dynamics of genome size evolution in birds and mammals. *PNAS* 114(8):E1460–E69
91. Kapusta A, Suh A. 2017. Evolution of bird genomes—a transposon’s-eye view. *Ann. N. Y. Acad. Sci.* 1389(1):164–85
92. Hughes AL, Hughes MK. 1995. Small genomes for better flyers. *Nature* 377(6548):391
93. Ray DA, Feschotte C, Pagan HJ, Smith JD, Pritham EJ, et al. 2008. Multiple waves of recent DNA transposon activity in the bat, *Myotis lucifugus*. *Genome Res.* 18:717–28
94. Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, et al. 2001. Initial sequencing and analysis of the human genome. *Nature* 409(6822):860–921
95. Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, et al. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437(7057):376–80
96. Glenn TC. 2011. Field guide to next-generation DNA sequencers. *Mol. Ecol. Resour.* 11(5):759–69
97. Manley LJ, Ma D, Levine SS. 2016. Monitoring error rates in Illumina sequencing. *J. Biomol. Tech.* 27(4):125–28
98. Lee H, Gurtowski J, Yoo S, Nattestad M, Marcus S, et al. 2016. Third-generation sequencing and the future of genomics. *bioRxiv* 048603
99. Heather JM, Chain B. 2016. The sequence of sequencers: the history of sequencing DNA. *Genomics* 107(1):1–8
100. Rhoads A, Au KF. 2015. PacBio sequencing and its application. *Genom. Proteom. Bioinforma.* 13(5):278–89
101. Gordon D, Huddleston J, Chaisson MJ, Hill CM, Kronenberg ZN, et al. 2016. Long-read sequence assembly of the gorilla genome. *Science* 352(6281):aae0344
102. Shi L, Guo Y, Dong C, Huddleston J, Yang H, et al. 2016. Long-read sequencing and de novo assembly of a Chinese genome. *Nat. Commun.* 7:12065
103. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *Genome Res.* 27:722–36
104. Salmela L, Rivals E. 2014. LorDEC: accurate and efficient long read error correction. *Bioinformatics* 30(24):3506–14
105. Deshpande V, Fung EDK, Pham S, Bafna V. 2013. Cerulean: a hybrid assembly using high throughput short and long reads. In *Algorithms in Bioinformatics*, Vol. 8126, ed. A Darling, J Stoye, pp. 349–63. Berlin: Springer
106. Koren S, Schatz MC, Walenz BP, Martin J, Howard JT, Ganapathy G, et al. Hybrid error correction and *de novo* assembly of single-molecule sequencing reads. *Nat. Biotechnol.* 30:693–700
107. Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, et al. 2011. A high-resolution map of human evolutionary constraint using 29 mammals. *Nature* 478(7370):476–82
108. Simmons NB. 2005. Order Chiroptera. In *Mammal Species of the World: A Taxonomic and Geographic Reference*, ed. DE Wilson, DM Reeder, pp. 312–529. Baltimore, MD: Johns Hopkins Univ. Press. 3rd ed.
109. Tsang SM, Cirranello AL, Bates PJ, Simmons NB. 2016. The roles of taxonomy and systematics in bat conservation. In *Bats in the Anthropocene: Conservation of Bats in a Changing World*, ed. CC Voight, T Kingston, pp. 503–38. New York: Springer Int. 1st ed.
110. Mayr E. 1942. *Systematics and the Origin of Species*. New York: Columbia Univ. Press
111. Hoffmann FG, Owen JG, Baker RJ. 2003. mtDNA perspective of chromosomal diversification and hybridization in Peters’ tent-making bat (*Uroderma bilobatum*: Phyllostomidae). *Mol. Ecol.* 12(11):2981–93
112. Afonso E, Goyadin A-C, Giraudoux P, Farny G. 2017. Investigating hybridization between the two sibling bat species *Myotis myotis* and *M. blythii* from guano in a natural mixed maternity colony. *PLOS ONE* 12(2):e0170534

113. Bachanek J, Postawa T. 2010. Morphological evidence for hybridization in the sister species *Myotis myotis* and *Myotis oxygnathus* (Chiroptera: Vespertilionidae) in the Carpathian Basin. *Acta Chiropterologica* 12(2):439–48
114. Berthier P, Excoffier L, Ruedi M. 2006. Recurrent replacement of mtDNA and cryptic hybridization between two sibling bat species *Myotis myotis* and *Myotis blythii*. *Proc. R. Soc. B Biol. Sci.* 273:3101–9
115. Koubínová D, Irwin N, Hulva P, Koubek P, Zima J. 2013. Hidden diversity in Senegalese bats and associated findings in the systematics of the family Vespertilionidae. *Front. Zool.* 10:48
116. Baker BJ, Bradley RD. 2006. Speciation in mammals and genetic species concept. *J. Mammology* 87(4):643–62
117. Mayer F, von Helversen O. 2001. Cryptic diversity in European bats. *Proc. R. Soc. B Biol. Sci.* 268:1825–32
118. Kacprzyk J, Teeling EC, Kelleher C, Volleth M. 2016. Wing membrane biopsies for bat cytogenetics: finding of $2n = 54$ in Irish *Rhinolophus hipposideros* (Rhinolophidae, Chiroptera, Mammalia) supports two geographically separated chromosomal variants in Europe. *Cytogenet. Genome Res.* 148(4):279–83
119. Eid J, Fehr A, Gray J, Luong K, Lyle J, et al. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323(5910):133–38
120. Clarke J, Wu HC, Jayasinghe L, Patel A, Reid S, et al. 2009. Continuous base identification for single-molecule nanopore DNA sequencing. *Nat. Nanotechnol.* 4(4):265–70
121. Chin CS, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, et al. 2016. Phased diploid genome assembly with single-molecule real-time sequencing. *Nat. Methods* 13(12):1050–54
122. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, et al. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. *bioRxiv* 071282
123. Chaisson MJ, Huddleston J, Dennis MY, Sudmant PH, Malig M, et al. 2015. Resolving the complexity of the human genome using single-molecule sequencing. *Nature* 517(7536):608–11
124. Bishara A, Liu Y, Weng Z, Kashaf-Haghighi D, Newburger DE, et al. 2015. Read clouds uncover variation in complex regions of the human genome. *Genome Res.* 25(10):1570–80
125. Weisenfeld NI, Kumar V, Shah P, Church DM, Jaffe DB. 2017. Direct determination of diploid genome sequences. *Genome Res.* 27(5):757–67
126. Adey A, Kitzman JO, Burton JN, Daza R, Kumar A, et al. 2014. In vitro, long-range sequence information for de novo genome assembly via transposase contiguity. *Genome Res.* 24(12):2041–49
127. Lieberman-Aiden E, Van Berkum NL, Williams L, Imakaev M, Ragoczy T, et al. 2009. Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* 326(5950):289–93
128. Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, et al. 2017. De novo assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science* 356(6333):92–95
129. Koepfli KP, Paten B, O'Brien SJ. 2015. The Genome 10K Project: a way forward. *Annu. Rev. Anim. Biosci.* 3(1):57–111
130. Schwenk K, Padilla DK, Bakken GS, Full RJ. 2009. Grand challenges in organismal biology. *Integr. Comp. Biol.* 49(1):7–14
131. Zhang G, Li C, Li Q, Li B, Larkin DM, et al. 2014. Comparative genomics reveals insights into avian genome evolution and adaptation. *Science* 346(6215):1311–20
132. Jarvis ED, Mirarab S, Aberer AJ, Li B, Houde P, et al. 2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. *Science* 346(6215):1320–31
133. Misof B, Liu S, Meusemann K, Peters RS, Donath A, et al. 2014. Phylogenomics resolves the timing and pattern of insect evolution. *Science* 346(6210):763–67
134. Wickett NJ, Mirarab S, Nguyen N, Warnow T, Carpenter E, et al. 2014. Phylotranscriptomic analysis of the origin and early diversification of land plants. *PNAS* 111(45):E4859–68
135. Foley NM, Springer MS, Teeling EC. 2016. Mammal madness: Is the mammal tree of life not yet resolved? *Philos. Trans. R. Soc. B* 371(1699):20150140
136. Dool SE, Puechmaille SJ, Foley NM, Allegrini B, Bastian A, et al. 2016. Nuclear introns outperform mitochondrial DNA in inter-specific phylogenetic reconstruction: lessons from horseshoe bats (Rhinolophidae: Chiroptera). *Mol. Phylogenet. Evol.* 97:196–212
137. Hahn MW, Nakhleh L. 2016. Irrational exuberance for resolved species trees. *Evolution* 70(1):7–17

138. Platt RN, Faircloth BC, Sullivan KAM, Kieran TJ, Glenn TC, et al. 2017. Conflicting evolutionary histories of the mitochondrial and nuclear genomes in New World *Myotis* bats. *Syst. Biol.* In press
139. Edwards SV, Potter S, Schmitt CJ, Bragg JG, Moritz C. 2016a. Reticulation, divergence, and the phylogeography-phylogenetics continuum. *PNAS* 113(29):8025–32
140. Prum RO, Berv JS, Dornburg A, Field DJ, Townsend JP, et al. 2015. A comprehensive phylogeny of birds (*Aves*) using targeted next-generation DNA sequencing. *Nature* 526:569–73
141. Li H, Durbin R. 2011. Inference of human population history from individual whole-genome sequences. *Nature* 475(7357):493–96
142. Schiffels S, Durbin R. 2014. Inferring human population size and separation history from multiple genome sequences. *Nat. Genet.* 46(8):919–25
143. Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, et al. 2010. A draft sequence of the Neandertal genome. *Science* 328(5979):710–22
144. Gopalakrishnan S, Castruita JAS, Sinding MHS, Kuderna LF, Rääkkönen J, et al. 2017. The wolf reference genome sequence (*Canis lupus lupus*) and its implications for *Canis* spp. population genomics. *BMC Genom.* 18(1):495
145. Ruedi M, Stadelmann B, Gager Y, Douzery EJ, Francis CM, et al. 2013. Molecular phylogenetic reconstructions identify East Asia as the cradle for the evolution of the cosmopolitan genus *Myotis* (Mammalia, Chiroptera). *Mol. Phylogenet. Evol.* 69(3):437–49
146. Moore JH, Asselbergs FW, Williams SM. 2010. Bioinformatics challenges for genome-wide association studies. *Bioinformatics* 26(4):445–55
147. Hsu PD, Lander ES, Zhang F. 2014. Development and applications of CRISPR-Cas9 for genome engineering. *Cell* 157(6):1262–78
148. White SA, Fisher SE, Geschwind DH, Scharff C, Holy TE. 2006. Singing mice, songbirds, and more: models for FOXP2 function and dysfunction in human speech and language. *J. Neurosci.* 26:10376–79
149. Haesler S, Rochefort C, Georgi B, Licznarski P, Osten P, et al. 2007. Incomplete and inaccurate vocal imitation after knockdown of *FoxP2* in songbird basal ganglia nucleus Area X. *PLoS Biol.* 5(12):e321
150. Schulz SB, Haesler S, Scharff C, Rochefort C. 2010. Knockdown of *FoxP2* alters spine density in Area X of the zebra finch. *Genes Brain Behav.* 9(7):732–40
151. Scharff C, Petri J. 2011. Evo-devo, deep homology and *FoxP2*: implications for the evolution of speech and language. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 366(1574):2124–40
152. Wohlgenuth S, Adam I, Scharff C. 2014. *FoxP2* in songbirds. *Curr. Opin. Neurobiol.* 28:86–93
153. Adam I, Mendoza E, Kobalz U, Wohlgenuth S, Scharff C. 2016. *FoxP2* directly regulates the reelin receptor *VLDLR* developmentally and by singing. *Mol. Cell. Neurosci.* 74:96–105
154. Furey TS. 2012. ChIP-seq and beyond: new and improved methodologies to detect and characterize protein–DNA interactions. *Nat. Rev. Genet.* 13(12):840–52
155. Kouzine F, Levens D, Baranello L. 2014. DNA topology and transcription. *Nucleus* 5(3):195–202
156. Scacheri CA, Scacheri PC. 2015. Mutations in the non-coding genome. *Curr. Opin. Pediatr.* 27(6):659–64
157. Platt RN, Vandewege MW, Kern C, Schmidt CJ, Hoffmann FG, et al. 2014. Large numbers of novel miRNAs originate from DNA transposons and are coincident with a large species radiation in bats. *Mol. Biol. Evol.* 31(6):1536–45
158. Baker RJ, Bininda-Emonds OR, Mantilla-Meluk H, Porter CA, Van Den Bussche RA. 2012. Molecular timescale of diversification of feeding strategy and morphology in New World leaf-nosed bats (Phyllostomidae): a phylogenetic perspective. See Reference 29, pp. 385–409
159. Dumont ER, Dávalos LM, Goldberg A, Santana SE, Rex K, et al. 2012. Morphological innovation, diversification and invasion of a new adaptive zone. *Proc. R. Soc. Lond. B Biol. Sci.* 279:1797–805
160. Rojas D, Warsi OM, Dávalos LM. 2016. Bats (Chiroptera: Noctilionoidea) challenge a recent origin of extant neotropical diversity. *Syst. Biol.* 65(3):432–48
161. Foote AD, Liu Y, Thomas GW, Vinař T, Alföldi J, et al. 2015. Convergent evolution of the genomes of marine mammals. *Nat. Genet.* 47(3):272–75
162. Zhou Q, Zhang J, Bachtrog D, An N, Huang Q, et al. 2014. Complex evolutionary trajectories of sex chromosomes across bird taxa. *Science* 346(6215):1246338

163. ENCODE Proj. Consort. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* 489(7414):57–74
164. Chi KR. 2016. The dark side of the human genome. *Nature* 538(7624):275–77
165. Rao SS, Huntley MH, Durand NC, Stamenova EK, Bochkov ID, et al. 2014. A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* 159(7):1665–80
166. Seki R, Li C, Fang Q, Hayashi S, Egawa S, et al. 2017. Functional roles of Aves class-specific *cis*-regulatory elements on macroevolution of bird-specific features. *Nat. Commun.* 8:14229
167. Pennisi E. 2017. Sequencing all life captivates biologists. *Science* 355(6328):89–95
168. Dong D, Lei M, Hua P, Pan YH, Mu S, et al. 2016. The genomes of two bat species with long constant frequency echolocation calls. *Mol. Biol. Evol.* 34:20–34
169. Seim I, Fang X, Xiong Z, Lobanov AV, Huang Z, et al. 2013. Genome analysis reveals insights into physiology and longevity of the Brandt's bat *Myotis brandtii*. *Nat. Commun.* 4:2212
170. Halpin K, Young PL, Field HE, Mackenzie JS. 2000. Isolation of Hendra virus from pteropid bats: a natural reservoir of Hendra virus. *J. Gen. Virol.* 81:1927–32
171. Halpin K, Hyatt AD, Fogarty R, Middleton D, Bingham J, et al. 2011. Pteropid bats are confirmed as the reservoir hosts of henipaviruses: a comprehensive experimental study of virus transmission. *Am. J. Trop. Med. Hyg.* 85:946–51
172. Weir LD, Annand JE, Reid AP, Broder CC. 2014. Recent observations on Australian bat lyssavirus tropism and viral entry. *Viruses* 6:909–26
173. Eggerbauer E, Pfaff F, Finke S, Höper D, Beer M, et al. 2017. Comparative analysis of European bat lyssavirus 1 pathogenicity in the mouse model. *PLOS Negl. Trop. Dis.* 11:e0005668
174. Towner JS, Pourrut X, Albariño CG, Nkogue CN, Bird BH, et al. 2007. Marburg virus infection detected in a common African bat. *PLOS ONE* 2:e764
175. Ge X-Y, Li J-L, Yang X-L, Chmura AA, Zhu G, et al. 2013. Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* 503:535–38
176. Ithete NL, Stoffberg S, Corman VM, Cottontail VM, Richards LR, et al. 2013. Close relative of human middle east respiratory syndrome coronavirus in bat, South Africa. *Emerg. Infect. Dis. J.* 19:1697



New From Annual Reviews:

Annual Review of Cancer Biology

cancerbio.annualreviews.org • Volume 1 • March 2017

ONLINE NOW!


Co-Editors: **Tyler Jacks**, *Massachusetts Institute of Technology*

Charles L. Sawyers, *Memorial Sloan Kettering Cancer Center*

The *Annual Review of Cancer Biology* reviews a range of subjects representing important and emerging areas in the field of cancer research. The *Annual Review of Cancer Biology* includes three broad themes: Cancer Cell Biology, Tumorigenesis and Cancer Progression, and Translational Cancer Science.

TABLE OF CONTENTS FOR VOLUME 1:

- *How Tumor Virology Evolved into Cancer Biology and Transformed Oncology*, Harold Varmus 
- *The Role of Autophagy in Cancer*, Naiara Santana-Codina, Joseph D. Mancias, Alec C. Kimmelman
- *Cell Cycle-Targeted Cancer Therapies*, Charles J. Sherr, Jiri Bartek
- *Ubiquitin in Cell-Cycle Regulation and Dysregulation in Cancer*, Natalie A. Borg, Vishva M. Dixit
- *The Two Faces of Reactive Oxygen Species in Cancer*, Colleen R. Reczek, Navdeep S. Chandel
- *Analyzing Tumor Metabolism In Vivo*, Brandon Faubert, Ralph J. DeBerardinis
- *Stress-Induced Mutagenesis: Implications in Cancer and Drug Resistance*, Devon M. Fitzgerald, P.J. Hastings, Susan M. Rosenberg
- *Synthetic Lethality in Cancer Therapeutics*, Roderick L. Beijersbergen, Lodewyk F.A. Wessels, René Bernards
- *Noncoding RNAs in Cancer Development*, Chao-Po Lin, Lin He
- *p53: Multiple Facets of a Rubik's Cube*, Yun Zhang, Guillermina Lozano
- *Resisting Resistance*, Ivana Bozic, Martin A. Nowak
- *Deciphering Genetic Intratumor Heterogeneity and Its Impact on Cancer Evolution*, Rachel Rosenthal, Nicholas McGranahan, Javier Herrero, Charles Swanton
- *Immune-Suppressing Cellular Elements of the Tumor Microenvironment*, Douglas T. Fearon
- *Overcoming On-Target Resistance to Tyrosine Kinase Inhibitors in Lung Cancer*, Ibiayi Dagogo-Jack, Jeffrey A. Engelman, Alice T. Shaw
- *Apoptosis and Cancer*, Anthony Letai
- *Chemical Carcinogenesis Models of Cancer: Back to the Future*, Melissa Q. McCreery, Allan Balmain
- *Extracellular Matrix Remodeling and Stiffening Modulate Tumor Phenotype and Treatment Response*, Jennifer L. Leight, Allison P. Drain, Valerie M. Weaver
- *Aneuploidy in Cancer: Seq-ing Answers to Old Questions*, Kristin A. Knouse, Teresa Davoli, Stephen J. Elledge, Angelika Amon
- *The Role of Chromatin-Associated Proteins in Cancer*, Kristian Helin, Saverio Minucci
- *Targeted Differentiation Therapy with Mutant IDH Inhibitors: Early Experiences and Parallels with Other Differentiation Agents*, Eytan Stein, Katharine Yen
- *Determinants of Organotropic Metastasis*, Heath A. Smith, Yibin Kang
- *Multiple Roles for the MLL/COMPASS Family in the Epigenetic Regulation of Gene Expression and in Cancer*, Joshua J. Meeks, Ali Shilatifard
- *Chimeric Antigen Receptors: A Paradigm Shift in Immunotherapy*, Michel Sadelain



Contents

Weird Animals, Sex, and Genome Evolution <i>Jennifer A. Marshall Graves</i>	1
Bat Biology, Genomes, and the Bat1K Project: To Generate Chromosome-Level Genomes for All Living Bat Species <i>Emma C. Teeling, Sonja C. Vernes, Liliana M. Dávalos, David A. Ray, M. Thomas P. Gilbert, Eugene Myers, and Bat1K Consortium</i>	23
The Divergent Genomes of Teleosts <i>Vyidianathan Ravi and Byrappa Venkatesh</i>	47
Biobanking Genetic Material for Agricultural Animal Species <i>H.D. Blackburn</i>	69
Viable Cell Culture Banking for Biodiversity Characterization and Conservation <i>Oliver A. Ryder and Manabu Onuma</i>	83
Lessons from the Barn to the Operating Suite: A Comprehensive Review of Animal Models for Fetal Surgery <i>Sandra K. Kabagambe, Chelsey J. Lee, Laura F. Goodman, Y. Julia Chen, Melissa A. Vanover, and Diana L. Farmer</i>	99
Detection of Emerging Zoonotic Pathogens: An Integrated One Health Approach <i>Brian H. Bird and Jonna A.K. Mazet</i>	121
Development and Function of the Mucosal Immune System in the Upper Respiratory Tract of Neonatal Calves <i>Rabwa Osman, Nilusha Malmuthuge, Patricia Gonzalez-Cano, and Philip Griebel</i> ...	141
Equine Intestinal Mucosal Pathobiology <i>Anthony Blikslager and Liara Gonzalez</i>	157
Metabolic and Endocrine Role of Adipose Tissue During Lactation <i>J.P. McNamara and K. Huber</i>	177
Nutrient-Sensing Biology in Mammals and Birds <i>Eugeni Roura and Simon R. Foster</i>	197

Impact of Micronutrients on the Immune Response of Animals <i>Allen D. Smith, Kiran S. Panickar, Joseph F. Urban, Jr., and Harry D. Dawson</i>	227
Measuring Male-to-Male Differences in Fertility or Effects of Semen Treatments <i>Rupert P. Amann, Richard G. Saacke, Guy F. Barbato, and Dagmar Waberski</i>	255
Reproductive Impacts of Endocrine-Disrupting Chemicals on Wildlife Species: Implications for Conservation of Endangered Species <i>Christopher W. Tubbs and Caitlin E. McDonough</i>	287
Catfish Biology and Farming <i>Rex A. Dunham and Ahmed Elaswad</i>	305

Errata

An online log of corrections to *Annual Review of Animal Biosciences* articles may be found at <http://www.annualreviews.org/errata/animal>