

Massive spin 2 excitations in $AdS_6 \times S^2$ warped spacetimes

Michael Gutperle,^a Christoph F. Uhlemann^a and Oscar Varela^{b,c,d}

^a*Mani L. Bhaumik Institute for Theoretical Physics, Department of Physics and Astronomy, University of California, 475 Portola Plaza, Los Angeles, CA, 90095 U.S.A.*

^b*Department of Physics, Utah State University, Logan, UT, 84322 U.S.A.*

^c*Departamento de Física Teórica and Instituto de Física Teórica UAM/CSIC, Universidad Autónoma de Madrid, Cantoblanco, Madrid, 28049 Spain*

^d*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, Potsdam, D-14476 Germany*

E-mail: gutperle@ucla.edu, uhlemann@physics.ucla.edu, oscar.varela@usu.edu

ABSTRACT: We study (massive) spin-2 fluctuations around warped AdS_6 solutions with 16 supersymmetries in type IIB supergravity. We identify two classes of fluctuations which are universally present for all solutions of this form. The holographically dual operators have scaling dimensions $\Delta = 5 + 3\ell$ and $\Delta = 6 + 3\ell$, where the integer ℓ encodes the R -symmetry charge. They are identified as descendant states in respective BPS multiplets (the current multiplet for $\Delta = 5$). We also compute the normalization of the energy-momentum tensor two-point function and show that it is related to the S^5 partition function of the dual $5d$ SCFTs.

KEYWORDS: AdS-CFT Correspondence, Gauge-gravity correspondence

ARXIV EPRINT: [1805.11914](https://arxiv.org/abs/1805.11914)

Contents

1	Introduction	1
2	Type IIB $AdS_6 \times S^2$ warped solution	2
3	The spectrum of spin 2 excitations	3
3.1	Regularity conditions	5
3.2	Universal minimal solutions	5
3.3	Universal non-minimal solutions	6
4	Implications for the dual SCFTs	7
4.1	Superconformal multiplets	7
4.2	Normalization of two-point functions	8
5	Discussion	10
A	T-dual of massive IIA solution	11
A.1	Metric perturbations	11

1 Introduction

Supersymmetric conformal field theories (SCFT) in five dimensions exhibit many interesting properties: they are intrinsically strongly coupled in the UV and many have relevant deformations that flow to non-renormalizable gauge theories in the IR. Generically, they do not have exactly marginal deformations and no conventional Lagrangian description. These features are challenging for (perturbative) QFT methods, but the existence and properties of such theories can be studied using string theory and M-theory [1–3].

AdS/CFT is a particularly powerful tool to quantitatively study these theories, and supergravity solutions describing five dimensional SCFTs have been found in type IIA [4–6] and type IIB supergravity [7–9].¹ The type IIB solutions were identified with a large class of SCFTs which are obtained from the conformal limit of (p, q) 5-brane webs [17, 18]. The IIB supergravity solutions have since been studied and extended. In [21] the S^5 partition function and entanglement entropy for a spherical region were calculated holographically. In [22] probe (p, q) strings in the supergravity background have been studied. In [19] supergravity solutions were constructed with punctures and non-trivial $SL(2, \mathbb{R})$ monodromy, and they were related in a precise way to 5-brane webs with additional 7-branes in [20].

An important part of the basic data of any SCFT is the spectrum of short and long operators and how they fit into representations of the superconformal algebra. It is well

¹See [11–16] for earlier work on type IIB solutions.

known that five dimensional superconformal symmetry is a special case, with a unique superconformal algebra $F(4)$ [23–25]. Recently, the superconformal multiplets of $F(4)$ have been classified in [26, 27]. On general grounds, it is expected that linearized fluctuations of the supergravity fields around an AdS vacuum fall into superconformal multiplets, with the complete spectrum determined by the Kaluza-Klein (KK) reduction. A famous example is the KK-reduction of type IIB supergravity on its $AdS_5 \times S^5$ vacuum [28, 29] and the relation to the spectrum of half-BPS operators of $\mathcal{N} = 4$ SYM. The AdS_6 solutions of type IIB supergravity are, however, more complicated: they only preserve half the number of supersymmetries, the metric is a more involved warped product of $AdS_6 \times S^2$ over a Riemann surface, and they involve non-trivial configurations of the axion-dilaton scalar and complex two-form field. A general analysis of the coupled equations of motion and the resulting KK spectrum for these warped compactifications is therefore challenging, and we instead focus on a special class of fluctuations which can be studied by itself.

The structure of the present paper is as follows: in section 2 we briefly review the type IIB solutions of [7–9, 19] and their relation to 5-brane webs. In section 3 we study fluctuations of the metric field which correspond to massless or massive symmetric, transverse and traceless rank two tensors on AdS_6 . Crucially, the linearized equation of motion for these fluctuations decouples from all other supergravity fields [30], and we identify a particular class of solutions which is universally present for the type IIB AdS_6 solutions discussed above. As we discuss in section 4, in the dual SCFTs these fluctuations source the stress tensor (in the massless case) as well as spin two tensor operators of higher scaling dimension. We explicitly obtain the “central charge” C_T which appears in the two point function of the stress tensor from our supergravity fluctuation. We close with a discussion in section 5.

2 Type IIB $AdS_6 \times S^2$ warped solution

In this section we review aspects of the supergravity solutions constructed in [7] which will be important in the present paper. These solutions provide the general local form of warped AdS_6 solutions with 16 supersymmetries in type IIB supergravity. The geometry is a warped product of $AdS_6 \times S^2$ over a Riemann surface Σ . The complex scalar and the complex two-form potential (proportional to the volume form on S^2) depend on the coordinates on Σ . The metric is given by

$$ds^2 = f_6^2 ds_{AdS_6}^2 + f_2^2 ds_{S^2}^2 + 4\rho^2 |dw|^2. \tag{2.1}$$

The metric functions and the remaining supergravity fields are all expressed in terms of two locally holomorphic functions \mathcal{A}_\pm on Σ , via the composite quantities

$$\begin{aligned} \kappa^2 &= -|\partial_w \mathcal{A}_+|^2 + |\partial_w \mathcal{A}_-|^2, & \partial_w \mathcal{B} &= \mathcal{A}_+ \partial_w \mathcal{A}_- - \mathcal{A}_- \partial_w \mathcal{A}_+, \\ \mathcal{G} &= |\mathcal{A}_+|^2 - |\mathcal{A}_-|^2 + \mathcal{B} + \bar{\mathcal{B}}, & R + \frac{1}{R} &= 2 + \frac{6 \kappa^2 \mathcal{G}}{|\partial_w \mathcal{G}|^2}. \end{aligned} \tag{2.2}$$

Note that \mathcal{G} and κ^2 are related as follows

$$\partial_{\bar{w}} \partial_w \mathcal{G} = -\kappa^2. \tag{2.3}$$

In Einstein frame, the metric functions f_6^2 , f_2^2 and ρ^2 are given by

$$f_6^2 = \sqrt{6\mathcal{G}} \left(\frac{1+R}{1-R} \right)^{\frac{1}{2}}, \quad f_2^2 = \frac{1}{9} \sqrt{6\mathcal{G}} \left(\frac{1-R}{1+R} \right)^{\frac{3}{2}}, \quad \rho^2 = \frac{\kappa^2}{\sqrt{6\mathcal{G}}} \left(\frac{1+R}{1-R} \right)^{\frac{1}{2}}. \quad (2.4)$$

The expressions for the axion-dilaton scalar and the complex two-form $C_{(2)}$ can be found in [7], but are not needed in the following. For arbitrary locally holomorphic \mathcal{A}_\pm this yields a solution to the BPS equations (and also to the equations of motion [10]). For physically regular solutions, κ^2 and \mathcal{G} satisfy

$$\kappa^2 > 0, \quad \mathcal{G} > 0, \quad (2.5)$$

at interior points of Σ . If Σ has a boundary, they further satisfy the boundary conditions

$$\kappa^2|_\Sigma = \mathcal{G}|_\Sigma = 0, \quad (2.6)$$

which ensure that the S^2 collapses on $\partial\Sigma$ to smoothly close off the $10d$ geometry. Large classes of physically regular solutions were constructed in [8, 9, 19], and related to (p, q) 5-brane webs in type IIB string theory. The Riemann surface Σ is a disc in these solutions, and the differentials $\partial_w \mathcal{A}_\pm$ have poles at isolated points on the boundary of the disc. The technical details of these constructions will be largely irrelevant for the discussion in the present paper. The results hold in particular for these solutions, but are not limited to them. We will make use of the regularity conditions (2.5) and (2.6), and discuss the poles where necessary, e.g. in relation to regularity conditions in section 3.1 and 3.3.

3 The spectrum of spin 2 excitations

As discussed in the introduction, determining the complete KK-spectrum of a warped AdS supergravity solution is challenging. However, as shown in [30], the spectrum of spin-2 fluctuations is determined from a universal equation which depends on the geometry only, not on the other fields in type IIB supergravity. The original motivation of [30] was to study localized spin 2 excitations on solutions dual to defect conformal field theories. The result on the decoupling, however, can be applied to general warped spacetimes with AdS factors, see e.g. [31–36] for application in various cases.

While the formulae in [30] were for AdS_4 , it is straightforward to generalize the analysis to the case which we are interested in, namely the fluctuations around the warped $AdS_6 \times S^2 \times \Sigma$ solutions. The perturbed $10d$ metric takes the form

$$ds^2 = f_6^2 (ds_{AdS_6}^2 + h_{\mu\nu} dx^\mu dx^\nu) + \hat{g}_{ab} dy^a dy^b, \quad (3.1)$$

with $h_{\mu\nu}$ a transverse traceless fluctuation in the unit-radius AdS_6 part of the background geometry, and the metric on the 4-dimensional internal space in (2.1) given by

$$\hat{g}_{ab} dy^a dy^b = f_2^2 ds_{S^2}^2 + 4\rho^2 |dw|^2. \quad (3.2)$$

The $6d$ part of the metric perturbation is expanded in modes on unit-radius AdS_6 ,

$$h_{\mu\nu}(x, y) = h_{\mu\nu}^{[tt]}(x) \psi(y), \quad \square_{AdS_6} h_{\mu\nu}^{[tt]} = (M^2 - 2) h_{\mu\nu}^{[tt]}. \quad (3.3)$$

The starting point is eq. (2.23) of [30], which holds for a non-compact part of arbitrary dimension. With g denoting the full $10d$ metric and $M, N = 0, \dots, 9$, it is given by

$$\frac{1}{\sqrt{-g}} \partial_M \sqrt{-g} g^{MN} \partial_N h_{\mu\nu} = 0. \quad (3.4)$$

With the mode expansion (3.3), this reduces to an equation for ψ , given by

$$-\frac{1}{f_6^4 \sqrt{\hat{g}}} \partial_a (f_6^6 \sqrt{\hat{g}} \hat{g}^{ab} \partial_b) \psi = M^2 \psi. \quad (3.5)$$

The only effect of having AdS_6 instead of AdS_4 are the different powers of the AdS warp factor as compared to [30]. With the explicit form of the internal space metric in (3.2),² eq. (3.5) evaluates to

$$\frac{1}{f_6^4 f_2^2 \rho^2} \partial_a (f_6^6 f_2^2 \eta^{ab} \partial_b \psi) + \frac{f_6^2}{f_2^2} \nabla_{S^2}^2 \psi + M^2 \psi = 0, \quad (3.6)$$

where $a, b = w, \bar{w}$ from now on and

$$\eta^{w\bar{w}} = \eta^{\bar{w}w} = \frac{1}{2}. \quad (3.7)$$

Somewhat remarkably, the composite quantity R cancels in all the combinations of metric factors, which are given explicitly by

$$\frac{f_6^2}{f_2^2} = 9 + \frac{6|\partial\mathcal{G}|^2}{\kappa^2 \mathcal{G}}, \quad f_6^4 f_2^2 \rho^2 = \frac{2}{3} \kappa^2 \mathcal{G}, \quad f_2 f_6^3 = 2\mathcal{G}. \quad (3.8)$$

We further expand ψ in spherical harmonics on S^2

$$\psi(y) = \phi_{\ell m}(w, \bar{w}) Y_{\ell m}(S^2). \quad (3.9)$$

This turns eq. (3.6) into

$$6\partial_a (\mathcal{G}^2 \eta^{ab} \partial_b \phi_{\ell m}) - \ell(\ell+1) (9\kappa^2 \mathcal{G} + 6|\partial\mathcal{G}|^2) \phi_{\ell m} + M^2 \kappa^2 \mathcal{G} \phi_{\ell m} = 0. \quad (3.10)$$

This equation can be further simplified with the field redefinition

$$\phi_{\ell m} = \mathcal{G}^\ell \chi_{\ell m}. \quad (3.11)$$

After multiplying (3.10) by \mathcal{G}^ℓ , it becomes

$$\partial_a \left(\mathcal{G}^{2\ell+2} \partial^a \chi_{\ell m} \right) + \frac{1}{6} (M^2 - 3\ell(3\ell+5)) \kappa^2 \mathcal{G}^{2\ell+1} \chi_{\ell m} = 0. \quad (3.12)$$

This equation determines the transverse traceless spin-2 fluctuations around generic type IIB warped $AdS_6 \times S^2 \times \Sigma$ solutions reviewed in section 2.

²The Σ part of the metric in (2.1) may be written more explicitly as $g_\Sigma = 2\rho^2(dw \otimes d\bar{w} + d\bar{w} \otimes dw)$, from which the components of the metric and inverse metric are extracted.

3.1 Regularity conditions

We implement the following regularity requirements for the metric fluctuations. In the interior of Σ , the perturbation should be finite. At generic points on the boundary of Σ , $\partial\Sigma$, the S^2 closes off, as ensured by the boundary condition (2.6). These boundary points are completely regular points in the $10d$ geometry, and as such the metric fluctuation at these points should be finite. Locally, coordinates can be chosen such that Σ corresponds to the upper half plane, with the real line as the boundary. In view of the decomposition (3.9) and (3.11), the requirement for finiteness of the metric perturbation then implies

$$\lim_{\text{Im}(w) \rightarrow 0} \mathcal{G}^\ell \chi_{\ell m} < \infty. \tag{3.13}$$

For the solutions constructed in [7–9], the poles on the boundary $\partial\Sigma$, at which the external (p, q) 5-branes of the associated 5-brane web emerge, call for special attention. We will insist that the metric fluctuation (3.1) does not change the asymptotic behavior of the AdS_6 radius near the poles. This ensures that the poles in the perturbed solution still admit an interpretation as (p, q) 5-branes. The AdS_6 radius in Einstein frame vanishes near the poles, $f_6^2 \rightarrow 0$, as shown in section 3.9 of [9], while it diverges in string frame. Demanding this behavior to remain unmodified simply amounts to requiring (3.13) also at the poles: as long as $\phi_{\ell m}$ is finite at the poles, the AdS_6 radius in the perturbed Einstein-frame metric still vanishes at the same rate as the poles are approached, due to the overall f_6^2 in the ansatz for the metric perturbation (3.1).

For the solutions of [19] there are additional punctures in Σ , around which the supergravity fields undergo non-trivial $SL(2, \mathbb{R})$ monodromy and which encode additional 7-branes in the associated 5-brane web. The Einstein-frame metric, however, is single-valued. We keep the requirement that the metric fluctuation be finite all through the interior of Σ , which ensures that the interpretation of the punctures as 7-branes is unmodified. Since \mathcal{G} is single-valued and also finite all through Σ for these solutions, the punctures then do not play a special role in the fluctuation equation (3.12).

3.2 Universal minimal solutions

Constructing the general solution to the partial differential equation (3.12) is, still, challenging. However, with the equation in this particular form, we can immediately identify a particularly simple class of solutions, given by

$$\chi_{\ell m} = 1, \quad \phi_{\ell m} = \mathcal{G}^\ell, \quad M^2 = 3\ell(3\ell + 5). \tag{3.14}$$

Since \mathcal{G} vanishes on all of $\partial\Sigma$ by (2.6), the regularity condition (3.13) excludes such constant solutions for $\ell < 0$ (which would not be regular on S^2 to begin with). But for $\ell \geq 0$ these solutions satisfy the regularity conditions discussed in section 3.1. This special class of solutions, in fact, provides the solution with minimal M^2 for each ℓ : multiplying eq. (3.12) by $\chi_{\ell m}$ and integrating over Σ yields

$$\int_{\Sigma} d^2w \left[\chi_{\ell m} \partial_a \left(\mathcal{G}^{2\ell+2} \partial^a \chi_{\ell m} \right) + \frac{1}{6} \left(M^2 - 3\ell(3\ell + 5) \right) \kappa^2 \mathcal{G}^{2\ell+1} \chi_{\ell m}^2 \right] = 0. \tag{3.15}$$

Integrating the first term by parts yields a boundary term $\propto \chi_{\ell m} \mathcal{G}^{2\ell+2} \partial^a \chi_{\ell m}$, which vanishes by (3.13). We thus find, using that $\chi_{\ell m}$ is real,

$$\int_{\Sigma} d^2 w \left[-\mathcal{G}^{2\ell+2} |\partial \chi_{\ell m}|^2 + \frac{1}{6} (M^2 - 3\ell(3\ell + 5)) \kappa^2 \mathcal{G}^{2\ell+1} \chi_{\ell m}^2 \right] = 0. \quad (3.16)$$

Noting that κ^2 and \mathcal{G} are non-negative in all of Σ and positive in the interior, we conclude that

$$M^2 \geq 3\ell(3\ell + 5). \quad (3.17)$$

The solutions in (3.14) saturate this inequality and therefore indeed produce minimal M^2 for a given choice of ℓ, m .

3.3 Universal non-minimal solutions

In this section we discuss a second class of solutions which is also universally present but produces non-minimal M^2 . It is reminiscent of a second branch of solutions found for the type IIA solution in [36], but differs in crucial details. As starting point we take eq. (3.13), and rewrite it, noting (3.7), as follows

$$\mathcal{G} \partial_w \partial_{\bar{w}} \chi_{\ell m} + (2\ell + 2) (\partial_a \mathcal{G}) \partial^a \chi_{\ell m} + \frac{1}{6} (M^2 - 3\ell(3\ell + 5)) \kappa^2 \chi_{\ell m} = 0. \quad (3.18)$$

We make an ansatz for $\chi_{\ell m}$ as the difference of a holomorphic and an anti-holomorphic part,

$$\chi_{\ell m}(w, \bar{w}) = \chi_{\ell m}^+(w) - \chi_{\ell m}^-(\bar{w}). \quad (3.19)$$

In particular, this makes $\chi_{\ell m}$ harmonic, and the first term in (3.18) drops out. Evaluating the remaining terms more explicitly yields

$$(\ell + 1) ((\partial_{\bar{w}} \mathcal{G}) \partial_w \chi_{\ell m}^+ - (\partial_w \mathcal{G}) \partial_{\bar{w}} \chi_{\ell m}^-) + \frac{1}{6} (M^2 - 3\ell(3\ell + 5)) \kappa^2 (\chi_{\ell m}^+ - \chi_{\ell m}^-) = 0. \quad (3.20)$$

Using the explicit expression for $\partial_w \mathcal{G}$, given by

$$\partial_w \mathcal{G} = (\bar{\mathcal{A}}_+ - \mathcal{A}_-) \partial_w \mathcal{A}_+ + (\mathcal{A}_+ - \bar{\mathcal{A}}_-) \partial_w \mathcal{A}_-, \quad (3.21)$$

it is straightforward to show that eq. (3.20) admits a solution when $\chi_{\ell m}^+ = \mathcal{A}_+$ and $\chi_{\ell m}^- = \bar{\mathcal{A}}_-$. It is given by

$$\chi_{\ell m} = \mathcal{A}_+ - \bar{\mathcal{A}}_-, \quad M^2 = (3\ell + 1)(3\ell + 6). \quad (3.22)$$

Note that these $\chi_{\ell m}$ are not real, and the linearized equation is solved by the real and imaginary parts separately. They are regular in the interior of Σ for the solutions of [8, 9, 19]. The \mathcal{A}_{\pm} have logarithmic divergences at the poles on the boundary of Σ , and associated branch cuts along the boundary. The discontinuities across the branch cuts, however, cancel in the combination $\mathcal{A}_+ - \bar{\mathcal{A}}_-$, such that the $\chi_{\ell m}$ are single valued. This leaves only the logarithmic divergences at the poles. Crucially, the profile of the metric

fluctuation is given by $\phi_{\ell m}$ as defined in (3.11), not by $\chi_{\ell m}$ directly. The solutions for $\phi_{\ell m}$ are

$$\phi_{\ell m}^{(1)} = \mathcal{G}^\ell \operatorname{Re}(\mathcal{A}_+ - \bar{\mathcal{A}}_-), \quad \phi_{\ell m}^{(2)} = \mathcal{G}^\ell \operatorname{Im}(\mathcal{A}_+ - \bar{\mathcal{A}}_-). \quad (3.23)$$

For $\ell = 0$ the logarithmic singularity at the poles remains.³ For $\ell \geq 1$, however, since \mathcal{G} vanishes on the boundary of Σ and at the poles (faster than logarithmically, as can be seen from the near-pole behavior derived in section 3.9 of [7]), these solutions are perfectly regular.

4 Implications for the dual SCFTs

In this section we discuss the operators that the universal spin-2 fluctuations identified above source in the dual 5d SCFTs, and the superconformal multiplets they belong to. We also obtain the normalization of the two-point functions for these operators and discuss the relation to the five-sphere partition function.

4.1 Superconformal multiplets

For each of the graviton perturbations discussed in the previous section there is an operator in the spectrum of the dual SCFT, which we will call $T_{\ell, m}^{\mu\nu}$. From the usual relation between the AdS mass of a bulk field and the scaling dimension of the dual operator, $M^2 = \Delta(\Delta - 5)$, we conclude that the dual operators for the minimal solutions (3.14) have scaling dimension

$$\Delta(T_{\min, \ell, m}^{\mu\nu}) = 5 + 3\ell. \quad (4.1)$$

To identify the multiplets that these dual operators correspond to we consult with the classification of [27].⁴ We first note that the Dynkin label R in the convention of [27] is always integer, rather than half integer, and related to our ℓ by $R = 2\ell$.⁵ In our type IIB solution the R-symmetry is realized as the isometries of the S^2 and the scalar spherical harmonics on S^2 have integer ℓ , this implies that R takes even integer values. The fluctuation spectrum of the supergravity solution contains states of up to spin 2. The only short multiplets in table 22 of [27] that fulfill this requirement are B_2 and A_4 . The B_2 representation with $R = 0$ reproduces the $6d$ (gauged) supergravity multiplet. This multiplet should be present in the spectrum, as we expect our type IIB supergravity solutions to admit a truncation to $6d$ gauged supergravity. The arguments of [37] actually suggest the stronger statement that there is a truncation which is consistent at the full non-linear level. The linear analysis is sufficient for our purposes. From the perspective of the dual SCFTs, having a $6d$ (gauged) supergravity multiplet in the spectrum simply amounts to the field theories having an energy-momentum tensor.

³The complete metric perturbation (3.1) in Einstein frame is finite at the poles, due to the factor f_6^2 . But the perturbation dominates the AdS_6 part of the background metric and actually blows up in string frame.

⁴For definiteness, we use the notation and conventions of [27] instead of the slightly different ones in [26] to denote the multiplets and representations.

⁵The supercharge in their eq. (2.56) has $R = 1$, but, being a spinor, would correspond to $\ell = 1/2$.

The natural $F(4)$ supermultiplet candidate to host the operators $T_{\ell,m}^{\mu\nu}$ is thus B_2 . From [27], the structure of the B_2 supercurrent multiplet is

$$\boxed{[0, 0]_3^{(0)}} \xrightarrow{Q} \boxed{[1, 0]_{\frac{7}{2}}^{(0)}} \xrightarrow{Q} \boxed{[0, 1]_4^{(2)} \oplus [2, 0]_4^{(0)}} \xrightarrow{Q} \boxed{[1, 1]_{\frac{9}{2}}^{(1)}} \xrightarrow{Q} \boxed{[0, 2]_5^{(0)}}. \quad (4.2)$$

The primary in this multiplet $[0, 0]_3^{(0)}$ is a Lorentz scalar with R-charge $R = 0$ and scaling dimension $\Delta = 3$. Note that the operator dual to the massless graviton fluctuation, $[0, 2]_5^{(0)}$, arises as the Q^4 descendant in this multiplet. Generalizing this structure, the minimal massive spin two fluctuations found in section 3.2 with $\ell > 0$ fit into a B_2 multiplet where the scalar primary $[0, 0]_{3+3\ell}^{(2\ell)}$ has scaling dimension

$$\Delta_{B_2, \text{primary}} = 3 + 3\ell. \quad (4.3)$$

The massive spin-2 fluctuations can again be identified as the Q^4 descendant $[0, 2]_{5+3\ell}^{2\ell}$. That means the operators dual to the massive spin 2 fluctuations are not primaries, but descendants in this multiplet, with the Q^4 precisely accounting for the difference of 2 between the scaling dimensions (4.1) and (4.3).

The universal non-minimal fluctuation, presented in section 3.3, is dual to a spin-2 operator with dimension

$$\Delta(T_{\text{non min}, \ell, m}^{\mu\nu}) = 6 + 3\ell. \quad (4.4)$$

Following the same logic as for the minimal solution we conclude that this operator is part of an A_4 multiplet with $R = 3\ell$, where it is again a Q^4 descendant of a scalar primary $[0, 0]_{4+3\ell}^{(2\ell)}$ with conformal dimension

$$\Delta_{A_4, \text{primary}} = 4 + 3\ell. \quad (4.5)$$

4.2 Normalization of two-point functions

We now compute the normalization of the two-point functions of the operators $T_{\ell,m}^{\mu\nu}$ sourced by the universal graviton fluctuations of section 3.2 and 3.3. For the minimal solution with $\ell = 0$, corresponding to the massless graviton fluctuation, the dual operator is the energy-momentum tensor, whose normalization can be fixed independently, such that the normalization of the two-point function becomes physically meaningful. This is what [38] called the central charge C_T , which is related to the partition function on squashed spheres and therefore accessible for field theory computations using supersymmetric localization.

Holographically, the normalization of the energy-momentum tensor two-point function can be extracted from the effective action for the $6d$ graviton. We start from the Einstein-frame action as given e.g. in [39], whose form we only need schematically,

$$S_{\text{IIB}} = \frac{1}{2\kappa_{10}^2} \int d^{10}x \sqrt{-g} (R + \dots). \quad (4.6)$$

On general grounds, expanding the action (4.6) to quadratic order in a perturbation δg yields

$$\delta^2 S_{\text{IIB}} = \frac{1}{\kappa_{10}^2} \int d^{10}x \sqrt{-g} \delta g^{MN} P_{MN}[\delta g] + \text{boundary terms}, \quad (4.7)$$

where $P_{MN}[\delta g]$ is the equation of motion operator acting on δg , and the indices in δg^{MN} are raised with the full $10d$ background metric. $P[\delta g]$ is given by (3.4), up to a factor f_6^2 owing to the fact that it was derived in [30] for the Weyl rescaled metric where the AdS factor has unit radius. Noting also the explicit f_6^2 in the perturbation ansatz (3.1), we have

$$P_{\mu\nu}[\delta g] = f_6^2 \frac{1}{\sqrt{-g}} \partial_M \sqrt{-g} g^{MN} \partial_N h_{\mu\nu}, \quad \delta g^{\mu\nu} = f_6^{-2} h^{\mu\nu}, \quad (4.8)$$

where the indices on h are now raised with the unit-radius AdS_6 metric. Thus, dropping boundary terms,

$$\delta^2 S_{\text{IIB}} = \frac{1}{\kappa_{10}^2} \int d^{10}x h^{\mu\nu} \partial_M \sqrt{-g} g^{MN} \partial_N h_{\mu\nu}. \quad (4.9)$$

Using the expansion $h_{\mu\nu} = (h_{\ell m}^{[tt]})_{\mu\nu} Y_{\ell m} \phi_{\ell m}$, with normalization $\int_{S^2} Y_{\ell m} Y_{\ell' m'} = \text{Vol}_{S^2} \delta_{\ell\ell'} \delta_{mm'}$ such that $Y_{00} = 1$ and $\phi_{00} = 1$ for the minimal solutions in (3.14), yields the effective action for the $6d$ fluctuations

$$\delta^2 S_{\text{IIB}} = \sum_{\ell, m} C_\ell \int d^6x \sqrt{-g_{AdS_6}} (h_{\ell, m}^{[tt]})^{\mu\nu} [\square_{AdS_6} - (3\ell(3\ell + 5) - 2)] (h_{\ell, m}^{[tt]})_{\mu\nu}. \quad (4.10)$$

The constants C_ℓ evaluate to

$$C_\ell = \frac{1}{\kappa_{10}^2} \text{Vol}_{S^2} \int d^2w 4f_6^4 f_2^2 \rho^2 (\phi_{\ell m})^2 = \frac{8}{3\kappa_{10}^2} \text{Vol}_{S^2} \int d^2w \kappa^2 \mathcal{G} (\phi_{\ell m})^2, \quad (4.11)$$

with $dw^2 = \frac{i}{2} dw \wedge d\bar{w}$ and where (3.8) has been used to obtain the second equality. The effective 6d gravitational coupling is related to C_0 for the minimal $\ell = 0$ solution with $\phi_{00} = 1$, corresponding to the massless graviton, by $C_0 = 1/\kappa_6^2$.

As discussed recently in section 4 of [41], C_T is thus directly related to C_0 for the minimal solution as defined above. The result (4.11) also shows that this coefficient C_0 , up to a universal numerical factor that is independent of the choice of solution, is precisely the finite part of the entanglement entropy for a spherical region, as computed in section IV of [21]. As verified explicitly in [21], the finite part of this entanglement entropy is given by the S^5 partition function of the dual SCFTs. This establishes a direct relation of C_T to the S^5 partition function, and shows that the two notions of ‘‘central charge’’ are equivalent for the theories dual to the supergravity solutions discussed here.

The C_ℓ are in particular finite for the physically regular solutions constructed in [8, 9, 19]. This follows from the discussion in [21], where it was shown that the integral in (4.11) for $\ell = 0$ and $\phi_{\ell m} = 1$ is finite, despite the presence of poles in the differentials $\partial_w \mathcal{A}_\pm$ on the boundary $\partial\Sigma$. Since the $\phi_{\ell m}$ are well behaved for the minimal solutions with $\ell \geq 0$ and the non-minimal solutions with $\ell > 0$, as discussed in section 3.2 and 3.3, respectively, this immediately implies finiteness of the integral in (4.11) for all these solutions. This extends to the non-minimal solution with $\ell = 0$ as well, since the logarithmic singularities near the poles in ϕ_{00} of (3.23) are mild enough for the complete integrand in (4.11) to be integrable.

5 Discussion

We have identified two special classes of (massive) spin-2 excitations around the warped $AdS_6 \times S^2 \times \Sigma$ solutions to type IIB supergravity constructed in [9]. Determining the full spectrum of spin-2 fluctuations is a non-trivial problem, and the full results are expected to depend crucially on the details of the specific solution at hand. Remarkably, the fluctuations identified here are universally present, for all solutions of this form. The fluctuations are expressed directly in terms of the basic building blocks of the background solution and satisfy natural regularity conditions and boundary conditions on $\partial\Sigma$. In the dual $5d$ SCFTs they correspond to two sets of universally present spin-2 operators, which are Q^4 descendants in B_2 and A_4 multiplets in the notation of [27].

These results are reminiscent of the recently obtained results on spin-2 fluctuations [36] around the massive type IIA solution of [4], but differ in crucial details. The scaling dimensions (4.1) and (4.4) superficially agree with the scaling dimensions found in [36]. However, as discussed in section 3.2, we only find multiplets with even integer R -charge, whereas in [36] even and odd integer R -charges appear. This can be explained by the fact that the R -symmetry is realized differently: in the type IIB solution the R -symmetry is realized directly as the isometries of S^2 , while it appears as part of the isometries of a half S^4 in the massive IIA solutions. A second point concerns regularity of the fluctuations dual to A_4 multiplets, of which we find two sets. While those were found to be singular, albeit normalizable, in [36], the corresponding fluctuations for the type IIB solutions identified here are regular for $\ell > 0$.

The massive type IIA solution can formally be T-dualized along an S^1 inside the half S^4 , to obtain a solution in type IIB. The resulting singular solution is contained as a special case in the solutions discussed here, and we have shown in appendix A how the PDE for the spin-2 fluctuations reduces to an ODE in that case. The implementation of regularity conditions, however, is unclear for this background and we did not attempt a general analysis. It may be interesting in that context to also study the non-Abelian T-dual.

Finally, we have studied generalizations of the notion of central charge to $5d$ SCFTs. Two objects which have been studied in that context are the coefficient C_T appearing in the two-point function of the energy-momentum tensor and the five-sphere partition function. Drawing on results obtained earlier, we showed that the two notions of central charge are equivalent for the $5d$ SCFTs described by classical supergravity on the solutions of [9, 19].

We close the paper with some open questions and directions for future work:

As discussed above, the fluctuations identified here correspond to Q^4 descendants in short multiplets with scalar primaries. It would be interesting to identify the other members of these multiplets as well. While the full KK reduction is a technically very challenging task, the fluctuations corresponding to the remaining members of these multiplets should be related by supersymmetry transformations to the ones identified here. In particular, they should also be universal, and likely have expressions in terms of the basic building blocks of the background solutions. This should make them accessible and provide potentially interesting insights on the structure of the full KK spectrum.

An immediate question from the perspective of the dual SCFTs is whether and how the operators identified here can be realized in terms of Lagrangian fields in gauge theory deformations. The supergravity solutions of [9, 19] are naturally associated to (p, q) 5-brane webs, many of which have gauge theory deformations to quiver theories. It would be interesting to understand what the universality of the operators we found implies in that context and how they are realized.

Acknowledgments

We are grateful to Constantin Bachas, Eric D’Hoker, Martin Fluder, Achilleas Passias and Paul Richmond for useful conversations. The work of MG and CFU is supported in part by the National Science Foundation under grant PHY-16-19926. OV is supported by NSF grant PHY-1720364 and, partially, by grant FPA2015-65480-P (MINECO/FEDER UE) from the Spanish Government. MG is grateful to the International Solvay Institutes, ULB, Brussels for hospitality while this paper was finalized.

A T-dual of massive IIA solution

We briefly discuss the T-dual of the massive type IIA solution, as given in appendix A of [12]. A choice of \mathcal{A}_\pm realizing this solution was given in section 5.6 of [7], as

$$\mathcal{A}_\pm = \frac{1}{2}aw^2 \mp bw. \tag{A.1}$$

The parameters a, b were given for $L = 1$, where L is the curvature radius of AdS_6 , in [7]. For generic L they are

$$a = \frac{27}{16}L^4m^{1/3}, \quad b = \frac{9L^2}{8m^{1/3}}, \tag{A.2}$$

with m encoding the Romans mass. The relevant quantities for the discussion of spin-2 fluctuations are κ^2 and \mathcal{G} , given by

$$\kappa^2 = 2ab(1 - Z)^{1/3}, \quad \mathcal{G} = \frac{ab}{3}Z, \quad Z = 1 - (w + \bar{w})^3. \tag{A.3}$$

With the field identifications and real coordinates discussed in [7], defined by $\cos\theta = w + \bar{w}$ and $\phi_3 = \frac{ia}{2bm}(w - \bar{w})$, this produces the solution as given in appendix A of [12], including the factors of L .

A.1 Metric perturbations

The equation determining the spin-2 fluctuations is (3.12), with κ^2 and \mathcal{G} given in (A.3). The constants a, b drop out, and the equation becomes

$$Z^{-2\ell-1}\partial_a \left(Z^{2\ell+2}\partial^a \chi_{\ell m} \right) + (M^2 - 3\ell(3\ell + 5))(1 - Z)^{1/3} \chi_{\ell m} = 0. \tag{A.4}$$

The geometry is invariant under shifts in $w - \bar{w}$, and the general solution can therefore be expanded as

$$\chi_{\ell m}(w, \bar{w}) = f(w + \bar{w})e^{ik(w-\bar{w})}. \quad (\text{A.5})$$

Eq. (A.4) becomes, with a real coordinate $x = w + \bar{w}$ such that $Z = 1 - x^3$,

$$Z^{-2\ell-1}\partial_x \left(Z^{2\ell+2}\partial_x f \right) - k^2 Z f + (M^2 - 3\ell(3\ell + 5)) x f = 0. \quad (\text{A.6})$$

Setting $z = x^3$ turns this equation into

$$z(1-z)\partial_z^2 f - \frac{2}{3}((3\ell+4)z-1)\partial_z f + \frac{1}{9}(M^2 - 3\ell(3\ell+5))f - k^2 f \frac{1-z}{z^{1/3}} = 0. \quad (\text{A.7})$$

For $k = 0$ this is a hypergeometric differential equation in standard form. The minimal solutions discussed in section 3.2 correspond to constant f with $k = 0$. The real part of the non-minimal solutions for $\chi_{\ell m}$ in section 3.3 corresponds to $f = z^{1/3}$ with $k = 0$, while the imaginary part has non-trivial dependence on $w - \bar{w}$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] N. Seiberg, *Five-dimensional SUSY field theories, nontrivial fixed points and string dynamics*, *Phys. Lett. B* **388** (1996) 753 [[hep-th/9608111](#)] [[INSPIRE](#)].
- [2] D.R. Morrison and N. Seiberg, *Extremal transitions and five-dimensional supersymmetric field theories*, *Nucl. Phys. B* **483** (1997) 229 [[hep-th/9609070](#)] [[INSPIRE](#)].
- [3] K.A. Intriligator, D.R. Morrison and N. Seiberg, *Five-dimensional supersymmetric gauge theories and degenerations of Calabi-Yau spaces*, *Nucl. Phys. B* **497** (1997) 56 [[hep-th/9702198](#)] [[INSPIRE](#)].
- [4] A. Brandhuber and Y. Oz, *The D4–D8 brane system and five-dimensional fixed points*, *Phys. Lett. B* **460** (1999) 307 [[hep-th/9905148](#)] [[INSPIRE](#)].
- [5] O. Bergman and D. Rodríguez-Gómez, *5d quivers and their AdS₆ duals*, *JHEP* **07** (2012) 171 [[arXiv:1206.3503](#)] [[INSPIRE](#)].
- [6] A. Passias, *A note on supersymmetric AdS₆ solutions of massive type IIA supergravity*, *JHEP* **01** (2013) 113 [[arXiv:1209.3267](#)] [[INSPIRE](#)].
- [7] E. D’Hoker, M. Gutperle, A. Karch and C.F. Uhlemann, *Warped AdS₆ × S² in Type IIB supergravity I: Local solutions*, *JHEP* **08** (2016) 046 [[arXiv:1606.01254](#)] [[INSPIRE](#)].
- [8] E. D’Hoker, M. Gutperle and C.F. Uhlemann, *Holographic duals for five-dimensional superconformal quantum field theories*, *Phys. Rev. Lett.* **118** (2017) 101601 [[arXiv:1611.09411](#)] [[INSPIRE](#)].
- [9] E. D’Hoker, M. Gutperle and C.F. Uhlemann, *Warped AdS₆ × S² in Type IIB supergravity II: Global solutions and five-brane webs*, *JHEP* **05** (2017) 131 [[arXiv:1703.08186](#)] [[INSPIRE](#)].

- [10] D. Corbino, E. D'Hoker and C.F. Uhlemann, *AdS₂ × S⁶ versus AdS₆ × S² in Type IIB supergravity*, *JHEP* **03** (2018) 120 [[arXiv:1712.04463](#)] [[INSPIRE](#)].
- [11] Y. Lozano, E.Ó. Colgáin, D. Rodríguez-Gómez and K. Sfetsos, *Supersymmetric AdS₆ via T Duality*, *Phys. Rev. Lett.* **110** (2013) 231601 [[arXiv:1212.1043](#)] [[INSPIRE](#)].
- [12] Y. Lozano, E.Ó. Colgáin and D. Rodríguez-Gómez, *Hints of 5d Fixed Point Theories from Non-Abelian T-duality*, *JHEP* **05** (2014) 009 [[arXiv:1311.4842](#)] [[INSPIRE](#)].
- [13] F. Apruzzi, M. Fazzi, A. Passias, D. Rosa and A. Tomasiello, *AdS₆ solutions of type-II supergravity*, *JHEP* **11** (2014) 099 [Erratum *JHEP* **05** (2015) 012] [[arXiv:1406.0852](#)] [[INSPIRE](#)].
- [14] Ö. Kelekci, Y. Lozano, N.T. Macpherson and E.Ó. Colgáin, *Supersymmetry and non-Abelian T-duality in type-II supergravity*, *Class. Quant. Grav.* **32** (2015) 035014 [[arXiv:1409.7406](#)] [[INSPIRE](#)].
- [15] H. Kim, N. Kim and M. Suh, *Supersymmetric AdS₆ Solutions of Type IIB Supergravity*, *Eur. Phys. J. C* **75** (2015) 484 [[arXiv:1506.05480](#)] [[INSPIRE](#)].
- [16] H. Kim and N. Kim, *Comments on the symmetry of AdS₆ solutions in string/M-theory and Killing spinor equations*, *Phys. Lett. B* **760** (2016) 780 [[arXiv:1604.07987](#)] [[INSPIRE](#)].
- [17] O. Aharony, A. Hanany and B. Kol, *Webs of (p, q) five-branes, five-dimensional field theories and grid diagrams*, *JHEP* **01** (1998) 002 [[hep-th/9710116](#)] [[INSPIRE](#)].
- [18] O. Aharony and A. Hanany, *Branes, superpotentials and superconformal fixed points*, *Nucl. Phys. B* **504** (1997) 239 [[hep-th/9704170](#)] [[INSPIRE](#)].
- [19] E. D'Hoker, M. Gutperle and C.F. Uhlemann, *Warped AdS₆ × S² in Type IIB supergravity III: Global solutions with seven-branes*, *JHEP* **11** (2017) 200 [[arXiv:1706.00433](#)] [[INSPIRE](#)].
- [20] M. Gutperle, A. Trivella and C.F. Uhlemann, *Type IIB 7-branes in warped AdS₆: partition functions, brane webs and probe limit*, *JHEP* **04** (2018) 135 [[arXiv:1802.07274](#)] [[INSPIRE](#)].
- [21] M. Gutperle, C. Marasinou, A. Trivella and C.F. Uhlemann, *Entanglement entropy vs. free energy in IIB supergravity duals for 5d SCFTs*, *JHEP* **09** (2017) 125 [[arXiv:1705.01561](#)] [[INSPIRE](#)].
- [22] J. Kaidi, *(p, q)-strings probing five-brane webs*, *JHEP* **10** (2017) 087 [[arXiv:1708.03404](#)] [[INSPIRE](#)].
- [23] W. Nahm, *Supersymmetries and their Representations*, *Nucl. Phys. B* **135** (1978) 149 [[INSPIRE](#)].
- [24] V.G. Kac, *Lie Superalgebras*, *Adv. Math.* **26** (1977) 8 [[INSPIRE](#)].
- [25] S. Shnider, *The Superconformal Algebra In Higher Dimensions*, *Lett. Math. Phys.* **16** (1988) 377 [[INSPIRE](#)].
- [26] M. Buican, J. Hayling and C. Papageorgakis, *Aspects of Superconformal Multiplets in D > 4*, *JHEP* **11** (2016) 091 [[arXiv:1606.00810](#)] [[INSPIRE](#)].
- [27] C. Cordova, T.T. Dumitrescu and K. Intriligator, *Multiplets of Superconformal Symmetry in Diverse Dimensions*, [[arXiv:1612.00809](#)] [[INSPIRE](#)].
- [28] H.J. Kim, L.J. Romans and P. van Nieuwenhuizen, *The Mass Spectrum of Chiral N = 2 D = 10 Supergravity on S⁵*, *Phys. Rev. D* **32** (1985) 389 [[INSPIRE](#)].

- [29] M. Günaydin and N. Marcus, *The Spectrum of the S^5 Compactification of the Chiral $N = 2$, $D = 10$ Supergravity and the Unitary Supermultiplets of $U(2, 2/4)$* , *Class. Quant. Grav.* **2** (1985) L11 [[INSPIRE](#)].
- [30] C. Bachas and J. Estes, *Spin-2 spectrum of defect theories*, *JHEP* **06** (2011) 005 [[arXiv:1103.2800](#)] [[INSPIRE](#)].
- [31] I.R. Klebanov, S.S. Pufu and F.D. Rocha, *The Squashed, Stretched and Warped Gets Perturbed*, *JHEP* **06** (2009) 019 [[arXiv:0904.1009](#)] [[INSPIRE](#)].
- [32] J.-M. Richard, R. Terrisse and D. Tsimpis, *On the spin-2 Kaluza-Klein spectrum of $AdS_4 \times S^2(\mathcal{B}_4)$* , *JHEP* **12** (2014) 144 [[arXiv:1410.4669](#)] [[INSPIRE](#)].
- [33] A. Passias and A. Tomasiello, *Spin-2 spectrum of six-dimensional field theories*, *JHEP* **12** (2016) 050 [[arXiv:1604.04286](#)] [[INSPIRE](#)].
- [34] J. Schmude and O. Vasilakis, *Superconformal Symmetry in the Kaluza-Klein Spectrum of Warped AdS_3* , *JHEP* **10** (2016) 096 [[arXiv:1605.00636](#)] [[INSPIRE](#)].
- [35] Y. Pang, J. Rong and O. Varela, *Spectrum universality properties of holographic Chern-Simons theories*, *JHEP* **01** (2018) 061 [[arXiv:1711.07781](#)] [[INSPIRE](#)].
- [36] A. Passias and P. Richmond, *Perturbing $AdS_6 \times_w S^4$: linearised equations and spin-2 spectrum*, [arXiv:1804.09728](#) [[INSPIRE](#)].
- [37] J.P. Gauntlett and O. Varela, *Consistent Kaluza-Klein reductions for general supersymmetric AdS solutions*, *Phys. Rev. D* **76** (2007) 126007 [[arXiv:0707.2315](#)] [[INSPIRE](#)].
- [38] C.-M. Chang, M. Fluder, Y.-H. Lin and Y. Wang, *Spheres, Charges, Instantons and Bootstrap: A Five-Dimensional Odyssey*, *JHEP* **03** (2018) 123 [[arXiv:1710.08418](#)] [[INSPIRE](#)].
- [39] J. Polchinski, *String theory. Volume 2: Superstring theory and beyond*, Cambridge Monographs on Mathematical Physics, [Cambridge University Press](#), Cambridge U.K. (2007) [ISBN: 9780511252280, 9780521633048, 9780521672283] [[INSPIRE](#)].
- [40] F. Benini, S. Benvenuti and Y. Tachikawa, *Webs of five-branes and $N = 2$ superconformal field theories*, *JHEP* **09** (2009) 052 [[arXiv:0906.0359](#)] [[INSPIRE](#)].
- [41] C.-M. Chang, M. Fluder, Y.-H. Lin and Y. Wang, *Romans Supergravity from Five-Dimensional Holograms*, *JHEP* **05** (2018) 039 [[arXiv:1712.10313](#)] [[INSPIRE](#)].