

Corrigendum

Decomposition of time-dependent fluorescence signals reveals codon-specific kinetics of protein synthesis

Nadin Haase¹, Wolf Holtkamp², Reinhard Lipowsky¹, Marina Rodnina² and Sophia Rudolf^{1,*}

¹Theory and Bio-Systems, Max Planck Institute of Colloids and Interfaces, Am Muehlenberg 1, 14476 Potsdam, Germany and ²Physical Biochemistry, Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Goettingen, Germany

Nucleic Acids Research, 2018, gky740, <https://doi.org/10.1093/nar/gky740>

The authors wish to correct an error in their article. In Figure 3, the data set for the ‘no EF-G’ experiment is plotted only for 1s, instead of for 10s as indicated. A new Figure 3 is provided below. The article has been updated online and in print. This correction does not influence the results and overall conclusions of the article.

*To whom correspondence should be addressed. Tel: +49 331 567 9638; Fax: +49 331 567 9602; Email: sophia.rudorf@mpikg.mpg.de




Figure 3. Measured fluorescence signatures of the in-vitro synthesis of BOF-fMetPhen peptides (colored lines) and best fit in terms of least squares of the theoretical translation model (black lines). The label 'no EF-G' indicates translation of poly(U) mRNA in the absence of EF-G.