

CLOSED STRINGS AS SINGLE-VALUED OPEN STRINGS: A GENUS-ZERO DERIVATION

OLIVER SCHLOTTERER, OLIVER SCHNETZ

ABSTRACT. Based on general mathematical assumptions we give an independent, elementary proof of a theorem by Francis Brown and Clément Dupont in [1] which states that tree-level amplitudes of closed and open strings are related through the single-valued map ‘sv’. This relation can be traced back to the underlying moduli-space integrals over punctured Riemann surfaces of genus zero. The sphere integrals J in closed-string amplitudes and the disk integrals Z in open-string amplitudes are shown to obey $J = \text{sv } Z$.

1. INTRODUCTION

The study of scattering amplitudes grew into a fertile and rapidly developing research area at the interface of particle physics, mathematics and string theory. A wealth of modern mathematical concepts including periods, motives and elliptic functions became a common theme in scattering amplitudes of quantum field theory and string theory: Field-theory amplitudes encounter various flavors of polylogarithms via Feynman integrals, and string amplitudes are formulated in terms of moduli-space integrals for punctured Riemann surfaces. In contrast to field theory, the infinite number of vibration modes in string spectra introduces transcendental numbers already into the tree level of string perturbation theory.

More specifically, the low-energy expansions of tree-level amplitudes of both open and closed strings involve multiple zeta values (MZVs),

$$(1) \quad \zeta(k_1, k_2, \dots, k_r) = \sum_{0 < l_1 < l_2 < \dots < l_r}^{\infty} l_1^{-k_1} l_2^{-k_2} \dots l_r^{-k_r}, \quad k_1, k_2, \dots, k_r \in \mathbb{N}, \quad k_r \geq 2,$$

characterized by depth r and weight $k_1 + k_2 + \dots + k_r$. MZVs are the periods of the moduli space $M_{0,n}$ of n -punctured genus-zero surfaces [2]: For open strings, MZVs arise from iterated integrals over the boundary of a disk, and closed-string tree amplitudes are obtained from complex integration over punctures on a sphere. From the work of Kawai, Lewellen and Tye (KLT) in 1986 [3], the sphere integrals for closed strings are known to factorize into bilinears in disk integrals for open strings. However, the approach of KLT does not manifest whether the ‘squaring procedure’ for disk integrals induces any cancellations for certain classes of MZVs. From the observations of [4], only the so-called single-valued subclass of MZVs (see [5]) seems to persist in the final results for the sphere integrals in closed-string tree amplitudes. The purpose of this work is to give a general proof for these conjectural selection rules. In fact, as will be detailed below, the closed-string amplitudes are tied to open-string amplitudes by the ‘single-valued map’.

While four-point tree-level scattering of open strings gives rise to all the Riemann zeta values $\zeta(m)$, $2 \leq m \in \mathbb{N}$ in the low-energy expansion, the analogous closed-string four-point function only involves odd zeta values $\zeta(2k+1)$, $k \in \mathbb{N}$. Here, the cancellations of integer powers of π^2 can be tracked by the closed-form representation of the four-point amplitudes in terms of gamma functions of the kinematic data. In open-string amplitudes with $n \geq 5$ external legs, in turn, the MZVs in the low-energy expansions include higher-depth instances and follow a more elaborate structure that can be understood in terms of motivic MZVs [4] and the Drinfeld

associator [6, 7]¹. And it took until 2012 that an all-order conjecture² for the selection rules on the MZVs in closed-string n -point amplitudes could be made [4], based on an order-by-order inspection of the output of the KLT relations.

According to the observations in [4], closed-string low-energy expansions are conjectured to follow from the *single-valued* map [5] of the MZVs in the disk integrals of open-string amplitudes [16, 17]. A defining property of the resulting *single-valued MZVs* is their descent from single-valued multiple polylogarithms at unit argument. The procedure of Francis Brown [18] to eliminate the monodromies from harmonic polylogarithms induces a map on MZVs which is referred to as the single-valued map sv [19, 5]. From the conjectures in [4, 16, 17], the MZVs in the low-energy expansion of n -point sphere integrals J can be obtained from specific disk integrals Z via the sv map. As will be detailed below, the relation [17] (see (33), an independent proof by F. Brown and C. Dupont was recently announced in [1])

$$(2) \quad J = sv Z$$

associates certain anti-meromorphic functions of the punctures on the sphere with cyclic orderings of the punctures on the disk boundary, following a Betti-deRham duality.

In this work, we will deliver an inductive proof under general mathematical assumptions that sphere integrals are single-valued versions of disk integrals and, equivalently, that closed-string tree-level amplitudes are single-valued open-string amplitudes. The driving force for the proof is the notion of single-valued integration [19] along with its properties that originate from motivic algebraic geometry [5, 20]. The Betti-deRham duality between anti-meromorphic integrands on the sphere and integration cycles on the disk boundary will arise naturally from the Stokes theorem.

The sv relations between disk and sphere integrals can be applied to closed strings in the supersymmetric, heterotic and bosonic theories [17] and triggered several directions of follow-up research. For instance, single-valued open-string amplitudes govern amplitude relations mixing gauge and gravitational states of the heterotic string [21] as well as the recent double-copy description of bosonic and heterotic strings [22]. Moreover, the appearance of single-valued MZVs in the sigma-model approach to effective gauge interactions of type-I and heterotic strings has been studied in [23]. The proof in this work will place these results on firm grounds without the need to rely on a conjectural status for the key relations between sphere integrals and single-valued disk integrals.

The authors would like to emphasize that the proof in this article is *not* a proof in a full mathematical sense. Mostly, one has to keep in mind that the single-valued map is only defined to exist in a ‘motivic’ framework. In the motivic setup integrals are lifted to objects in algebraic geometry thereby bypassing general transcendental problems. Because the singular divisor in the disk and sphere integrals are not normal crossing, it is a non-trivial step to set up a motivic theory for these objects. Alternatively, one can assume standard transcendental conjectures for the related integrals.

Moreover, we use three natural properties of the single-valued map in section 3.1. These properties are thoroughly tested. Properties (i) and (ii) are proved or mostly proved in the stated literature. Property (iii) is proved in the text using a standard property of the ‘ f -alphabet’ which may not be fully proved in the mathematical literature (the f -alphabet has not yet drawn much attention in mathematics).

Finally, we use the existence of a subtraction scheme whose existence we do not fully prove. Subtractions schemes are extensively studied in the much more complicated case of quantum field theory. So, we considered it more beneficial to provide the reader with explicit examples

¹State-of-the-art methods to compute the low-energy expansion of n -point disk integrals include matrix representations of the Drinfeld associator [7] and recursions for off-shell versions of the disk integrals [8] (building upon the approach via polylogarithm manipulations in [9]). For certain multiplicities n , explicit results are available for download via [10], and one can also use the connection between disk integrals and hypergeometric functions to extract low-energy expansions, see e.g. [11, 12, 13, 14] and references therein.

²Also see [15] for earlier work on MZVs at weight ≤ 8 in closed-string five- and six-point functions.

in appendix A rather than going through all technical details at full length. The proof of (2) by F. Brown and C. Dupont also establishes a rigorous approach to subtraction in the context of genus-zero string theory [1].

In spite of these restrictions we informally use the word ‘proof’ in this article.

2. REVIEWING THE BASES AND RELATIONS OF DISK AND SPHERE INTEGRALS

In this section, we review the classes of disk and sphere integrals that are related through the sv map. These moduli-space integrals encode the low-energy regime of string tree-level amplitudes through their series expansion in the dimensionless Mandelstam invariants

$$(3) \quad s_{ij} := 2\alpha' k_i \cdot k_j = s_{ji}, \quad s_{ij} \in \mathbb{R},$$

where α' denotes the inverse string tension. The external momenta k_i are Lorentz vectors referring to massless external states $i = 1, 2, \dots, n$ of an n -point amplitude subject to $k_i^2 = 0$ and momentum conservation $\sum_{i=1}^n k_i = 0$. These kinematic constraints imply

$$(4) \quad s_{i,i} = 0, \quad \sum_{i=1}^n s_{ij} = 0 \quad \forall j = 1, 2, \dots, n,$$

so that only $\frac{n}{2}(n-3)$ Mandelstam invariants are independent.

2.1. Four-point integrals: an inviting example. The simplest appearance of MZVs in string perturbation theory occurs in the four-point tree amplitude of open strings. After peeling off suitable kinematic factors, the amplitude boils down to the disk integral

$$(5) \quad Z_{4\text{pt}} := \int_0^1 \frac{dz}{z} z^{s_{12}} (1-z)^{s_{23}} = \frac{\Gamma(s_{12})\Gamma(1+s_{23})}{\Gamma(1+s_{12}+s_{23})}$$

and its permutations w.r.t. the external momenta. The α' -expansion of the integral $Z_{4\text{pt}}$ – i.e. the simultaneous series expansion in the dimensionless s_{ij} variables (3) – follows from the Γ -function identity $\log \Gamma(1+x) = -\gamma x + \sum_{k=2}^{\infty} \frac{\zeta(k)}{k} (-x)^k$,

$$(6) \quad \begin{aligned} Z_{4\text{pt}} &= \frac{1}{s_{12}} \exp\left(\sum_{k=2}^{\infty} \frac{\zeta(k)}{k} (-1)^k [s_{12}^k + s_{23}^k - (s_{12} + s_{23})^k]\right) \\ &= \frac{1}{s_{12}} - \zeta(2)s_{23} + \zeta(3)s_{23}(s_{12} + s_{23}) + O(\alpha'^3), \end{aligned}$$

and involves all Riemann zeta values (while the Euler Mascheroni constant γ cancels).

The simplest appearance of MZVs in a closed-string setup is the following complex integral in the four-point tree amplitude

$$(7) \quad J_{4\text{pt}} := \frac{1}{\pi} \int_{\mathbb{C}} \frac{d^2z}{z\bar{z}(1-\bar{z})} |z|^{2s_{12}} |1-z|^{2s_{23}} = \frac{\Gamma(s_{12})\Gamma(1+s_{23})\Gamma(1+s_{13})}{\Gamma(1-s_{12})\Gamma(1-s_{23})\Gamma(1-s_{13})},$$

where \bar{z} is the complex conjugate of $z = x + iy$ and $d^2z := dx dy$. The α' -expansion takes a particularly symmetric form in terms of $s_{13} = -s_{12} - s_{23}$, see (3),

$$(8) \quad \begin{aligned} J_{4\text{pt}} &= \frac{1}{s_{12}} \exp\left(-2 \sum_{k=1}^{\infty} \frac{\zeta(2k+1)}{2k+1} [s_{12}^{2k+1} + s_{23}^{2k+1} + s_{13}^{2k+1}]\right) \\ &= \frac{1}{s_{12}} + 2\zeta(3)s_{23}(s_{12} + s_{23}) + O(\alpha'^4), \end{aligned}$$

and the first line of (8) manifests the cancellation of even Riemann-zeta values.

2.2. The integrals for n points. The above four-point integrals fall into the following general classes of n -point disk integrals $Z(\tau|\rho)$ and sphere integrals $J(\tau|\rho)$,

(9)

$$Z(\tau|\rho) := \int_{-\infty \leq z_{\tau(1)} \leq z_{\tau(2)} \leq \dots \leq z_{\tau(n)} \leq \infty} \frac{dz_1 dz_2 \dots dz_n}{\text{vol SL}_2(\mathbb{R})} \frac{(-1)^{n-3} \prod_{1 \leq i < j \leq n} |z_{i,j}|^{s_{ij}}}{z_{\rho(1),\rho(2)} z_{\rho(2),\rho(3)} \dots z_{\rho(n-1),\rho(n)} z_{\rho(n),\rho(1)}},$$

(10)

$$J(\tau|\rho) := \int_{\mathbb{C}^n} \frac{d^2 z_1 d^2 z_2 \dots d^2 z_n}{\pi^{n-3} \text{vol SL}_2(\mathbb{C})} \frac{\prod_{1 \leq i < j \leq n} |z_{i,j}|^{2s_{ij}}}{(z_{\rho(1),\rho(2)} z_{\rho(2),\rho(3)} \dots z_{\rho(n),\rho(1)}) (\bar{z}_{\tau(1),\tau(2)} \bar{z}_{\tau(2),\tau(3)} \dots \bar{z}_{\tau(n),\tau(1)})},$$

where $z_{i,j} := z_i - z_j$. Both types of integrals are indexed by two permutations $\rho, \tau \in S_n$ of the legs $\{1, 2, \dots, n\}$. The absolute value in the integrand $\prod_{1 \leq i < j \leq n} |z_{i,j}|^{s_{ij}}$ of (9) ensures that only positive numbers are raised to the power of s_{ij} , regardless of the integration domain characterized by $z_{\tau(i)} < z_{\tau(i+1)}$.

The inverse factor of $\text{vol SL}_2(\mathbb{R})$ in the disk integrals (9) is implemented by dropping three integrations over any z_i, z_j, z_k (with $i, j, k \in \{1, 2, \dots, n\}$), inserting $|z_{i,j} z_{i,k} z_{j,k}|$ and fixing $(z_i, z_j, z_k) \rightarrow (0, 1, \infty)$. Its analogue $(\text{vol SL}_2(\mathbb{C}))^{-1}$ in the sphere integral (10) instructs to insert $|z_{i,j} z_{i,k} z_{j,k}|^2$. The limit $z_k \rightarrow \infty$ is non-singular by the Mandelstam identity (4) and the choice of cyclic ‘Parke–Taylor’ denominators in (9) and (10).

Note that the four-point integrals (5) and (7) can be recovered from the general definition via

$$(11) \quad Z_{4\text{pt}} = -Z(1, 2, 3, 4|1, 2, 4, 3), \quad J_{4\text{pt}} = -J(1, 2, 3, 4|1, 2, 4, 3)$$

after fixing $(z_1, z_3, z_4) \rightarrow (0, 1, \infty)$ and identifying $z_2 \rightarrow z$.

By the results of refs. [24, 9], the n -point tree-level amplitudes of open and closed superstrings are expressible in terms of the integrals (9) and (10), also see [25, 22] for analogous statements on bosonic and heterotic strings.

2.3. Relations of disk and sphere integrals. One can infer from the right-hand sides of (9) and (10) that the disk and sphere integrals $Z(\tau|\rho)$ and $J(\tau|\rho)$ only depend on the cyclic equivalence class of the permutations τ, ρ . The cyclic denominators manifest that

$$(12) \quad Z(\tau|1, 2, 3, \dots, n) = Z(\tau|2, 3, \dots, n, 1), \quad J(\tau|1, 2, 3, \dots, n) = J(\tau|2, 3, \dots, n, 1) \quad \forall \tau \in S_n,$$

and the same is true for the first entry of the sphere integrals [by reality $J(\tau|\rho) = J(\rho|\tau)$]. Also, the integration domain of the disk integrals (9) is cyclically invariant

$$(13) \quad Z(1, 2, 3, \dots, n|\rho) = Z(2, 3, \dots, n, 1|\rho) \quad \forall \rho \in S_n.$$

Still, the number $(n-1)!$ of cyclically inequivalent permutations in S_n overcounts the number of inequivalent disk and sphere integrals: Different choices of the cyclic denominators are related via integration-by-parts relations which lead to a basis of $(n-3)!$ inequivalent permutations of $(z_{1,2} z_{2,3} \dots z_{n,1})^{-1}$. For disk integrals, dropping total derivatives w.r.t. the punctures yields [9]

$$(14) \quad \sum_{j=2}^{n-1} k_1 \cdot (k_2 + k_3 + \dots + k_j) Z(\tau|2, 3, \dots, j, 1, j+1, \dots, n-1, n) = 0 \quad \forall \tau \in S_n,$$

and the same relations hold for both entries of the sphere integrals. Since the first entry of the disk integrals (9) refers to an integration cycle $-\infty \leq z_{\tau(1)} \leq z_{\tau(2)} \leq \dots \leq z_{\tau(n)} \leq \infty$ rather than a choice of integrand, i.e. $Z(\tau|\rho) \neq Z(\rho|\tau)$, monodromy properties of the Koba–Nielsen factor $\prod_{1 \leq i < j \leq n} |z_{i,j}|^{s_{ij}}$ yield [26, 27]

$$(15) \quad \sum_{j=2}^{n-1} \sin [2\pi \alpha' k_1 \cdot (k_2 + k_3 + \dots + k_j)] Z(2, 3, \dots, j, 1, j+1, \dots, n-1, n|\rho) = 0 \quad \forall \rho \in S_n.$$

The combinatorics of these monodromy relations follow the structure of (14) except for the promotion of the coefficients $k_1 \cdot (k_2 + k_3 + \dots + k_j)$ to a trigonometric function. Hence, permutations of (15) leave $(n-3)!$ independent integration cycles [26, 27].

By combining permutations of (14) and (15), the moduli-space integrals $Z(\tau|\rho)$ and $J(\tau|\rho)$ can be expressed in a basis of $(n-3)! \times (n-3)!$ elements. For both entries, one can fix legs $n-1, n, 1$ in adjacent positions and take $\rho = 1, \beta, n-1, n$ with permutations $\beta \in S_{n-3}$ of $\{2, 3, \dots, n-2\}$ as a convenient basis choice. These relations can be understood in the framework of intersection theory, where $(n-3)!$ arises as the dimension of twisted homologies and cohomologies [28, 29].

2.4. Kawai–Lewellen–Tye relations. Using the representations of the four-point integrals (5) and (7) in terms of Γ functions, one can observe via $\sin(\pi x) = \frac{\pi}{\Gamma(1-x)\Gamma(x)}$ that

$$(16) \quad J(1, 2, 3, 4|1, 2, 4, 3) = -\frac{1}{\pi} Z(1, 2, 3, 4|1, 2, 4, 3) \sin(\pi s_{12}) Z(1, 2, 4, 3|1, 2, 3, 4).$$

This is the simplest instance of the KLT relations [3] between sphere integrals and bilinears in disk integrals which can be derived by suitable deformations of the complex integration contours. Their generalization to n -points do not depend on the rational factors besides $\prod_{1 \leq i < j \leq n} |z_{i,j}|^{2s_{ij}}$ in the integrand of $J(\tau|\rho)$ and may be described in terms of a $(n-3)! \times (n-3)!$ KLT matrix $S_{\alpha'}(\sigma|\beta)_1$ [3, 30, 31]

$$(17) \quad J(\tau|\rho) = \sum_{\sigma, \beta \in S_{n-3}} Z(1, \sigma, n, n-1|\tau) S_{\alpha'}(\sigma|\beta)_1 Z(1, \beta, n-1, n|\rho).$$

The KLT matrix $S_{\alpha'}(\sigma|\beta)_1$ is indexed by permutations $\sigma, \beta \in S_{n-3}$ of $\{2, 3, \dots, n-2\}$ and admits a recursive definition [31, 32]

$$(18) \quad \begin{aligned} S_{\alpha'}(2|2)_1 &= -\frac{1}{\pi} \sin(\pi s_{12}) = -\frac{1}{\pi} \sin(2\pi\alpha' k_1 \cdot k_2) \\ S_{\alpha'}(A, j|B, j, C)_1 &= -\frac{1}{\pi} \sin(2\pi\alpha' k_j \cdot (k_1 + k_B)) S_{\alpha'}(A|B, C)_1. \end{aligned}$$

Here, we are employing the notation $A = a_1 a_2 \dots a_p$ and $B = b_1 b_2 \dots b_q$ for words of length $p, q \geq 0$ composed of external-state labels a_i and b_j as their letters. We also use $k_B = \sum_{j=1}^q k_{b_j}$ for the overall momentum associated with the word $B = b_1 b_2 \dots b_q$. The recursive step in (18) removes the last leg j in the first entry of $S_{\alpha'}(\cdot|\cdot)_1$ which is not necessarily in the last position in the second entry. The subscript of $S_{\alpha'}(\sigma|\beta)_1$ indicates that the entries in (18) depend on both k_1 and the momenta $k_2, k_3 \dots, k_{n-2}$ associated with the permutations σ, β .

Similar to the integration-by-parts and monodromy relations (14) and (15), the KLT relations (17) can be elegantly understood in terms of intersection theory [28] where they follow from the twisted period relations [33]. Another recent mathematical approach to the KLT relations can be found in [1].

The permutations $1, \sigma, n, n-1$ and $1, \beta, n-1, n$ in (17) reflect a particular basis choice of twisted homologies that is tailored to simplify the KLT matrix (18): The three legs $1, n-1, n$ are kept in adjacent positions, and the sets of integration cycles for $Z(1, \sigma, n, n-1|\tau)$ and $Z(1, \beta, n-1, n|\rho)$ in (17) are related through the transposition $n-1 \leftrightarrow n$. With this choice of bases, the entries of $S_{\alpha'}(\sigma|\beta)_1$ do not depend on k_{n-1} or k_n .

Given the α' -expansion of the disk integrals $Z(\tau|\rho)$, the KLT relations (17) in principle determine the analogous expansion of $J(\tau|\rho)$. However, already the four-point example (16) reveals the shortcoming of the KLT relations that both of its ingredients $Z(\tau|\rho)$ and $S_{\alpha'}(\sigma|\beta)_1$ carry spurious contributions of $\zeta(2k)$, $k \in \mathbb{N}$, which are absent in the final result (8).

At $n \geq 5$ points, similar cancellations have been observed [4] by inserting explicit α' -expansions of disk integrals into KLT formulae equivalent to (17). In the following we will not use the KLT relations. We rather give a general proof that the observed patterns of MZVs in sphere integrals are governed by the single-valued map.

3. THE MAIN RESULT

3.1. Single-valued iterated integrals and single-valued MZVs. The notion of single-valued (motivic³) MZVs is based on the representation of generic MZVs (1) in terms of multiple (harmonic) polylogarithms at unit argument (see [34] for the general definition of iterated integrals I)

$$(19) \quad I(0, a_1 a_2 \dots a_w, z) = \int_0^z \frac{dt}{t - a_w} I(0, a_1 a_2 \dots a_{w-1}, t), \quad I(0, z) = 1,$$

$$(20) \quad \zeta(n_1, n_2, \dots, n_r) = (-1)^r I(0, \underbrace{100 \dots 0}_{n_1} \underbrace{100 \dots 0}_{n_2} \dots \underbrace{100 \dots 0}_{n_r}, 1),$$

where $z \in \mathbb{C}$. For each choice of $a_1, a_2, \dots, a_w \in \{0, 1\}$, a construction by Francis Brown [18] provides a unique single-valued iterated integral $\mathcal{I}(0, a_1 a_2 \dots a_w, z)$. The latter can be considered as iteratively performing ‘single-valued integrations’ from the base point 0 to z in complete analogy to the analytic integration in (19).

In such single-valued multiple polylogarithms the monodromies of (19) around $t = 0, 1, \infty$ are annihilated by anti-holomorphic admixtures, e.g.

$$(21) \quad \mathcal{I}(0, 1, z) = I(0, 1, z) + I(0, 1, \bar{z}), \quad \mathcal{I}(0, 10, z) = I(0, 10, z) + I(0, 0, z)I(0, 1, \bar{z}) + I(0, 01, \bar{z}), \\ \mathcal{I}(0, 100, z) = I(0, 100, z) + I(0, 00, z)I(0, 1, \bar{z}) + I(0, 0, z)I(0, 01, \bar{z}) + I(0, 001, \bar{z}).$$

While the holomorphic differentials $\frac{\partial}{\partial z}$ of $I(0, \dots, z)$ are preserved by the $\mathcal{I}(0, \dots, z)$, the general connection between \mathcal{I} and I is more complicated than suggested in the above examples (a Maple implementation is [35]). By analogy with (20), single-valued MZVs (and the corresponding single-valued map sv) are defined as single-valued multiple polylogarithms at unit argument [19, 5],

$$(22) \quad \zeta_{\text{sv}}(n_1, n_2, \dots, n_r) = (-1)^r \mathcal{I}(0, \underbrace{100 \dots 0}_{n_1} \underbrace{100 \dots 0}_{n_2} \dots \underbrace{100 \dots 0}_{n_r}, 1) \\ \text{sv} : \zeta(n_1, n_2, \dots, n_r) \rightarrow \zeta_{\text{sv}}(n_1, n_2, \dots, n_r).$$

At the level of Riemann zeta values, single-valued MZVs (22) take the simple form

$$(23) \quad \zeta_{\text{sv}}(2k) = 0, \quad \zeta_{\text{sv}}(2k+1) = 2\zeta(2k+1),$$

while higher-depth instances such as

$$(24) \quad \zeta_{\text{sv}}(3, 5) = -10\zeta(3)\zeta(5), \quad \zeta_{\text{sv}}(3, 5, 3) = 2\zeta(3, 5, 3) - 2\zeta(3)\zeta(3, 5) - 10\zeta(3)^2\zeta(5),$$

are most conveniently understood in terms of the f -alphabet for MZVs [5, 36].

In the f -alphabet (motivic) iterated integrals become words in some alphabet which reflects the number-theoretical contents of the iterated integral. The f -alphabet exists for arbitrary $a_1, \dots, a_w \in \mathbb{C}$, in which case the iterated integrals (19) are hyperlogarithms. (Single) logarithms are primitive, i.e. they are represented by a single letter (of weight one). The product becomes shuffle \mathfrak{m} , and there is some admixture of polynomial type from pure periods (integrals without boundary which in the case of hyperlogarithms are polynomials in $2\pi i$).

Iterated integrals in several analytic variables are represented by words with purely analytic letters⁴. In an f -alphabet with purely analytic letters the sv map on a word w is given by

$$(25) \quad \text{sv } w = \sum_{w=uv} \bar{u} \mathfrak{m} v,$$

where \bar{u} is u in reversed order (and $\bar{\bullet}$ is complex conjugation). Moreover, $\text{sv } 2\pi i = 0$. In physical terminology the f -alphabet can be considered as a complete symbol [39]. In particular, the conversion into the f -alphabet has trivial kernel, so that no information is lost when one uses the f -alphabet.

³The single-valued map is only proved to exist in the motivic context [5].

⁴In the context of quantum field theory, iterated integrals with non-analytic letters also play a prominent role [37, 38]. Handling these objects is more complicated. Here, we only need the straightforward analytic case.

In pure mathematics the sv map exists as evaluation of ‘deRham’ periods in a very general motivic context. Here, we only use sv as the map

$$(26) \quad \text{sv}: I(0, a_1 a_2 \dots a_w, z) \mapsto \mathcal{I}(0, a_1 a_2 \dots a_w, z)$$

(which is consistent with (22)). In general, there exist relations between iterated integrals (e.g. for MZVs). A priori it is unclear (surprising even) that the map sv is well-defined (i.e. it is consistent with all relations). However, the sv-map on $I(0, a_1 a_2 \dots a_w, z)$ can be proved to have the following three natural properties:

- (i) The sv-map is well-defined.
- (ii) The sv-map commutes with evaluation.
- (iii) The sv-map extends to several (analytic) variables. I.e. $\mathcal{I}(0, a_1 a_2 \dots a_w, z)$ is single-valued in all variables a_1, a_2, \dots, a_w, z of its letters.

These results have a deep origin in motivic algebraic geometry. The Ihara action [40] plays a major role in the proof of property (i) for iterated integrals. Property (i) is theorem 1.1 in [5] and property (ii) in the context of multiple polylogarithms is corollary 5.4 in [5]. More on the evaluation of hyperlogarithms at special values of the arguments can be found in [20].

Property (iii) can be proved in the f -alphabet [36]:

Proof of (iii). In the general hyperlogarithmic context, monodromies can be expressed in terms of an ‘infinitesimal’ object, $\mathcal{M} = \exp(m)$, Here, m can be considered as picking the part of the monodromy which is proportional to $2\pi i$. Note that m is a derivative (i.e. it obeys the Leibniz rule). In the f -alphabet, m is obtained from the first letter on the Betti side (here, the left-hand side) [39],

$$(27) \quad m(aw) = m(a)w,$$

where a is a letter and w is a word.

Expressions with trivial monodromy lie in the kernel of m .

For hyperlogarithms, the only functions represented by single letters in the f -alphabet are logarithms (all logarithms are ‘primitives’ of weight one). Hence, only words with logarithms (like $I(0, a_1, z) = \log(1 - z/a_1)$) as first letters contribute to the differential monodromy m . For such a logarithm the differential monodromy around $z = a_1$ is $2\pi i$. The complex conjugate letter $\log(1 - \bar{z}/\bar{a}_1)$ has differential monodromy $-2\pi i$ around $z = a_1$ (this also remains true if one considers the monodromy of the variable a_1 around a fixed value of z). From this we conclude that in the f -alphabet for hyperlogarithms single-valuedness means that all words not beginning in constants come in pairs with complex conjugate first letters.

For a letter a we define $\partial_a aw = w$ (clipping off the first Betti letter) and $\partial_a bw = 0$ if $b \neq a$. Note that ∂_a is a differential with respect to the shuffle product. Because of the monodromy property of the f -alphabet, the proof of property (iii) reduces to showing that

$$(28) \quad \partial_a \text{sv} w = \partial_{\bar{a}} \text{sv} w$$

for all words w and all letters a (with complex conjugate \bar{a}). From (25) we have

$$(29) \quad \partial_a \text{sv} w = \sum_{w=uv} [(\partial_a \bar{u}) \text{III} v + \bar{u} \text{III} \partial_a v] = \sum_{w=uv} \bar{u} \text{III} \partial_a v.$$

Likewise,

$$(30) \quad \partial_{\bar{a}} \text{sv} w = \sum_{w=uv} (\partial_{\bar{a}} \bar{u}) \text{III} v.$$

Both expressions on the right-hand sides are equivalent to

$$(31) \quad \sum_{w=uv} \bar{u} \text{III} v$$

which completes the proof. □

Note that property (iii) means that single-valued integration with respect to any variable of a single-valued iterated integral is single-valued in all variables. A priori, this property of ‘single-valued integration’ is as mysterious as properties (i) and (ii). Single-valued integration was originally introduced by Francis Brown using generating functions [5]. In practice, it is more convenient to use a bootstrap algorithm first defined in [19]. A practical and fully general approach uses a commutative hexagon [37, 38].

Also note that property (iii) relates $\mathcal{I}(0, a_1 a_2 \dots a_w, z)$ to the single-valued multiple polylogarithms in more than one variable constructed in [41, 42].

3.2. The claim. The single-valued map (23) of Riemann zeta values relates the four-point integrals of section 2.1 at the level of their α' -expansions in (6) and (8),

$$(32) \quad J(1, 2, 3, 4|1, 2, 4, 3) = \text{sv } Z(1, 2, 3, 4|1, 2, 4, 3),$$

where sv is understood on the expansion in the parameters s_{ij} . By $\zeta_{\text{sv}}(2k) = 0$, the sv map rationalizes the trigonometric functions $\sin(\pi s_{ij}) = \pi s_{ij} \exp\left(-2 \sum_{k=1}^{\infty} \frac{\zeta(2k)}{2k} s_{ij}^{2k}\right)$ in the monodromy relations (15), $\text{sv } \sin(\pi s_{ij})/\pi = s_{ij}$. Hence, the observation (32) extends to all four-point disk and sphere integrals of the general form (9) and (10).

The general conjecture we want to prove in this work concerns the striking connection between n -point disk and sphere integrals in (9) and (10) via [17]

$$(33) \quad J(\tau|\rho) = \text{sv } Z(\tau|\rho) \quad \forall \tau, \rho \in S_n,$$

see [4] for equivalent conjectures on an $(n-3)! \times (n-3)!$ basis of disk and sphere integrals. In other words, (33) identifies sphere integrals J as single-valued disk integrals sv Z , where the anti-meromorphic part $(\bar{z}_{\tau(1),\tau(2)} \dots \bar{z}_{\tau(n),\tau(1)})^{-1}$ of the sphere integrand reflects the ordering of the integration cycle $-\infty \leq z_{\tau(1)} \leq z_{\tau(2)} \leq \dots \leq z_{\tau(n)} \leq \infty$ on the disk boundary. Again, the action of the sv map on the trigonometric functions $\text{sv } \sin(\pi s_{ij})/\pi = s_{ij}$ ensures consistency of (33) with the integration-by-parts and monodromy relations (14) and (15) [17].

Reducing the sphere integral $J(\tau|\rho)$ to a single-valued disk integral has both a conceptual and a practical advantage over the KLT formula: The low-energy expansion of (33) bypasses the spurious appearance of MZVs beyond $\zeta_{\text{sv}}(n_1, \dots, n_r)$, and the summation over $(n-3)! \times (n-3)!$ terms⁵ on the right-hand side of (17) is replaced by a single term sv $Z(\tau|\rho)$.

3.3. The proof. As the main result of this work, this section is dedicated to a proof of (33). We emphasize again that the proof is subject to the restrictions detailed at the end of the introduction.

For ease of notation, we assume the first slot of the integrals $Z(\tau|\rho)$ and $J(\tau|\rho)$ to comprise the identity permutation $\tau = 1, 2, \dots, n$. This assumption does not cause any loss of generality since all the other disk and sphere integrals with the same relative permutation $\rho \circ \tau^{-1}$ can be inferred by relabellings of the subscripts $1 \leq i, j \leq n$ of s_{ij} . Moreover, it will be convenient to pick an SL_2 frame where $(z_1, z_{n-1}, z_n) \rightarrow (0, 1, \infty)$, such that

$$(34) \quad Z(1, 2, \dots, n|\rho) = (-1)^{n-3} \int_{0 \leq z_2 \leq z_3 \leq \dots \leq z_{n-2} \leq 1} dz_2 dz_3 \dots dz_{n-2} \prod_{1 \leq i < j < n} |z_{i,j}|^{s_{ij}} f(\rho)$$

$$(35) \quad J(1, 2, \dots, n|\rho) = -\frac{1}{\pi^{n-3}} \int_{\mathbb{C}^{n-3}} \frac{d^2 z_2 d^2 z_3 \dots d^2 z_{n-2}}{\bar{z}_{1,2} \bar{z}_{2,3} \dots \bar{z}_{n-3,n-2} \bar{z}_{n-2,n-1}} \prod_{1 \leq i < j < n} |z_{i,j}|^{2s_{ij}} f(\rho).$$

The form of the meromorphic integrand

$$(36) \quad f(\rho) := \lim_{z_n \rightarrow \infty} \frac{z_n^2}{z_{\rho(1),\rho(2)} z_{\rho(2),\rho(3)} \dots z_{\rho(n-1),\rho(n)} z_{\rho(n),\rho(1)}},$$

does not affect the subsequent arguments. The values $\bar{z}_1 = 0$ and $\bar{z}_{n-1} = 1$ are meant to be inserted in the denominator of (35) and subsequent expressions.

⁵The KLT formula (17) may also be rewritten more compactly with $(n-3)! \left(\lceil \frac{n}{2} \rceil - 2\right)! \left(\lfloor \frac{n}{2} \rfloor - 1\right)!$ terms [30, 31].

The integrals Z (open string) and J (closed string) are connected by a Betti-deRham duality [43, 44]: In (34) the chain of integration is bounded by the identities $z_i = z_{i+1}$ for $i = 1, \dots, n-2$. Likewise, the integrand in (35) has the anti-meromorphic singular divisor $\cup_{i=1}^{n-2} \{\bar{z}_i = \bar{z}_{i+1}\}$ which is the deRham version of the chain of integration in (34). Accordingly, $J(\tau|\rho)$ becomes the deRham analogue of $Z(\tau|\rho)$. It is explained in [5] that single-valued MZVs are evaluations of deRham periods (after a projection from motivic periods into deRham periods which suppresses $2\pi i$, see also [20]). So, it is natural that J is the image of Z under the single-valued map. These statements, however, do not have the status of a theorem so that we need a proof of the result (33).

Proof of (33). We will iteratively integrate (34) and (35) over the variables z_2, z_3, \dots, z_{n-2} . Let $Z_i(z_{i+1}, \dots, z_{n-2})$ and $J_i(z_{i+1}, \dots, z_{n-2})$ denote the result after the $(i-1)$ st integration, i.e.

$$(37) \quad Z_i(z_{i+1}, \dots, z_{n-2}) := (-1)^{n-3} \int_{0 \leq z_2 \leq z_3 \leq \dots \leq z_i \leq z_{i+1}} dz_2 dz_3 \dots dz_i \prod_{1 \leq a < b < n} |z_{a,b}|^{s_{ab}} f(\rho)$$

$$(38) \quad J_i(z_{i+1}, \dots, z_{n-2}) := -\frac{1}{\pi^{n-3}} \int_{\mathbb{C}^{i-1}} \frac{d^2 z_2 d^2 z_3 \dots d^2 z_i}{\bar{z}_{1,2} \bar{z}_{2,3} \dots \bar{z}_{n-3,n-2} \bar{z}_{n-2,n-1}} \prod_{1 \leq a < b < n} |z_{a,b}|^{2s_{ab}} f(\rho).$$

The functions $Z_1(z_2, \dots, z_{n-2})$ and $J_1(z_2, \dots, z_{n-2})$ at $i = 1$ are given by the integrands of (34) and (35), respectively.

We will show by induction that

$$(39) \quad J_i(z_{i+1}, \dots, z_{n-2}) = \frac{(-1)^{n-i+1} \text{sv } Z_i(z_{i+1}, \dots, z_{n-2})}{\pi^{n-2-i} \bar{z}_{1,i+1} \bar{z}_{i+1,i+2} \dots \bar{z}_{n-2,n-1}}$$

for all $i = 1, \dots, n-2$. Because $Z(1, 2, \dots, n|\rho) = Z_{n-2}(\emptyset)$ and $J(1, 2, \dots, n|\rho) = J_{n-2}(\emptyset)$, this implies the theorem (33) (because $\bar{z}_{1,n-1} = -1$).

Note that the absolute values in the numerators of (37) and (38) play completely different roles in both cases. In Z_1 there exist no complex conjugate variables and $|z_{a,b}|$ with $a < b$ is $-z_{a,b}$. In fact, the only motivation for employing the absolute values for disk integrals stems from (9), where the integrand does not need any explicit reference to the permutation τ of the integration cycle. We consider the numerator as a generating series of logarithms with the expansion parameters s_{ab} . In J_1 the numerator is a generating series of logarithms in $|z_{a,b}|^2 = z_{a,b} \bar{z}_{a,b}$. Because

$$(40) \quad \text{sv } \log(x - y) = \log[(x - y)(\bar{x} - \bar{y})]$$

for any complex numbers or variables x, y , equation (39) holds for $i = 1$.

Now, assume (39) holds for i . In the calculation of Z_{i+1} , the integrand may have a singularity at $z_{i+1} = z_{i+2}$ or at $z_{i+1} = 0$. In these cases, one has to subtract the asymptotic expansion at the singular locus which will be exemplified in appendix A.⁶ The subtraction at $z_{i+1} = z_{i+2}$ is of the form $c|z_{i+1,i+2}|^{s-1}$ for some $c = c(z_{i+2}, \dots, z_{n-2})$ which is constant in z_{i+1} but may depend on the integration variables z_{i+2}, \dots, z_{n-2} of later steps. The exponent in $|z_{i+1,i+2}|^{s-1}$ refers to a sum $s = \sum s_{ab}$ for some pairs a, b that are determined by previous integration steps. Assuming that⁷ $s > 0$, the subtraction can trivially be integrated from $0 = z_1$ to z_{i+2} yielding $-\frac{c}{s} |z_{i+1,i+2}|^s$ (providing a pole in $s = 0$). The analogous result holds for a singularity at $z_{i+1} = 0$.

The systematics of the kinematic poles of disk integrals generated in this way have been discussed in the literature from various perspectives [45, 9, 8]. Note that for the present proof, we only need the existence of such a subtraction scheme, i.e. the four- and five-point examples in appendix A are merely displayed for illustrative purposes. The closed-string analogues of the disk integrals with kinematic poles can be addressed with almost identical subtraction schemes,

⁶A general approach by F. Brown and C. Dupont to this subtraction of singularities uses ‘dihedral coordinates’ [1].

⁷Negative values of s can be addressed via analytic continuation, based on the same form of the primitive that arises for $s > 0$.

where the primitives involve factors of $|z_{i+1,i+2}|^{2s}$ rather than $|z_{i+1,i+2}|^s$. All the intermediate steps of the open-string and closed-string subtraction scheme are related through the sv map as one can see from the Taylor expansions of $|z_{i+1,i+2}|^{2s}$ and $|z_{i+1,i+2}|^s$.

After the subtraction, the integrands of (37) and (38) have an integrable expansion at $s_{ij} = 0$ and we can consider the integrand as a generating series in the s_{ij} . With this prescription we define the primitive F_i of Z_i with respect to z_{i+1} and obtain

$$(41) \quad Z_{i+1} = \int_0^{z_{i+2}} dz_{i+1} Z_i = F_i(z_{i+2}) - F_i(0).$$

In general, the right-hand side of (41) is a series of Laurent type whose coefficients are iterated integrals in the letters $0, 1, z_k$ for $k = i+2, \dots, n-2$.

By the inductive assumption we have

$$(42) \quad J_{i+1} := \int_{\mathbb{C}} d^2 z_{i+1} J_i = \int_{\mathbb{C}} d^2 z_{i+1} \frac{(-1)^{n-i+1} \text{sv} Z_i}{\pi^{n-2-i} \bar{z}_{1,i+1} \bar{z}_{i+1,i+2} \dots \bar{z}_{n-2,n-1}}.$$

We calculate the integral with the residue theorem of section 2.8 in [19]. To do so we need a single-valued primitive of the integrand with respect to the holomorphic variable z_{i+1} . By single-valued integration – see property (iii) in section 3.1 – this primitive is

$$(43) \quad \mathcal{F}_i := \frac{(-1)^{n-i+1} \text{sv} F_i}{\pi^{n-2-i} \bar{z}_{1,i+1} \bar{z}_{i+1,i+2} \dots \bar{z}_{n-2,n-1}}.$$

Because the denominator of \mathcal{F}_i is of degree two in \bar{z}_{i+1} , its anti-residue at infinity (the residue with respect to the anti-holomorphic variable \bar{z}_{i+1}) vanishes. Moreover, \mathcal{F}_i has simple poles at $\bar{z}_{i+1} = \bar{z}_1 = 0$ and at $\bar{z}_{i+1} = \bar{z}_{i+2}$ whose anti-residues are obtained by substitution. From the residue theorem in [19] (using Stokes' theorem)⁸ we obtain

$$(44) \quad \begin{aligned} J_{i+1} &= \int_{\mathbb{C}} d^2 z_{i+1} \frac{\partial}{\partial z_{i+1}} \mathcal{F}_i \\ &= \frac{(-2\pi i)}{2i} \frac{(-1)^{n-i+1} [(\text{sv} F_i)(z_{i+2}) - (\text{sv} F_i)(0)]}{\pi^{n-2-i} \bar{z}_{1,i+2} \dots \bar{z}_{n-2,n-1}} \\ &= \frac{(-1)^{n-i} \text{sv} Z_{i+1}}{\pi^{n-3-i} \bar{z}_{1,i+2} \dots \bar{z}_{n-2,n-1}}. \end{aligned}$$

Because the evaluation of F_i commutes with the sv-map – see property (ii) in section 3.1 – this reproduces the shifted form $i \rightarrow i+1$ of the inductive assumption (39) and therefore completes the induction. \square

The proof confirms the result of [46, 2] that the Laurent series of Z has MZV coefficients and provides a method to calculate them which closely follows the lines of [9, 8]. At the same time, it clarifies that the coefficients of J are single-valued MZVs which can be inferred from open-string results on Z without any reference to KLT relations (17).

4. CONCLUSIONS

In this work, we have proved that the moduli-space integrals in n -point tree-level amplitudes of open and closed strings are related by the sv map, confirming the conjectures of [4, 16, 17]. More precisely, sphere integrals are expressed as single-valued disk integrals, where the singular parts of the anti-meromorphic sphere integrand are traded for an integration cycle on the disk boundary related by Betti-deRham duality. Our proof puts an intriguing web of connections between

⁸Schematically, after using Stokes' theorem we use the residue theorem in the following way in passing to the second line of (44)

$$\oint_{\partial(\mathbb{C} \setminus \{z_a, z_c\})} d\bar{z}_b \frac{f(z_b)}{\bar{z}_{ab} \bar{z}_{bc}} = -\frac{2\pi i}{\bar{z}_{ac}} (f(z_c) - f(z_a)),$$

where the function f is regular at $z_b = z_a, z_c$. Note that the 'boundary' of $\mathbb{C} \setminus \{z_a, z_c\}$ has negative orientation. A proof of this identity is in [19], see theorem 2.29.

low-energy interactions of gauge- and gravity states in different string theories [17, 21, 22] on firm grounds. These results go beyond the reach of the KLT relations (17) as well as the known string dualities [47, 48, 49] and call for various directions of follow-up research.

In the same way as the notion of a single-valued map applies to a variety of periods [20], the sv relations between string tree-level amplitudes should have an echo at loop level. At genus one, this gives rise to expect a relation between elliptic multiple zeta values [50] in open-string α' -expansions [51, 52] and modular graph functions in closed-string expansions [53, 54, 55, 56]⁹.

Single-valued polylogarithms and MZVs were found to play a key role in one-loop amplitudes of closed superstrings [59, 56]. Moreover, first explicit connections between open- and closed-string results at genus one were established in [60], along with an empirically motivated conjecture for the form of an elliptic single-valued map. Since the proof of this work only relies on general properties of the genus-zero integrals – such as singularities of the integrands and the existence of suitable primitives – it is conceivable that similar methods can be applied to timely research problems at genus one and beyond.

At higher genus, the α' -expansion of moduli-space integrals of closed strings was pioneered in [61, 62], and the last months witnessed tremendous progress in understanding their systematics and degenerations [63, 64]. However, a higher-genus framework of elliptic multiple zeta values is still lacking, so the knowledge of open-string low-energy expansions is very limited. We hope that the ideas of the proof in this work are helpful to identify a language for loop-level integrals in open- and closed-string amplitudes that is tailored to expose their relations.

ACKNOWLEDGEMENTS

We are grateful to the Hausdorff Institute Bonn for providing stimulating atmosphere, support, and hospitality through the Hausdorff Trimester Program “Periods in Number Theory, Algebraic Geometry and Physics” and the workshop “Amplitudes and Periods” where this work was initiated. Moreover, we would like to thank Francis Brown, Pierre Vanhove and Federico Zerbini for inspiring discussions. The research of Oliver Schlotterer was supported in part by Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development Canada and by the Province of Ontario through the Ministry of Research, Innovation and Science. Oliver Schnetz is supported by DFG grant SCHN 1240/2.

APPENDIX A. POLE SUBTRACTIONS

In this appendix, we illustrate the subtraction of singularities in the successive integration over disk punctures, see the discussion below (40). In the representation (34) of disk integrals, the rational function $f(\rho)$ defined in (36) may contribute a pole in $z_{i+1, i+2}$ or z_{i+1} to the integrand of $\int_0^{z_{i+2}} dz_{i+1}$ in the induction step of the main proof. An explicit realization of subtraction schemes will now be spelled out for certain four- and five-point integrals which reflect the key features of the strategy at n points. Still, we reiterate that the proof in section 3.3 only requires the existence of a subtraction scheme, i.e. the details of the subsequent examples are just given to illustrate the general mechanism.

Similar subtractions were done in the more complicated framework of ϕ^4 quantum field theory in [37] to obtain the seven loop beta-function (see Figure 7 and Conjecture 4.12). In tree-level amplitudes of string theories, the singularities are logarithmic once the disk and sphere integrals are brought into the form of $Z(\tau|\rho)$ and $J(\tau|\rho)$ via integration by parts. Since there is no need for dimensional regularization in string tree-level amplitudes, the analogue of Conjecture 4.12 in [37] becomes a lemma that follows from blowing up all singular loci in the integrand. See [65] for the application of the concept of blowing up singularities in the context of quantum field theory.

⁹Modular graph functions are believed to fall into the more general framework of non-holomorphic modular forms described in [57, 58].

A.1. Four-point examples. In an $\text{SL}_2(\mathbb{R})$ frame with $(z_1, z_3, z_4) \rightarrow (0, 1, \infty)$, we consider the following instances of the disk and sphere integrals (34) and (35) with a single kinematic pole,

$$(45) \quad Z(1, 2, 3, 4|1, 2, 4, 3) = - \int_0^1 dz_2 \frac{z_2^{s_{12}} (1-z_2)^{s_{23}}}{z_2} = - \int_0^1 dz_2 \frac{z_2^{s_{12}}}{z_2} \left(\underbrace{(1-z_2)^{s_{23}} - 1}_{(i)} + \underbrace{1}_{(ii)} \right)$$

$$(46) \quad Z(1, 2, 3, 4|1, 4, 2, 3) = - \int_0^1 dz_2 \frac{z_2^{s_{12}} (1-z_2)^{s_{23}}}{1-z_2} = - \int_0^1 dz_2 \frac{(1-z_2)^{s_{23}}}{1-z_2} \left(\underbrace{z_2^{s_{12}} - 1}_{(iii)} + \underbrace{1}_{(iv)} \right)$$

$$(47) \quad J(1, 2, 3, 4|1, 2, 4, 3) = \int_{\mathbb{C}} \frac{d^2 z_2 |z_2|^{2s_{12}} |1-z_2|^{2s_{23}}}{\pi z_2 \bar{z}_2 (\bar{z}_2 - 1)} = \int_{\mathbb{C}} \frac{d^2 z_2 |z_2|^{2s_{12}} \left(\underbrace{|1-z_2|^{2s_{23}} - 1}_{(v)} + \underbrace{1}_{(vi)} \right)}{\pi z_2 \bar{z}_2 (\bar{z}_2 - 1)}$$

$$(48) \quad J(1, 2, 3, 4|1, 4, 2, 3) = \int_{\mathbb{C}} \frac{d^2 z_2 |z_2|^{2s_{12}} |1-z_2|^{2s_{23}}}{\pi (1-z_2) \bar{z}_2 (\bar{z}_2 - 1)} = \int_{\mathbb{C}} \frac{d^2 z_2 |1-z_2|^{2s_{23}} \left(\underbrace{|z_2|^{2s_{12}} - 1}_{(vii)} + \underbrace{1}_{(viii)} \right)}{\pi (1-z_2) \bar{z}_2 (\bar{z}_2 - 1)},$$

where the shorthands (i) to (viii) refer to the full-fledged integrals after isolating the highlighted terms in the sums (...) of the integrand, e.g.

$$(49) \quad (v) = \int_{\mathbb{C}} \frac{d^2 z_2 |z_2|^{2s_{12}} (|1-z_2|^{2s_{23}} - 1)}{\pi z_2 \bar{z}_2 (\bar{z}_2 - 1)}.$$

The subtractions on the right-hand side are tailored to isolate the field-theory limits

$$(50) \quad Z(1, 2, 3, 4|1, 2, 4, 3) = -\frac{1}{s_{12}} + O(\alpha'), \quad J(1, 2, 3, 4|1, 2, 4, 3) = -\frac{1}{s_{12}} + O(\alpha')$$

$$(51) \quad Z(1, 2, 3, 4|1, 4, 2, 3) = -\frac{1}{s_{23}} + O(\alpha'), \quad J(1, 2, 3, 4|1, 4, 2, 3) = -\frac{1}{s_{23}} + O(\alpha'),$$

which can be straightforwardly generated from the integrals

$$(52) \quad (ii) = - \int_0^1 dz_2 z_2^{s_{12}-1} = - \frac{z_2^{s_{12}}}{s_{12}} \Big|_{z_2=0}^{z_2=1} = -\frac{1}{s_{12}}$$

$$(53) \quad (iv) = - \int_0^1 dz_2 (1-z_2)^{s_{23}-1} = \frac{(1-z_2)^{s_{23}}}{s_{23}} \Big|_{z_2=0}^{z_2=1} = -\frac{1}{s_{23}}$$

$$(54) \quad (vi) = \int_{\mathbb{C} \setminus \{0,1\}} \frac{d^2 z_2 |z_2|^{2s_{12}}}{\pi z_2 \bar{z}_2 (\bar{z}_2 - 1)} = \frac{1}{\pi s_{12}} \int_{\mathbb{C} \setminus \{0,1\}} d^2 z_2 \frac{\partial}{\partial z_2} \frac{|z_2|^{2s_{12}}}{\bar{z}_2 (\bar{z}_2 - 1)}$$

$$= \frac{1}{2\pi i s_{12}} \oint_{\partial(\mathbb{C} \setminus \{0,1\})} \frac{d\bar{z}_2 |z_2|^{2s_{12}}}{\bar{z}_2 (\bar{z}_2 - 1)}$$

$$= \frac{1}{2\pi i s_{12}} \left\{ - \frac{2\pi i |z_2|^{2s_{12}}}{\bar{z}_2 - 1} \Big|_{z_2=0} - \frac{2\pi i |z_2|^{2s_{12}}}{\bar{z}_2} \Big|_{z_2=1} \right\} = -\frac{1}{s_{12}}.$$

The evaluation of (viii) is completely analogous to (vi) and yields $-1/s_{23}$. Note that, following the proof in section 3.3, the meromorphic parts of the primitives in (52) and (54) are identical. So, the fact that (vi) = sv(ii) is clear from the general arguments given above and confirmed by the inspection of the final result $-1/s_{12}$ in both cases, where the action of sv trivializes.

The integrands in the curly bracket of (i), (v) and (iii), (vii) are designed to be regular as $z_2 \rightarrow 0$ and $z_2 \rightarrow 1$, respectively. This renders the integrated expressions non-singular w.r.t. s_{ij} , and the arguments in the proof in section 3.3 can be applied to the series in $\log(z_{ij})$ and $\log|z_{ij}|^2$ without the need for further subtractions: Along with each monomial in $s_{12}^m s_{23}^n$ with $m, n \geq 0$, the

holomorphic primitives of $(\log z_2)^m (\log(1-z_2))^n / z_2$ and $(\log |z_2|^2)^m (\log |1-z_2|^2)^n / z_2$ are related by the sv map and ultimately evaluated at $z_2 = 1$. Hence, at the level of the resulting MZVs,

$$(55) \quad (v) = \text{sv}(i) , \quad (vii) = \text{sv}(iii) .$$

A.2. Five-point examples: non-overlapping singularities. Starting from five-point disk and sphere integrals, the residues of the kinematic poles are by themselves series in s_{ij} with MZV coefficients. As a first example, we consider the integral

$$(56) \quad \begin{aligned} Z(1, 2, 3, 4, 5|1, 2, 5, 3, 4) &= \int_0^1 dz_3 \int_0^{z_3} dz_2 \frac{z_2^{s_{12}} z_3^{s_{13}} z_{32}^{s_{23}} (1-z_2)^{s_{24}} (1-z_3)^{s_{34}}}{z_2 (z_3-1)} \\ &= -\frac{1}{s_{12}s_{34}} + O(\alpha^0) \end{aligned}$$

in an $\text{SL}_2(\mathbb{R})$ frame with $(z_1, z_4, z_5) \rightarrow (0, 1, \infty)$, where the poles s_{12}^{-1} and s_{34}^{-1} stem from different endpoints $z_2 \rightarrow 0$ and $z_3 \rightarrow 1$ of the integration domain $0 \leq z_2 \leq z_3 \leq 1$. As an analogue of the subtraction scheme in (45) to (48), we rewrite the integrand of (56) as

$$(57) \quad \begin{aligned} Z(1, 2, 3, 4, 5|1, 2, 5, 3, 4) &= \int_0^1 dz_3 \frac{z_3^{s_{13}} (1-z_3)^{s_{34}}}{z_3-1} \\ &\int_0^{z_3} dz_2 \frac{z_2^{s_{12}}}{z_2} \left(\underbrace{z_{32}^{s_{23}} (1-z_2)^{s_{24}} - z_3^{s_{23}}}_{(\alpha)} + \underbrace{z_3^{s_{23}}}_{(\beta)} \right) . \end{aligned}$$

The contribution of (β) involves a straightforward integral over z_2 similar to (52) along with an integral over z_3 of four-point type, cf. (46),

$$(58) \quad \begin{aligned} (\beta) &= \int_0^1 dz_3 \frac{z_3^{s_{13}+s_{23}} (1-z_3)^{s_{34}}}{(z_3-1)} \frac{z_2^{s_{12}}}{s_{12}} \Big|_{z_2=0}^{z_2=z_3} \\ &= \frac{1}{s_{12}} \int_0^1 dz_3 \frac{z_3^{s_{12}+s_{13}+s_{23}} (1-z_3)^{s_{34}}}{z_3-1} \\ &= \frac{1}{s_{12}} \left(Z(1, 2, 3, 4|1, 4, 2, 3) \Big|_{s_{12} \rightarrow s_{12}+s_{13}+s_{23}}^{s_{23} \rightarrow s_{34}} \right) . \end{aligned}$$

The leading order of $Z(1, 2, 3, 4|1, 4, 2, 3)$ in (51) then yields the low-energy limit $-(s_{12}s_{34})^{-1}$ of the integral $Z(1, 2, 3, 4, 5|1, 2, 5, 3, 4)$, see (56).

The integrand of the contribution of (α) in (57) is regular at $z_2 = 0$, so the integral over z_2

$$(59) \quad H(z_3) := \int_0^{z_3} dz_2 \frac{z_2^{s_{12}}}{z_2} (z_{32}^{s_{23}} (1-z_2)^{s_{24}} - z_3^{s_{23}})$$

does not involve any singularity in s_{ij} , and the α' -expansion can be performed at the level of the $\log(z_{ij})$ in the integrand. The limit $z_3 \rightarrow 1$ of (59) is smooth and again reproduces an integral of four-point type, cf. (i) in (45)

$$(60) \quad H(1) = \int_0^1 dz_2 \frac{z_2^{s_{12}}}{z_2} ((1-z_2)^{s_{23}+s_{24}} - 1) .$$

The contribution of (α) in (57) still yields a pole in s_{34} upon integration over z_3 . This pole can be traced back to the factor of $\frac{(1-z_3)^{s_{34}}}{(z_3-1)}$, and we isolate it by the subtraction scheme

$$(61) \quad (\alpha) = \int_0^1 dz_3 \frac{(1-z_3)^{s_{34}}}{z_3-1} \left(\underbrace{z_3^{s_{13}} H(z_3) - H(1)}_{(\gamma)} + \underbrace{H(1)}_{(\delta)} \right) .$$

The integral in (53) determines

$$(62) \quad (\delta) = -\frac{H(1)}{s_{34}} ,$$

and the integrand for the contribution (γ) to (61) is regular at $z_3 \rightarrow 1$ such that

$$(63) \quad (\gamma) = \int_0^1 dz_3 \frac{(1-z_3)^{s_{34}}}{z_3-1} (z_3^{s_{13}} H(z_3) - H(1))$$

is regular in s_{34} and can be α' -expanded at the level of the integrand.

In adapting the subtraction scheme to the corresponding sphere integral

$$(64) \quad \begin{aligned} J(1, 2, 3, 4, 5|1, 2, 5, 3, 4) &= -\frac{1}{\pi^2} \int_{\mathbb{C}^2} \frac{d^2 z_2 d^2 z_3 |z_3|^{2s_{13}} |1-z_3|^{2s_{34}}}{\bar{z}_{12} \bar{z}_{23} \bar{z}_{34} (z_3-1)} \\ &\times \frac{|z_2|^{2s_{12}}}{z_2} \left(\underbrace{|z_{23}|^{2s_{23}} |1-z_2|^{2s_{24}} - |z_3|^{2s_{23}}}_{(A)} + \underbrace{|z_3|^{2s_{23}}}_{(B)} \right), \end{aligned}$$

the primitives for all contributions $(\alpha), (\beta), (\gamma), (\delta)$ have the same meromorphic parts as in the case of $J(1, 2, 3, 4, 5|1, 2, 5, 3, 4)$. In analogy with (58), we have

$$(65) \quad \begin{aligned} (B) &= -\frac{1}{2i\pi^2 s_{12}} \int_{\mathbb{C}} d^2 z_3 \oint_{\partial(\mathbb{C} \setminus \{0, z_3\})} d\bar{z}_2 \frac{|z_2|^{2s_{12}} |z_3|^{2s_{13}+2s_{23}} |1-z_3|^{2s_{34}}}{\bar{z}_{12} \bar{z}_{23} \bar{z}_{34} (z_3-1)} \\ &= \frac{1}{\pi s_{12}} \int_{\mathbb{C}} d^2 z_3 \frac{|z_3|^{2(s_{12}+s_{13}+s_{23})} |1-z_3|^{2s_{34}}}{\bar{z}_{13} \bar{z}_{34} (z_3-1)} \\ &= \frac{1}{s_{12}} \left(J(1, 2, 3, 4|1, 4, 2, 3) \Big|_{s_{12} \rightarrow s_{12}+s_{13}+s_{23}}^{s_{23} \rightarrow s_{34}} \right), \end{aligned}$$

which gives the desired expression $\text{sv}(\beta)$.

The z_2 integral of (A),

$$(66) \quad I(z_3) := \frac{1}{\pi} \int_{\mathbb{C}} d^2 z_2 \frac{|z_2|^{2s_{12}}}{z_2 \bar{z}_{12} \bar{z}_{23}} (|z_{23}|^{2s_{23}} |1-z_2|^{2s_{24}} - |z_3|^{2s_{23}})$$

is regular and the general method in the proof of the main result applies. We obtain:

$$(67) \quad I(z_3) = -\frac{1}{z_{13}} \text{sv} H(z_3).$$

Upon insertion into (64), this implies

$$(68) \quad (A) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{d^2 z_3}{\bar{z}_{13} \bar{z}_{34}} \frac{|1-z_3|^{2s_{34}}}{(z_3-1)} \left(\underbrace{|z_3|^{2s_{13}} \text{sv} H(z_3) - \text{sv} H(1)}_{(C)} + \underbrace{\text{sv} H(1)}_{(D)} \right).$$

In analogy to (54) the integral in the last term gives

$$(69) \quad (D) = -\frac{\text{sv} H(1)}{s_{34}},$$

which is identical to $\text{sv}(\delta)$ by (62). The integral (C) in (68) is regular and can be expanded in α' in the integrand. By the general method in the proof of the main result, we obtain $(C) = \text{sv}(\gamma)$ and recover $J(1, 2, 3, 4, 5|1, 2, 5, 3, 4) = \text{sv} Z(1, 2, 3, 4, 5|1, 2, 5, 3, 4)$ term by term in the subtraction scheme.

A.3. Five-point examples: nested singularities. While the singularities of the five-point example in appendix A.2 stem from different regions $z_2 \rightarrow 0$ and $z_3 \rightarrow 1$, the following disk integral acquires kinematic poles in $s_{123} := s_{12} + s_{13} + s_{23}$ from the nested singularity¹⁰ in the

¹⁰In a five-point setup, one can still avoid the nested singularities by representing (70) in a different SL_2 frame, but this is no longer true at six points. We choose the SL_2 frame with $(z_1, z_4, z_5) \rightarrow (0, 1, \infty)$ here to illustrate that the nesting of singularities does not obstruct the existence of a subtraction scheme.

integration region where $z_2, z_3 \rightarrow 0$:

$$\begin{aligned}
Z_{\text{nest}} &= -Z(1, 2, 3, 4, 5|1, 2, 3, 5, 4) - Z(1, 2, 3, 4, 5|1, 3, 2, 5, 4) \\
(70) \quad &= \int_0^1 dz_3 \int_0^{z_3} dz_2 \frac{z_2^{s_{12}} z_3^{s_{13}} z_{32}^{s_{23}} (1-z_2)^{s_{24}} (1-z_3)^{s_{34}}}{z_2 z_3} \\
&= \frac{1}{s_{12}s_{123}} + O(\alpha'^0).
\end{aligned}$$

The first step of the subtraction scheme closely follows the lines of (57)

$$(71) \quad Z_{\text{nest}} = \int_0^1 dz_3 \frac{z_3^{s_{13}} (1-z_3)^{s_{34}}}{z_3} \int_0^{z_3} dz_2 \frac{z_2^{s_{12}}}{z_2} \left(\underbrace{z_{32}^{s_{23}} (1-z_2)^{s_{24}} - z_3^{s_{23}}}_{(p)} + \underbrace{z_3^{s_{23}}}_{(q)} \right),$$

and the evaluation of the second contribution (q) is almost identical to (β) in (58),

$$(72) \quad (q) = \int_0^1 dz_3 \frac{z_3^{s_{13}+s_{23}} (1-z_3)^{s_{34}}}{z_3} \frac{z_2^{s_{12}}}{s_{12}} \Big|_{z_2=0}^{z_2=z_3} = -\frac{1}{s_{12}} \left(Z(1, 2, 3, 4|1, 2, 4, 3) \Big|_{s_{12} \rightarrow s_{123}}^{s_{23} \rightarrow s_{34}} \right).$$

The low-energy limit $Z_{\text{nest}} = \frac{1}{s_{12}s_{123}} + O(\alpha'^0)$ in (70) then stems from the leading term of the four-point integral $Z(1, 2, 3, 4|1, 2, 4, 3)$ in (50) at shifted first argument $s_{12} \rightarrow s_{123}$.

In the subtraction scheme for

$$(73) \quad (p) = \int_0^1 dz_3 \frac{z_3^{s_{13}} (1-z_3)^{s_{34}} H(z_3)}{z_3},$$

it would be tempting to closely follow the treatment of (α) in (61) and to subtract the $z_3 \rightarrow 0$ limit of the quantity $H(z_3)$ in (59). However, this limit does not admit a regular α' -expansion and we shall instead write $H(z_3) = z_3^{s_{12}+s_{23}} h(z_3)$ (which extracts the exact scaling behavior of H at $z_3 = 0$). We set $z_2 = xz_3$ in the integral representation (59) of $H(z_3)$ and obtain¹¹

$$(74) \quad h(z_3) = \int_0^1 \frac{dx}{x} x^{s_{12}} ((1-x)^{s_{23}} (1-z_3x)^{s_{24}} - 1), \quad z_3 \leq 1.$$

Since (74) is regular as $z_3 \rightarrow 0$, the appropriate analogue of (61) is

$$(75) \quad (p) = \int_0^1 dz_3 \frac{z_3^{s_{123}}}{z_3} \left(\underbrace{(1-z_3)^{s_{34}} h(z_3) - h(0)}_{(r)} + \underbrace{h(0)}_{(t)} \right),$$

where the integrand in (r) is regular as $z_3 \rightarrow 0$. The integral can be performed order by order. Finally, the pole from the nested singularity

$$(76) \quad (t) = \frac{h(0)}{s_{123}} = \frac{1}{s_{123}} \int_0^1 \frac{dx}{x} x^{s_{12}} ((1-x)^{s_{23}} - 1),$$

has a residue identical to (i) in (45).

For the corresponding sphere integral

$$\begin{aligned}
J_{\text{nest}} &= -J(1, 2, 3, 4, 5|1, 2, 3, 5, 4) - J(1, 2, 3, 4, 5|1, 3, 2, 5, 4) \\
(77) \quad &= -\frac{1}{\pi^2} \int_{\mathbb{C}^2} \frac{d^2 z_2 d^2 z_3}{\bar{z}_{12} \bar{z}_{23} \bar{z}_{34}} \frac{|z_2|^{2s_{12}} |z_3|^{2s_{13}} |1-z_3|^{2s_{34}}}{z_2 z_3} \left(\underbrace{|z_{23}|^{2s_{23}} |1-z_2|^{2s_{24}} - |z_3|^{2s_{23}}}_{(P)} + \underbrace{|z_3|^{2s_{23}}}_{(Q)} \right),
\end{aligned}$$

¹¹Note that the leading terms of the α' -expansion of (74) are given by

$$\begin{aligned}
h(z_3) &= s_{24} I(0, 10, z_3) - s_{23} \zeta(2) + s_{24}^2 I(0, 110, z_3) + s_{23}^2 \zeta(3) - s_{12} s_{24} I(0, 100, z_3) \\
&\quad + s_{12} s_{23} \zeta(3) + s_{23} s_{24} [I(0, 110, z_3) - I(0, 100, z_3)] + O(\alpha'^3),
\end{aligned}$$

see (19) for the definition of the iterated integrals $I(0, a_1 a_2 \dots a_w, z)$.

the first step of the subtraction scheme is again almost identical to (64), resulting in

$$(78) \quad (Q) = -\frac{1}{s_{12}} \left(J(1, 2, 3, 4 | 1, 2, 4, 3) \Big|_{s_{12} \rightarrow s_{123}}^{s_{23} \rightarrow s_{34}} \right),$$

which matches $\text{sv}(q)$ by (72).

The z_2 integral of (P) is again given by (66), and we will use its representation in (67),

$$(79) \quad (P) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{d^2 z_3}{\bar{z}_{13} \bar{z}_{34}} \frac{|z_3|^{2s_{13}} |1 - z_3|^{2s_{34}}}{z_3} \text{sv} H(z_3).$$

Then, we use the single-valued analogue $\text{sv} H(z_3) = |z_3|^{2s_{12} + 2s_{23}} \text{sv} h(z_3)$ of the above rewriting $H(z_3) = z_3^{s_{12} + s_{23}} h(z_3)$ with $h(z_3)$ given by (74) and employ the following subtraction scheme:

$$(80) \quad (P) = \frac{1}{\pi} \int_{\mathbb{C}} \frac{d^2 z_3}{\bar{z}_{13} \bar{z}_{34}} \frac{|z_3|^{2s_{123}}}{z_3} \left(\underbrace{|1 - z_3|^{2s_{34}} \text{sv} h(z_3) - \text{sv} h(0)}_{(R)} + \underbrace{\text{sv} h(0)}_{(T)} \right).$$

The integrand in (R) is regular as $z_3 \rightarrow 0$ and we arrive at $(R) = \text{sv}(r)$ upon order-by-order integration, cf. (75). The last term in (80) can be trivially integrated to give

$$(81) \quad (T) = \frac{\text{sv} h(0)}{s_{123}},$$

which agrees with $\text{sv}(t)$ by (76). Hence, we have checked the relation $J_{\text{nest}} = \text{sv} Z_{\text{nest}}$ at the level of all the terms in the subtraction scheme.

REFERENCES

- [1] F. Brown, C. Dupont, “Superstring amplitudes in genus 0 and 1”, talk given by F. Brown in ‘String Math’, Sendai, June 18, 2018.
- [2] F. Brown, “Multiple zeta values and periods of moduli spaces $M_{0,n}(\mathbb{R})$,” *Annales Sci. Ecole Norm. Sup.* **42** (2009) 371 [math/0606419 [math.AG]].
- [3] H. Kawai, D. C. Lewellen and S. H. H. Tye, “A Relation Between Tree Amplitudes of Closed and Open Strings,” *Nucl. Phys. B* **269** (1986) 1.
- [4] O. Schlotterer and S. Stieberger, “Motivic Multiple Zeta Values and Superstring Amplitudes,” *J. Phys. A* **46** (2013) 475401 [arXiv:1205.1516 [hep-th]].
- [5] F. Brown, “Single-valued periods and multiple zeta values,” *Forum of Mathematics, Sigma* 2, e25 (2014) [arXiv:1309.5309 [math.NT]].
- [6] J. M. Drummond and E. Ragoucy, “Superstring amplitudes and the associator,” *JHEP* **1308** (2013) 135 [arXiv:1301.0794 [hep-th]].
- [7] J. Broedel, O. Schlotterer, S. Stieberger and T. Terasoma, “All order α' -expansion of superstring trees from the Drinfeld associator,” *Phys. Rev. D* **89** (2014) no.6, 066014 [arXiv:1304.7304 [hep-th]].
- [8] C. R. Mafra and O. Schlotterer, “Non-abelian Z -theory: Berends-Giele recursion for the α' -expansion of disk integrals,” *JHEP* **1701** (2017) 031 [arXiv:1609.07078 [hep-th]].
- [9] J. Broedel, O. Schlotterer and S. Stieberger, “Polylogarithms, Multiple Zeta Values and Superstring Amplitudes,” *Fortsch. Phys.* **61** (2013) 812, [arXiv:1304.7267 [hep-th]].
- [10] J. Broedel, O. Schlotterer and S. Stieberger, <http://wwwth.mpp.mpg.de/members/stieberg/mzv/index.html>
- [11] D. Oprisa and S. Stieberger, “Six gluon open superstring disk amplitude, multiple hypergeometric series and Euler-Zagier sums,” hep-th/0509042.
- [12] S. Stieberger and T. R. Taylor, “Multi-Gluon Scattering in Open Superstring Theory,” *Phys. Rev. D* **74** (2006) 126007 [hep-th/0609175].
- [13] R. H. Boels, “On the field theory expansion of superstring five point amplitudes,” *Nucl. Phys. B* **876** (2013) 215 [arXiv:1304.7918 [hep-th]].
- [14] G. Puhlfürst and S. Stieberger, “Differential Equations, Associators, and Recurrences for Amplitudes,” *Nucl. Phys. B* **902** (2016) 186 [arXiv:1507.01582 [hep-th]].
- [15] S. Stieberger, “Constraints on Tree-Level Higher Order Gravitational Couplings in Superstring Theory,” *Phys. Rev. Lett.* **106** (2011) 111601 [arXiv:0910.0180 [hep-th]].
- [16] S. Stieberger, “Closed superstring amplitudes, single-valued multiple zeta values and the Deligne associator,” *J. Phys. A* **47** (2014) 155401 [arXiv:1310.3259 [hep-th]].
- [17] S. Stieberger and T. R. Taylor, “Closed String Amplitudes as Single-Valued Open String Amplitudes,” *Nucl. Phys. B* **881** (2014) 269 [arXiv:1401.1218 [hep-th]].
- [18] F. Brown, “Single-valued multiple polylogarithms in one variable,” *C.R. Acad. Sci. Paris, Ser. I* **338** (2004) 527-532.

- [19] O. Schnetz, “Graphical functions and single-valued multiple polylogarithms,” *Comm. Num. Theor. Phys.* **8**, no. 4 (2014) 589 [arXiv:1302.6445 [math.NT]].
- [20] F. Brown, “Notes on motivic periods,” *Comm. Num. Theor. Phys.* **11**, no. 3 (2017) 557 [arXiv:1512.06410 [math.NT]].
- [21] O. Schlotterer, “Amplitude relations in heterotic string theory and Einstein-Yang-Mills,” *JHEP* **1611** (2016) 074 [arXiv:1608.00130 [hep-th]].
- [22] T. Azevedo, M. Chiodaroli, H. Johansson and O. Schlotterer, “Heterotic and bosonic string amplitudes via field theory,” arXiv:1803.05452 [hep-th].
- [23] W. Fan, A. Fotopoulos, S. Stieberger and T. R. Taylor, “SV-map between Type I and Heterotic Sigma Models,” *Nucl. Phys. B* **930** (2018) 195 [arXiv:1711.05821 [hep-th]].
- [24] C. R. Mafra, O. Schlotterer and S. Stieberger, “Complete N-Point Superstring Disk Amplitude I. Pure Spinor Computation,” *Nucl. Phys. B* **873** (2013) 419 [arXiv:1106.2645 [hep-th]].
- [25] Y. t. Huang, O. Schlotterer and C. Wen, “Universality in string interactions,” *JHEP* **1609** (2016) 155 [arXiv:1602.01674 [hep-th]].
- [26] N. E. J. Bjerrum-Bohr, P. H. Damgaard and P. Vanhove, “Minimal Basis for Gauge Theory Amplitudes,” *Phys. Rev. Lett.* **103** (2009) 161602 [arXiv:0907.1425 [hep-th]].
- [27] S. Stieberger, “Open & Closed vs. Pure Open String Disk Amplitudes,” arXiv:0907.2211 [hep-th].
- [28] S. Mizera, “Combinatorics and Topology of Kawai-Lewellen-Tye Relations,” *JHEP* **1708** (2017) 097 [arXiv:1706.08527 [hep-th]].
- [29] S. Mizera, “Scattering Amplitudes from Intersection Theory,” *Phys. Rev. Lett.* **120** (2018) no.14, 141602 [arXiv:1711.00469 [hep-th]].
- [30] Z. Bern, L. J. Dixon, M. Perelstein and J. S. Rozowsky, “Multileg one loop gravity amplitudes from gauge theory,” *Nucl. Phys. B* **546** (1999) 423 [hep-th/9811140].
- [31] N. E. J. Bjerrum-Bohr, P. H. Damgaard, T. Sondergaard and P. Vanhove, “The Momentum Kernel of Gauge and Gravity Theories,” *JHEP* **1101** (2011) 001 [arXiv:1010.3933 [hep-th]].
- [32] J. J. M. Carrasco, C. R. Mafra and O. Schlotterer, “Abelian Z-theory: NLSM amplitudes and α' -corrections from the open string,” *JHEP* **1706** (2017) 093 [arXiv:1608.02569 [hep-th]].
- [33] K. Cho and K. Matsumoto, “Intersection theory for twisted cohomologies and twisted Riemann’s period relations I”, *Nagoya Math. J.* **139** (1995) 67–86.
- [34] K. Chen, “Algebras of Iterated Path Integrals and Fundamental Groups,” *Transactions of the American Mathematical Society* **156** (1971) 359-379.
- [35] O. Schnetz, **HyperlogProcedures**, Maple procedures available on the homepage of the author (2018).
- [36] F. Brown, “On the decomposition of motivic multiple zeta values”, in: “Galois-Teichmüller theory and arithmetic geometry,” *Math. Soc. Japan, Tokyo* (2012), 31–58 [arXiv:1102.1310 [math.NT]].
- [37] O. Schnetz, “Numbers and Functions in Quantum Field Theory,” *Phys. Rev. D* **97**, no. 8, 085018 (2018) [arXiv:1606.08598 [hep-th]].
- [38] O. Schnetz, “Generalized single-valued hyperlogarithms,” in preparation.
- [39] O. Schnetz, “Algebraic integration,” talks in Les Houches and Erlangen, June 14 and June 22, 2018.
- [40] Y. Ihara, “The Galois representation arising from $\mathbb{P}^1 - \{0, 1, \infty\}$ and Tate twists of even degree,” *Galois groups over \mathbb{Q}* , 299-313, *Math. Sci. Res. Inst. Publ.* **16**, (1989).
- [41] J. Broedel, M. Sprenger and A. Torres Orjuela, “Towards single-valued polylogarithms in two variables for the seven-point remainder function in multi-Regge-kinematics,” *Nucl. Phys. B* **915** (2017) 394 [arXiv:1606.08411 [hep-th]].
- [42] V. Del Duca, S. Druc, J. Drummond, C. Duhr, F. Dulat, R. Marzucca, G. Papathanasiou and B. Verbeek, “Multi-Regge kinematics and the moduli space of Riemann spheres with marked points,” *JHEP* **1608** (2016) 152 [arXiv:1606.08807 [hep-th]].
- [43] A. A. Beilinson, A. N. Varchenko, A. B. Goncharov, V. V. Shekhtman, “Projective Geometry and K -theory,” *Leningrad Math J.* **2**, no. 3, 523–575 (1991).
- [44] O. Schnetz, “Graphical hyperlogarithms,” talk within the trimester “Periods in Number Theory, Algebraic Geometry and Physics”, HIM, Bonn, Feb. 27, 2018.
- [45] C. R. Mafra, O. Schlotterer and S. Stieberger, “Complete N-Point Superstring Disk Amplitude II. Amplitude and Hypergeometric Function Structure,” *Nucl. Phys. B* **873** (2013) 461 [arXiv:1106.2646 [hep-th]].
- [46] T. Terasoma, “Selberg Integrals and Multiple Zeta Values”, *Compositio Mathematica* **133** (2002) 1.
- [47] C. M. Hull and P. K. Townsend, “Unity of superstring dualities,” *Nucl. Phys. B* **438** (1995) 109 [hep-th/9410167].
- [48] J. Polchinski and E. Witten, “Evidence for heterotic - type I string duality,” *Nucl. Phys. B* **460** (1996) 525 [hep-th/9510169].
- [49] E. Witten, “String theory dynamics in various dimensions,” *Nucl. Phys. B* **443** (1995) 85 [hep-th/9503124].
- [50] B. Enriquez, “Analogues elliptiques des nombres multizetas”, *Bull. Soc. Math. France* **144** (2016) 395, [arXiv:1301.3042 [math.NT]].
- [51] J. Broedel, C. R. Mafra, N. Matthes and O. Schlotterer, “Elliptic multiple zeta values and one-loop superstring amplitudes,” *JHEP* **1507** (2015) 112 [arXiv:1412.5535 [hep-th]].

- [52] J. Broedel, N. Matthes, G. Richter and O. Schlotterer, “Twisted elliptic multiple zeta values and non-planar one-loop open-string amplitudes,” *J. Phys. A* **51** (2018) no.28, 285401 [arXiv:1704.03449 [hep-th]].
- [53] M. B. Green and P. Vanhove, “The Low-energy expansion of the one loop type II superstring amplitude,” *Phys. Rev. D* **61** (2000) 104011 [hep-th/9910056].
- [54] M. B. Green, J. G. Russo and P. Vanhove, “Low energy expansion of the four-particle genus-one amplitude in type II superstring theory,” *JHEP* **0802** (2008) 020 [arXiv:0801.0322 [hep-th]].
- [55] E. D’Hoker, M. B. Green and P. Vanhove, “On the modular structure of the genus-one Type II superstring low energy expansion,” *JHEP* **1508** (2015) 041 [arXiv:1502.06698 [hep-th]].
- [56] E. D’Hoker, M. B. Green, Ö. Gürdogan and P. Vanhove, “Modular Graph Functions,” *Comm. Num. Theor. Phys.* **11** (2017) 165 [arXiv:1512.06779 [hep-th]].
- [57] F. Brown, “A class of non-holomorphic modular forms I”, *Res. Math. Sci.* **5** (2018) 5:7 [arXiv:1707.01230 [math.NT]].
- [58] F. Brown, “A class of non-holomorphic modular forms II: equivariant iterated Eisenstein integrals” [arXiv:1708.03354 [math.NT]].
- [59] F. Zerbini, “Single-valued multiple zeta values in genus 1 superstring amplitudes,” *Comm. Num. Theor. Phys.* **10** (2016) 703 [arXiv:1512.05689 [hep-th]].
- [60] J. Broedel, O. Schlotterer and F. Zerbini, “From elliptic multiple zeta values to modular graph functions: open and closed strings at one loop,” arXiv:1803.00527 [hep-th].
- [61] E. D’Hoker and M. B. Green, “Zhang-Kawazumi Invariants and Superstring Amplitudes,” *Journal of Number Theory* **144** (2014) 111 [arXiv:1308.4597 [hep-th]].
- [62] E. D’Hoker, M. B. Green, B. Pioline and R. Russo, “Matching the D^6R^4 interaction at two-loops,” *JHEP* **1501** (2015) 031 [arXiv:1405.6226 [hep-th]].
- [63] E. D’Hoker, M. B. Green and B. Pioline, “Higher genus modular graph functions, string invariants, and their exact asymptotics,” arXiv:1712.06135 [hep-th].
- [64] E. D’Hoker, M. B. Green and B. Pioline, “Asymptotics of the D^8R^4 genus-two string invariant,” arXiv:1806.02691 [hep-th].
- [65] S. Bloch, H. Esnault, D. Kreimer, “On motives associated to graph polynomials,” *Comm. Math. Phys.* **267**, no. 1, 181-225 (2006) [math.AG/0510011].

OLIVER SCHLOTTERER, MAX-PLANCK-INSTITUT FÜR GRAVITATIONSPHYSIK, ALBERT-EINSTEIN-INSTITUT, AM MÜHLENBERG 1, 14476 POTSDAM, GERMANY – AND – PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, 31 CAROLINE STREET N, WATERLOO, ON N2L 2Y5, CANADA

OLIVER SCHNETZ, DEPARTMENT MATHEMATIK, CAUERSTRASSE 11, 91058 ERLANGEN, GERMANY
E-mail address: olivers@aei.mpg.de, schnetz@mi.uni-erlangen.de