

Positive Solutions of Systems of Signed Parametric Polynomial Inequalities

Hoon Hong

North Carolina State University, Raleigh, NC, USA

`hong@ncsu.edu`

Thomas Sturm

CNRS, Inria, and the University of Lorraine, Nancy, France
MPI Informatics and Saarland University, Saarbrücken, Germany

`thomas.sturm@loria.fr`

July 27, 2018

Abstract

We consider systems of strict multivariate polynomial inequalities over the reals. All polynomial coefficients are parameters ranging over the reals, where for each coefficient we prescribe its sign. We are interested in the existence of positive real solutions of our system for all choices of coefficients subject to our sign conditions. We give a decision procedure for the existence of such solutions. In the positive case our procedure yields a parametric positive solution as a rational function in the coefficients. Our framework allows to reformulate heuristic subtropical approaches for non-parametric systems of polynomial inequalities that have been recently used in qualitative biological network analysis and, independently, in satisfiability modulo theory solving. We apply our results to characterize the incompleteness of those methods.

1 Introduction

We investigate the problem of finding a *parametric positive* solution of a system of *signed parametric* polynomial inequalities, if exists. We illustrate the problem by means of two toy examples:

$$f(x) = c_2x^2 - c_1x + c_0, \quad g(x) = -c_2x^2 + c_1x - c_0,$$

where c_2, c_1, c_0 are parameters. An expression $z(c)$ is called a parametric positive solution of $f(x) > 0$ if for all $c > 0$ we have $z(c) > 0$ and $f(z(c)) > 0$. One easily verifies that $z(c) = \frac{c_1}{c_2}$ is a parametric positive solution of $f(x)$.

However, $g(x) > 0$ does not have any parametric positive solution since $g(x) > 0$ has no positive solution when, e.g, $c_2 = c_1 = c_0 = 1$. Of course, we are interested in tackling much larger cases with respect to numbers of variables, monomials, and polynomials.

The problem is important as systems of polynomial inequalities often arise in science and engineering applications, including, e.g., the qualitative analysis of biological or chemical networks [28, 11, 5, 10] or Satisfiability Modulo Theories (SMT) solving [20, 1, 12]. Surprisingly often one is indeed interested in positive solutions. For instance, unknowns in the biological and chemical context of [28, 11, 5, 10] are typically positive concentrations of species or reaction rates, where the direction of the reaction is known. In SMT solving, positivity is often not required but, in the satisfiable case, benchmarks typically have also positive solutions; comprehensive statistical data for several thousand benchmarks can be found in [12, Sect. 6]. In many areas systems have parameters and one desires to have parametric solutions. Hence, an efficient and reliable tool for finding parametric positive solutions can aid scientists and engineers in developing and investigating their mathematical models.

The problem of finding parametric positive solution is essentially that of quantifier elimination over the first order theory of real closed fields. In 1930, Alfred Tarski [26] showed that real quantifier elimination can be carried out algorithmically. Since then, there have been intensive research, producing profound theories, dramatically improved algorithms, and highly refined implementations in widely available computer algebra software such as Mathematica, Maple, Qepcad B, or Reduce, e.g., [26, 8, 2, 19, 13, 7, 14, 15, 9, 29, 25, 23, 6, 22, 16, 17]. However, existing general quantifier elimination algorithms are still too inefficient for tackling even small problems of finding parametric positive solutions.

The main contribution of this paper is to provide simple and efficient algorithmic criteria for deciding whether or not a given signed parametric system has a parametric positive solution. If so, we provide an explicit formula (rational function) for a parametric positive solution. The main challenge was eliminating many universal quantifiers in the problem statement. We tackled that challenge by, firstly, carefully approximating/bounding polynomials by suitable multiple of monomials and, secondly, tropicalizing, i.e., linearizing monomials by taking logarithms in the style of [27]. However, unlike standard tropicalization approaches, we determine sufficiently large *finite bases* for our logarithms, in order to get an explicit formula for parametric positive solutions.

Our main result also shines a new light on recent heuristic subtropical methods [24, 12]: We provide a precise characterization of their incompleteness in terms of the existence of parametric positive solutions for the originally non-parametric input problems considered there.

The paper is structured as follows. In Section 2, we motivate and present a compact and convenient notation for a systems of multivariate polynomials, which will be used throughout the paper. In Section 3, we precisely define the key notions of *signed parametric systems* and *parametric positive solutions*. Then we present and prove the main result of this paper, which shows how to check the existence of a parametric positive solution and, in the positive case,

how to find one. In Section 4, we apply use our framework and our result to re-analyze and improve the above-mentioned subtropical methods [24, 12].

2 Notation

The principal mathematical object studied in this paper are systems of multivariate polynomials over the real numbers. In order to minimize cumbersome indices, we are going to introduce some compact notations. Let us start with a motivation by means of a simple example.

Example 1. Consider the following system of three polynomials in two variables:

$$\begin{aligned} f_1 &= 2x_1^2x_2 - x_1^3 \\ f_2 &= -3x_1x_2^2 + 6x_1^3 \\ f_3 &= -x_1^2 + 5x_1x_2^2. \end{aligned}$$

We rewrite those polynomials by aligning their signs, coefficients, and monomials supports:

$$\begin{aligned} f_1 &= 1 \cdot 2 \cdot x_1^2x_2^1 + 0 \cdot 1 \cdot x_1^1x_2^2 + -1 \cdot 4 \cdot x_1^3x_2^0 \\ f_2 &= 0 \cdot 1 \cdot x_1^2x_2^1 + -1 \cdot 3 \cdot x_1^1x_2^2 + 1 \cdot 6 \cdot x_1^3x_2^0 \\ f_3 &= -1 \cdot 1 \cdot x_1^2x_2^1 + 1 \cdot 5 \cdot x_1^1x_2^2 + 0 \cdot 1 \cdot x_1^3x_2^0, \end{aligned}$$

where signs are represented by -1 , 0 , and 1 . Note that we are writing 0 coefficients as $0 \cdot 1$ for notational uniformity. Rewriting this in matrix-vector notation, we have

$$\begin{bmatrix} f_1 \\ f_2 \\ f_3 \end{bmatrix} = \left(\left(\begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{bmatrix} \circ \begin{bmatrix} 2 & 1 & 4 \\ 1 & 3 & 6 \\ 1 & 5 & 1 \end{bmatrix} \right) \begin{bmatrix} x_1^2x_2^1 \\ x_1^1x_2^2 \\ x_1^3x_2^0 \end{bmatrix} \right),$$

where \circ is the component-wise Hadamard product. Pushing this even further, we have

$$\begin{bmatrix} f_1 \\ f_2 \\ f_3 \end{bmatrix} = \left(\left(\begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{bmatrix} \circ \begin{bmatrix} 2 & 1 & 4 \\ 1 & 3 & 6 \\ 1 & 5 & 1 \end{bmatrix} \right) \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 3 & 0 \end{bmatrix} \right).$$

Thus we have arrived at a form

$$f = (s \circ c)x^e,$$

where

$$f = \begin{bmatrix} f_1 \\ f_2 \\ f_3 \end{bmatrix}, \quad s = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -1 & 1 \\ -1 & 1 & 0 \end{bmatrix}, \quad c = \begin{bmatrix} 2 & 1 & 4 \\ 1 & 3 & 6 \\ 1 & 5 & 1 \end{bmatrix},$$

$$x = [x_1 \quad x_2], \quad e = \begin{bmatrix} e_1 \\ e_2 \\ e_3 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 1 & 2 \\ 3 & 0 \end{bmatrix}.$$

This concludes our example.

In general, a system $f \in \mathbb{R}[x_1, \dots, x_d]^u$ of multivariate polynomials over the reals will be written compactly as

$$f = (s \circ c)x^e,$$

where

$$f = \begin{bmatrix} f_1 \\ \vdots \\ f_u \end{bmatrix}, \quad s = \begin{bmatrix} s_{11} & \cdots & s_{1v} \\ \vdots & & \vdots \\ s_{u1} & \cdots & s_{uv} \end{bmatrix}, \quad c = \begin{bmatrix} c_{11} & \cdots & c_{1v} \\ \vdots & & \vdots \\ c_{u1} & \cdots & c_{uv} \end{bmatrix},$$

$$x = [x_1 \quad \cdots \quad x_d], \quad e = \begin{bmatrix} e_1 \\ \vdots \\ e_v \end{bmatrix} = \begin{bmatrix} e_{11} & \cdots & e_{1d} \\ \vdots & & \vdots \\ e_{v1} & \cdots & e_{vd} \end{bmatrix}.$$

We call $s \in \{-1, 0, 1\}^{u \times v}$ the *sign matrix*, $c \in \mathbb{R}_+^{u \times v}$ the *coefficient matrix*, and $e \in \mathbb{N}^{v \times d}$ the *exponent matrix* of f .

3 Main Result

Definition 2 (Signed parametric systems). A *signed parametric system* is given by

$$f = (s \circ c)x^e,$$

where the sign matrix $s \in \{-1, 0, 1\}^{u \times v}$ and the exponent matrix $e \in \mathbb{N}^{v \times d}$ are specified but the coefficient matrix c is unspecified in the sense that it is left parametric. Formally c is a $u \times v$ -matrix of pairwise different indeterminates.

Definition 3 (Parametric positive solutions). Consider a signed parametric system $f = (s \circ c)x^e$. A *parametric positive solution* of $f(x) > 0$ is a function $z : \mathbb{R}_+^{u \times v} \rightarrow \mathbb{R}_+^d$ that maps each possible specification of the coefficient matrix c to a solution of the corresponding non-parametric system, i.e.,

$$\forall_{c>0} f(z(c)) > 0.$$

Theorem 4 (Main). *Let $f = (s \circ c)x^e$ be a signed parametric system. Let*

$$C(n) := \bigwedge_i \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot n \geq 1.$$

Then the following are equivalent:

(i) $f(x) > 0$ has a parametric positive solution.

(ii) $C(n)$ has a solution $n \in \mathbb{R}^d$.

(iii) $C(n)$ has a solution $n \in \mathbb{Z}^d$.

In the positive case, the following function z is a parametric positive solution of $f(x) > 0$:

$$z(c) = t^n, \quad \text{where } t = 1 + \sum_{\substack{s_{ij} > 0 \\ s_{ik} < 0}} \frac{c_{ik}}{c_{ij}}.$$

In fact, we even have $\forall_{c > 0} \forall_{r \geq t} f(r^n) > 0$.

Proof. We first show that (i) implies (ii):

$$\begin{aligned} \text{(i)} &\iff \forall_{c > 0} \exists_{x > 0} (s \circ c)x^e > 0 \\ &\iff \forall_{c > 0} \exists_{x > 0} \bigwedge_i \sum_{s_{ij} > 0} c_{ij} x^{e_j} > \sum_{s_{ik} < 0} c_{ik} x^{e_k} \\ &\implies \exists_{x > 0} \bigwedge_i \sum_{s_{ij} > 0} x^{e_j} > \sum_{s_{ik} < 0} 2v x^{e_k}, \quad \text{by instantiating } c \\ &\implies \exists_{x > 0} \bigwedge_i v \max_{s_{ij} > 0} x^{e_j} > \max_{s_{ik} < 0} 2v x^{e_k} \\ &\iff \exists_{x > 0} \bigwedge_i \max_{s_{ij} > 0} x^{e_j} > \max_{s_{ik} < 0} 2x^{e_k} \\ &\iff \exists_{x > 0} \bigwedge_i \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} x^{e_j} > 2x^{e_k} \\ &\iff \exists_{x > 0} \bigwedge_i \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} x^{e_j - e_k} > 2 \\ &\iff \exists_{x > 0} \bigwedge_i \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot \log_2 x > 1 \\ &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_i \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot n > 1, \quad \text{using } \log_2 : \mathbb{R}_+ \leftrightarrow \mathbb{R} \\ &\implies \text{(ii)}. \end{aligned}$$

Assume now that (ii) holds. The existence of solutions $n \in \mathbb{R}^d$ and $n \in \mathbb{Q}^d$ of $C(n)$ coincides due to the Linear Tarski Principle: Ordered fields admit quantifier elimination for linear formulas, and therefore \mathbb{Q} is an elementary

substructure of \mathbb{R} with respect to linear sentences [18]. Given a solution $n \in \mathbb{Q}^d$, we can use the principal denominator $\delta > 0$ of all coordinates of n to obtain a solution $\delta n \in \mathbb{Z}^d$. Hence (iii) holds.

We finally show that (iii) implies (i):

$$\begin{aligned}
(i) &\iff \forall_{c>0} \exists_{x>0} (s \circ c)x^e > 0 \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \sum_{s_{ij}>0} c_{ij}x^{e_j} > \sum_{s_{ik}<0} c_{ik}x^{e_k} \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \max_{s_{ij}>0} c_{ij}x^{e_j} > \bar{c}_i \max_{s_{ik}<0} x^{e_k}, \quad \text{where } \bar{c}_i = \sum_{s_{ik'}<0} c_{ik'} \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} c_{ij}x^{e_j} > \bar{c}_i x^{e_k} \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} x^{e_j-e_k} > \frac{\bar{c}_i}{c_{ij}} \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} x^{e_j-e_k} \geq t, \\
&\quad \text{where } t \text{ is as stated in the theorem} \\
&\iff \forall_{c>0} \exists_{x>0} \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} (e_j - e_k) \cdot \log_t x \geq 1, \\
&\iff \exists_n \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} (e_j - e_k) \cdot n \geq 1, \quad \text{using } \log_t : \mathbb{R}_+ \leftrightarrow \mathbb{R} \\
&\iff \exists_{n \in \mathbb{Z}^d} \bigwedge_i \bigwedge_{s_{ik}<0} \bigvee_{s_{ij}>0} (e_j - e_k) \cdot n \geq 1 \\
&\iff (iii).
\end{aligned}$$

In our proof of the implication from (iii) to (i) we have applied \log_t so that $n = \log_t x$ and, accordingly, $x = t^n$, where t is as stated in the theorem. Notice that any larger choice $r \geq t$ would work there as well. \square

Example 5. Consider $f = \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}$ with

$$\begin{aligned}
f_1 &= -c_{11}x_1^5 + c_{12}x_1^2x_2 - c_{13}x_1^2 + c_{15}x_2^2 \\
f_2 &= c_{21}x_1^5 + c_{22}x_1^2x_2 + c_{23}x_1^2 - c_{24}x_2^3.
\end{aligned}$$

That is

$$s = \begin{bmatrix} -1 & 1 & -1 & 0 & 1 \\ 1 & 1 & 1 & -1 & 0 \end{bmatrix}, \quad e = \begin{bmatrix} 5 & 0 \\ 2 & 1 \\ 2 & 0 \\ 0 & 3 \\ 0 & 2 \end{bmatrix}.$$

Then $C(n)$ has a solution $n \in \mathbb{Z}^2$, e.g.,

$$n = \begin{bmatrix} -12 & -11 \end{bmatrix}.$$

Hence $f = (s \circ c)x^e > 0$ has a parametric positive solution, e.g.,

$$z(c) = \begin{bmatrix} t^{-12} & t^{-11} \end{bmatrix},$$

where $t = 1 + \frac{c_{11}}{c_{12}} + \frac{c_{11}}{c_{15}} + \frac{c_{13}}{c_{12}} + \frac{c_{13}}{c_{15}} + \frac{c_{24}}{c_{21}} + \frac{c_{24}}{c_{22}} + \frac{c_{24}}{c_{23}}$.

Example 6. We slightly modify Example 5 and consider $f = \begin{bmatrix} f_1 \\ f_2 \end{bmatrix}$ with

$$\begin{aligned} f_1 &= -c_{11}x_1^5 + c_{12}x_1^2x_2 - c_{13}x_1^2 + c_{15}x_2^2 \\ f_2 &= c_{21}x_1^5 + c_{22}x_1^2x_2 - c_{23}x_1^2 - c_{24}x_2^3. \end{aligned}$$

That is

$$s = \begin{bmatrix} -1 & 1 & -1 & 0 & 1 \\ 1 & 1 & -1 & -1 & 0 \end{bmatrix}, \quad e = \begin{bmatrix} 5 & 0 \\ 2 & 1 \\ 2 & 0 \\ 0 & 3 \\ 0 & 2 \end{bmatrix}.$$

Then $C(n)$ does not have a solution $n \in \mathbb{Z}^2$. Hence $f = (s \circ c)x^e > 0$ does not have a parametric positive solution.

4 A Re-analysis of Subtropical Methods

For non-parametric systems of real polynomial inequalities, heuristic Newton polytope-based *subtropical methods* [24, 12] have been successfully applied in two quite different areas: Firstly, qualitative analysis of biological and chemical networks and, secondly, SMT solving.

In the first area, a positive solution of a very large single inequality could be computed. The left hand side polynomial there has more than $8 \cdot 10^5$ monomials in 10 variables with individual degrees up to 10. This computation was the hard step in finding an exact positive solution of the corresponding equation using a known positive point with negative value of the polynomial and applying the intermediate value theorem. To give a very rough idea of the biological background: The polynomial is a Hurwitz determinant originating from a system of ordinary differential equations modeling mitogen-activated protein kinase (MAPK) in the metabolism of a frog. Positive zeros of the Hurwitz determinant point at Hopf bifurcations, which are in turn indicators for possible oscillation of the corresponding reaction network. For further details see [11].

In the second area, a subtropical approach for systems of several polynomial inequalities has been integrated with an SMT solver. That combination could solve a surprisingly large percentage of SMT benchmarks very fast and thus

establishes an interesting heuristic preprocessing step for the SMT theory of QF_NRA (quantifier-free nonlinear arithmetic). For detailed statistics see [12, Sect. 6].

The goal of this section is, to make precise the connections between subtropical methods and our main result here, to use these connections to improve the subtropical methods, and to precisely characterize their incompleteness.

4.1 Subtropical Real Root Finding

In [24] we have studied an incomplete method for heuristically finding a positive solution for a single multivariate polynomial inequality with fixed integer coefficients:

$$[f_1] = (s \circ c)x^e \quad \text{where } s \in \{-1, 0, 1\}^{1 \times v}, \quad c \in \mathbb{Z}_+^{1 \times v}, \quad e \in \mathbb{N}^{v \times d}.$$

The method considers the positive and the negative support, which in terms of our notions is given by

$$S^+ = \{e_j \mid s_{1j} > 0\}, \quad S^- = \{e_k \mid s_{1k} < 0\}.$$

Then [24, Lemma 4] essentially states that $f_1(x) > 0$ has a positive solution if

$$C' := \bigvee_{e_j \in S^+} \bigvee_{n \in \mathbb{R}^d} \bigvee_{\gamma \in \mathbb{R}} \left(\begin{bmatrix} -e_j & 1 \end{bmatrix} \begin{bmatrix} n \\ \gamma \end{bmatrix} \leq -1 \wedge \bigwedge_{\substack{e_k \in S^+ \cup S^- \\ e_k \neq e_j}} \begin{bmatrix} e_k & -1 \end{bmatrix} \begin{bmatrix} n \\ \gamma \end{bmatrix} \leq -1 \right).$$

Unfortunately, in [24, Lemma 4] vectors $e_l = [0 \ \dots \ 0]$ corresponding to absolute summands are treated specially. We have noted already in [12, p.192] that an inspection of the proof shows that this is not necessary. Therefore we discuss here a slightly improved and simpler version without that special treatment, which has been explicitly stated as [12, Lemma 2].

The proof of the loop invariant (I₁) in [24, Theorem 5(ii)] shows that the positive support need not be considered in the conjunction:

$$C' \iff \bigvee_{e_j \in S^+} \bigvee_{n \in \mathbb{R}^d} \bigvee_{\gamma \in \mathbb{R}} \left(\begin{bmatrix} -e_j & 1 \end{bmatrix} \begin{bmatrix} n \\ \gamma \end{bmatrix} \leq 1 \wedge \bigwedge_{e_k \in S^-} \begin{bmatrix} e_k & -1 \end{bmatrix} \begin{bmatrix} n \\ \gamma \end{bmatrix} \leq -1 \right).$$

Starting with Fourier–Motzkin elimination [21, Sect. 12.2] of γ , we obtain

$$C' \iff \bigvee_{e_j \in S^+} \bigvee_{n \in \mathbb{R}^d} \bigwedge_{e_k \in S^-} (e_k - e_j) \cdot n \leq -2$$

$$\begin{aligned}
&\iff \bigvee_{e_j \in S^+} \exists_{n \in \mathbb{R}^d} \bigwedge_{e_k \in S^-} (e_j - e_k) \cdot n \geq 1 \\
&\iff \exists_{n \in \mathbb{R}^d} \bigvee_{e_j \in S^+} \bigwedge_{e_k \in S^-} (e_j - e_k) \cdot n \geq 1 \\
&\iff \exists_{n \in \mathbb{R}^d} \max_{e_j \in S^+} (e_j \cdot n) \geq \max_{e_k \in S^-} (e_k \cdot n + 1) \\
&\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{e_k \in S^-} \bigvee_{e_j \in S^+} (e_j - e_k) \cdot n \geq 1 \\
&\iff \exists_{n \in \mathbb{R}^d} C(n)
\end{aligned}$$

with $C(n)$ as in Theorem 4.

Corollary 7. Consider $f_1 \in \mathbb{Z}[x_1, \dots, x_d]$, say, $f_1 = (s \circ c)x^e$, where $s \in \{-1, 0, 1\}^{1 \times v}$, $c \in \mathbb{Z}_+^{1 \times v}$, $e \in \mathbb{N}^{v \times d}$. Let $f^* = [(s \circ c^*)x^e]$, where c^* is a $1 \times v$ -matrix of pairwise different indeterminates. Then the following are equivalent:

- (i) The algorithm **find-positive** [24, Algorithm 1] does not fail, and thus finds a rational solution of $f_1 > 0$ with positive coordinates.
- (ii) There is a row e_j of e with $s_{1j} > 0$ such that the following LP problem has a solution $n \in \mathbb{Q}^d$:

$$\bigwedge_{s_{1k} < 0} (e_j - e_k) \cdot n \geq 1.$$

- (iii) $f^* > 0$ has a parametric positive solution.

In the positive case, $f(r^n) > 0$ for all $r \geq 1 + v \sum_{s_{1k} < 0} c_{1k}$.

Proof. The equivalence between (i), (ii), and (iii) has been derived above.

According to Theorem 4, a solution for $f_1 > 0$ can be obtained by plugging c into the parametric positive solution for f^* . Since we have positive integer coefficients, we can bound t from above as follows.

$$t = 1 + \sum_{\substack{s_{1j} > 0 \\ s_{1k} < 0}} \frac{c_{1k}}{c_{1j}} \leq 1 + \sum_{\substack{s_{1j} > 0 \\ s_{1k} < 0}} \frac{c_{1k}}{1} \leq 1 + v \sum_{s_{1k} < 0} c_{1k}. \quad \square$$

In simple words the equivalence between (i) and (iii) in the corollary states the following: The incomplete heuristic [24, Algorithm 1] succeeds *if and only if* not only the inequality for the input polynomial has a solution as required, but also the inequality for all polynomials with the same monomials and signs of coefficients as the input polynomial.

We have added (ii) to the corollary, because we consider this form optimal for algorithmic purposes. Our special case of one single inequality allows to transform the conjunctive normal form provided by Theorem 4 into an equivalent disjunctive normal form without increasing size. This way, a decision procedure

can use finitely many LP solving steps [21] instead of employing more general methods like SMT solving [20].

Finally notice that the brute force search for a suitable t in **find-positive** [24, Algorithm 1, l.10–12] is not necessary anymore. Our corollary computes a suitable number from the coefficients.

4.2 Subtropical Satisfiability Checking

Subsequent work [12] takes an entirely geometric approach to generalize the work in [24] from one polynomial inequality to finitely many such inequalities. Consider a system with fixed integer coefficients in our notation:

$$f = \begin{bmatrix} f_1 \\ \vdots \\ f_u \end{bmatrix} = (s \circ c)x^e, \quad \text{where } s \in \{-1, 0, 1\}^{u \times v}, \quad c \in \mathbb{Z}_+^{u \times v}, \quad e \in \mathbb{N}^{v \times d}.$$

Then [12, Theorem 12] derives the following sufficient condition for the existence of a positive solution of $f > 0$:

$$C'' := \exists_{n \in \mathbb{R}^d} \exists_{\gamma_1 \in \mathbb{R}} \dots \exists_{\gamma_u \in \mathbb{R}} \bigwedge_{i=1}^u \left(\left(\bigvee_{s_{ij} > 0} ne_j + \gamma_i > 0 \right) \wedge \bigwedge_{s_{ik} < 0} ne_k + \gamma_i < 0 \right).$$

After an equivalence transformation, we can once more apply Fourier–Motzkin elimination [21, Sect. 12.2]:

$$\begin{aligned} C'' &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{i=1}^u \bigvee_{s_{ij} > 0} \exists_{\gamma_i \in \mathbb{R}} \left(ne_j + \gamma_i > 0 \wedge \bigwedge_{s_{ik} < 0} ne_k + \gamma_i < 0 \right) \\ &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{i=1}^u \bigvee_{s_{ij} > 0} \bigwedge_{s_{ik} < 0} (e_j - e_k) \cdot n > 0 \\ &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{i=1}^u \max_{s_{ij} > 0} e_j n > \max_{s_{ik} < 0} e_k n \\ &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{i=1}^u \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot n > 0 \\ &\iff \exists_{n \in \mathbb{R}^d} \bigwedge_{i=1}^u \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot n \geq 1 \\ &\iff \exists_{n \in \mathbb{R}^d} C(n) \end{aligned}$$

with $C(n)$ as in Theorem 4.

Corollary 8. *Consider $f \in \mathbb{Z}[x_1, \dots, x_d]^u$, say, $f = (s \circ c)x^e$, where $s \in \{-1, 0, 1\}^{u \times v}$, $c \in \mathbb{Z}_+^{u \times v}$, $e \in \mathbb{N}^{v \times d}$. Let $f^* = (s \circ c^*)x^e$, where c^* is a $u \times v$ -matrix of pairwise different indeterminates. Then the following are equivalent:*

- (i) The incomplete subtropical satisfiability checking method for several inequalities over QF_NRA (quantifier-free nonlinear real arithmetic) introduced in [12] succeeds on $f > 0$.
- (ii) The following SMT problem with unknowns n is satisfiable in QF_LRA (quantifier-free linear real arithmetic):

$$\bigwedge_{i=1}^u \bigwedge_{s_{ik} < 0} \bigvee_{s_{ij} > 0} (e_j - e_k) \cdot n \geq 1.$$

- (iii) $f^* > 0$ has a parametric positive solution.

In the positive case, $f(r^n) > 0$ for all $r \geq 1 + v \sum_{s_{ik} < 0} c_{ik}$.

Proof. The equivalence between (i), (ii), and (iii) has been derived above. About the solution r see the proof of Corollary 7. \square

The equivalence between (i) and (iii) in the corollary states the following: The procedure in [12] yields “sat” in contrast to “unknown” *if and only if* not only the input system is satisfiable, but that system with all real choices of coefficients with the same signs as in the input system. While there are no formal algorithms in [12], the work has been implemented within a combination of the veriT solver [4] with the library STROPSAT [12, Sect. 6]. Our characterization applies in particular to the completeness of this software.

We have added (ii) to the corollary, because we consider this form optimal for algorithmic purposes. Like the original input C'' used in [12] this is a conjunctive normal form, which is ideal for DPLL-based SMT solvers [20]. Recall that u is the number of inequalities in the input, and d is the number of variables. Let ι and κ be the numbers of positive and negative coefficients, respectively. Then compared to [12] we have reduced $d + u$ variables to d variables, and we have reduced $u\kappa$ clauses with ι atoms each plus u unit clauses to some different $u\kappa$ clauses with ι atoms each but without any additional unit clauses.

With the `:produce-model` option the SMT-LIB standard [3] supports solutions like the r^n provided by our corollary. The work in [12] does not address the computation of solutions. It only mentions that sufficiently large r will work, which implicitly suggests a brute-force search like the one in [24, Algorithm 1, 1.10–12].

Acknowledgments

This work has been supported by the European Union’s Horizon 2020 research and innovation programme under grant agreement No H2020-FETOPEN-2015-CSA 712689 SC-SQUARE and by the bilateral project ANR-17-CE40-0036 and DFG-391322026 SYMBIONT. The second author would like to thank Georg Regensburger for his hospitality and an interesting week of inspiring discussions around the topic, and Dima Grigoriev for getting him started on the subject.

References

- [1] E. Abraham, J. Abbott, B. Becker, A. M. Bigatti, M. Brain, B. Buchberger, A. Cimatti, J. H. Davenport, M. England, P. Fontaine, S. Forrest, A. Griggio, D. Kroening, W. M. Seiler, and T. Sturm. *SC²: Satisfiability checking meets symbolic computation*. In *CICM 2016*, volume 9791 of *LNCS*, pages 28–43. Springer, 2016.
- [2] D. S. Arnon. Algorithms for the geometry of semi-algebraic sets. Technical Report 436, Comput. Sci. Dept., University of Wisconsin-Madison, 1981. Ph.D. Thesis.
- [3] C. Barrett, P. Fontaine, and C. Tinelli. The SMT-LIB standard: Version 2.6. Technical report, Department of Computer Science, The University of Iowa, 2017.
- [4] T. Bouton, D. Caminha B. De Oliveira, D. Déharbe, and P. Fontaine. veriT: An open, trustable and efficient SMT-solver. In *CADE-22*, volume 5663 of *LNCS*, pages 151–156. Springer, 2009.
- [5] R. Bradford, J. Davenport, M. England, H. Errami, V. Gerdt, D. Grigoriev, C. Hoyt, M. Kosta, O. Radulescu, T. Sturm, and A. Weber. A case study on the parametric occurrence of multiple steady states. In *ISSAC 2017*, pages 45–52. ACM, 2017.
- [6] C. W. Brown. Improved projection for CAD’s of \mathbb{R}^3 . In *ISSAC 2000*, pages 48–53. ACM, 2000.
- [7] J. Canny. Some algebraic and geometric computations in PSPACE. In *Proceedings of the 20th Annual ACM Symposium on the Theory of Computing*, pages 460–467, 1988.
- [8] G. E. Collins. Quantifier elimination for the elementary theory of real closed fields by cylindrical algebraic decomposition. In *Automata Theory and Formal Languages. 2nd GI Conference*, volume 33 of *LNCS*, pages 134–183. Springer, 1975.
- [9] G. E. Collins and H. Hong. Partial cylindrical algebraic decomposition for quantifier elimination. *J. Symb. Comput.*, 12(3):299–328, 1991.
- [10] M. England, H. Errami, D. Grigoriev, O. Radulescu, T. Sturm, and A. Weber. Symbolic versus numerical computation and visualization of parameter regions for multistationarity of biological networks. In *CASC 2017*, volume 10490 of *LNCS*, pages 93–108. Springer, 2017.
- [11] H. Errami, M. Eiswirth, D. Grigoriev, W. M. Seiler, T. Sturm, and A. Weber. Detection of Hopf bifurcations in chemical reaction networks using convex coordinates. *J. Comput. Phys.*, 291:279–302, 2015.

- [12] P. Fontaine, M. Ogawa, T. Sturm, and X. T. Vu. Subtropical satisfiability. In *FROCOS 2017*, volume 10483 of *LNCS*, pages 189–206. Springer, 2017.
- [13] D. Grigoriev and N. Vorobjov. Solving systems of polynomial inequalities in subexponential time. *J. Symb. Comput.*, 5(1/2):37–64, 1988.
- [14] H. Hong. An improvement of the projection operator in cylindrical algebraic decomposition. In *ISSAC 1990*, pages 261–264. ACM, 1990.
- [15] H. Hong. *Improvements in CAD-based Quantifier Elimination*. PhD thesis, The Ohio State University, 1990.
- [16] H. Hong and M. S. E. Din. Variant quantifier elimination. *J. Symb. Comput.*, 47(7):883–901, 2012.
- [17] M. Košta. *New Concepts for Real Quantifier Elimination by Virtual Substitution*. Doctoral dissertation, Saarland University, Germany, 2016.
- [18] R. Loos and V. Weispfenning. Applying linear quantifier elimination. *THE Computer Journal*, 36(5):450–462, 1993.
- [19] S. McCallum. *An Improved Projection Operator for Cylindrical Algebraic Decomposition*. PhD thesis, University of Wisconsin-Madison, 1984.
- [20] R. Nieuwenhuis, A. Oliveras, and C. Tinelli. Solving SAT and SAT modulo theories: From an abstract Davis–Putnam–Logemann–Loveland procedure to DPLL(T). *JACM*, 53(6):937–977, 2006.
- [21] A. Schrijver. *Theory of Linear and Integer Programming*. John Wiley & Sons, Chichester, 1986.
- [22] A. Strzebonski. Cylindrical algebraic decomposition using validated numerics. *J. Symb. Comput.*, 41(9):1021–1038, 2006.
- [23] T. Sturm. *Real Quantifier Elimination in Geometry*. Doctoral dissertation, Universität Passau, Germany, 1999.
- [24] T. Sturm. Subtropical real root finding. In *ISSAC 2015*, pages 347–354. ACM, 2015.
- [25] T. Sturm and V. Weispfenning. Computational geometry problems in Redlog. In *Automated Deduction in Geometry*, volume 1360 of *LNAI*, pages 58–86. Springer, 1998.
- [26] A. Tarski. The completeness of elementary algebra and geometry. Reprinted by CNRS, Institute Blaise Pascal, Paris, 1967, 1930.
- [27] O. Viro. Dequantization of real algebraic geometry on logarithmic paper. *CoRR arXiv:math/0005163*, 2000.
- [28] A. Weber, T. Sturm, and E. O. Abdel-Rahman. Algorithmic global criteria for excluding oscillations. *Bull. Math. Biol.*, 73(4):899–916, 2011.

- [29] V. Weispfenning. A new approach to quantifier elimination for real algebra. In B. Caviness and J. Johnson, editors, *Quantifier Elimination and Cylindrical Algebraic Decomposition*. Springer, 1996.