

E_9 exceptional field theory. Part I. The potential

Guillaume Bossard,^a Franz Ciceri,^b Gianluca Inverso,^c Axel Kleinschmidt^{b,d} and Henning Samtleben^e

^a*Centre de Physique Théorique, Ecole Polytechnique, CNRS,
Université Paris-Saclay,
FR-91128 Palaiseau cedex, France*

^b*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut),
Am Mühlenberg 1, DE-14476 Potsdam, Germany*

^c*Centre for Research in String Theory, School of Physics and Astronomy,
Queen Mary University of London,
327 Mile End Road, London, E1 4NS, United Kingdom*

^d*International Solvay Institutes,
ULB-Campus Plaine CP231, BE-1050 Brussels, Belgium*

^e*Univ Lyon, Ens de Lyon, Univ Claude Bernard, CNRS,
Laboratoire de Physique, FR-69342 Lyon, France*

E-mail: guillaume.bossard@polytechnique.edu, franz.ciceri@aei.mpg.de,
g.inverso@qmul.ac.uk, axel.kleinschmidt@aei.mpg.de,
henning.samtleben@ens-lyon.fr

ABSTRACT: We construct the scalar potential for the exceptional field theory based on the affine symmetry group E_9 . The fields appearing in this potential live formally on an infinite-dimensional extended spacetime and transform under E_9 generalised diffeomorphisms. In addition to the scalar fields expected from $D = 2$ maximal supergravity, the invariance of the potential requires the introduction of new constrained scalar fields. Other essential ingredients in the construction include the Virasoro algebra and indecomposable representations of E_9 . Upon solving the section constraint, the potential reproduces the dynamics of either eleven-dimensional or type IIB supergravity in the presence of two isometries.

KEYWORDS: Extended Supersymmetry, String Duality, Supergravity Models

ARXIV EPRINT: [1811.04088](https://arxiv.org/abs/1811.04088)

Contents

1	Introduction	1
2	E_9 basic representation and Virasoro algebra	4
2.1	E_9 preliminaries	5
2.2	Virasoro algebra	6
3	Building blocks of E_9 exceptional field theory	9
3.1	$D = 2$ maximal supergravity fields	9
3.2	Generalised diffeomorphisms and scalar fields	12
3.2.1	Generalised Lie derivative	12
3.2.2	Unconstrained scalar fields and currents	14
3.2.3	Constrained scalar fields and shifted currents	16
4	The potential	16
4.1	The potential at $\tilde{\rho} = 0$	17
4.1.1	Rigid E_9 symmetry	18
4.1.2	Invariance under generalised diffeomorphisms	19
4.2	The potential at $\tilde{\rho} \neq 0$	23
4.2.1	Rigid $\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L-1})$ symmetry	23
4.2.2	Invariance under generalised diffeomorphisms	27
4.2.3	Invariance under Σ transformations	31
4.3	The potential in the unendlichbein formalism	34
5	Reduction to E_8 and consistency with supergravity	37
5.1	E_8 section and exceptional field theory	37
5.2	Parameterising \mathcal{M} and decomposition of the potential	39
5.3	Comparison with E_8 exceptional theory	45
6	Conclusions	46
A	Properties of the cocycle	48
A.1	E_9 group 1-cocycles in the co-adjoint	48
A.2	Generalisation to $\hat{E}_8 \times \text{SL}(2)$	51
A.3	Useful identities	53
B	Gauge algebra closure on $\langle \chi$	55

1 Introduction

Exceptional geometry is a way of unifying the local symmetries of supergravity theories by combining geometric diffeomorphisms with matter gauge transformations into a single so-called generalised Lie derivative [1–14]. This generalised Lie derivative generates generalised diffeomorphisms acting on the fields of the theory and requires the introduction of an extended space beyond the usual space-time geometry of gravity. The generalised Lie derivative forms a closed gauge algebra only when the so-called section condition is imposed on the fields, restricting their dependence on the extended space. Upon solving the section constraint explicitly one recovers the standard supergravity theories. However, exceptional geometry also offers the possibility of describing more complicated (local or global) situations that have been named non-geometric backgrounds.

The symmetry groups of maximal supergravities in $D = 11 - n$ dimensions belong to the (split real) exceptional series E_n [15–19]. For each exceptional symmetry group E_n one can construct an extended space that is described locally by a set of coordinates Y^M where M labels a representation of E_n [7]. Adjoining to these coordinates the ‘external’ D -dimensional space with coordinates x^μ and an appropriate notion of external diffeomorphisms one obtains a total space with coordinates (x^μ, Y^M) . Taking the fields from D -dimensional maximal supergravity, as prescribed by the tensor hierarchy [20, 21], one may try to construct an action invariant under generalised and external diffeomorphisms. As it turns out, closure of the gauge algebra of p -forms and invariance of the action require the introduction of additional p -forms of rank $p \geq D - 2$ beyond those of D -dimensional supergravity. These extra p -forms are covariantly constrained in the sense that they obey algebraic constraints analogous to those satisfied by the internal partial derivatives $\partial_M = \frac{\partial}{\partial Y^M}$ by virtue of the section constraint. Moreover, these fields do not constitute additional degrees of freedom, but are related by first order equations to the propagating fields of the theory. Combining all these ingredients leads to a unique theory called E_n exceptional field theory and that has been explicitly constructed for E_n with $n \leq 8$ [22–24]. Imposing a solution to the section condition relates exceptional field theory to maximal supergravity in eleven space-time dimensions, to type IIB supergravity or to their dimensional reductions depending on the choice of solution to the section condition.

The invariant ‘actions’ of exceptional field theories combine various terms. They carry an Einstein-Hilbert-type term, kinetic terms for the various matter fields including a non-linear sigma model for the scalars, a topological term for the p -forms, and a ‘potential’ term for the scalar fields. The scalar fields belong to the coset space $E_n/K(E_n)$, where $K(E_n)$ denotes the maximal compact subgroup of E_n , and may be parameterised by a symmetric matrix \mathcal{M}_{MN} which determines the internal generalised metric on the extended space. The ‘potential’ $V(\mathcal{M})$ is bilinear in the internal derivatives ∂_M with respect to the extended coordinates but does not carry derivatives ∂_μ with respect to the ‘external’ coordinates. Under generalised Scherk-Schwarz reduction [25–29] in the extended space, it is $V(\mathcal{M})$ that generates the scalar potential term of gauged supergravity. From the point of view of exceptional geometry, $V(\mathcal{M})$ is invariant under generalised diffeomorphisms up to a total derivative and plays to some extent the role of the curvature scalar on the extended space.

Its structure has been worked out up to and including E_8 and it is tied to the remaining terms in the exceptional field theory Lagrangian by external diffeomorphisms.

The first infinite-dimensional group in the E_n series is the affine symmetry group E_9 that is known to be a rigid symmetry of $D = 2$ ungauged maximal supergravity [30]. The associated exceptional field theory has not yet been constructed and the aim of the present paper is to begin filling this gap. The E_9 generalised Lie derivative was recently introduced in [13] and it acts on fields that depend on infinitely many coordinates Y^M . The latter transform under E_9 in the basic lowest weight representation. Closure of the algebra requires a section constraint of the generic form

$$Y^{MN}{}_{PQ} \partial_M \otimes \partial_N = 0 \tag{1.1}$$

with the internal derivatives acting on any pair of fields. Here, $Y^{MN}{}_{PQ}$ is a specific E_9 invariant tensor that can be expressed most easily in terms of quadratic combinations of the E_9 generators as we shall review in section 3.2.1.

The purpose of the present article is to construct the potential V for E_9 exceptional field theory. In $D = 2$, the scalar fields appearing in maximal supergravity parameterise the coset space

$$\frac{\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})}{K(E_9)}, \tag{1.2}$$

where \hat{E}_8 denotes the centrally extended loop group over E_8 . Its quotient by the maximal ‘compact’ subgroup¹ $K(E_9) = K(\hat{E}_8)$ contains the infinite tower of dual scalar fields from $D = 2$ maximal supergravity together with the conformal factor of the external metric. The factor $\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}$ is parameterised by two more scalar fields $\{\rho, \tilde{\rho}\}$ which, in $D = 2$ ungauged supergravity, are related by a free duality equation. The generator d associated with the dilaton ρ enhances \hat{E}_8 to

$$E_9 = \hat{E}_8 \times \mathbb{R}_d^+, \tag{1.3}$$

while the (Virasoro) generator L_{-1} associated with the axion $\tilde{\rho}$ acts as a translation generator on the loop parameter of the loop group \hat{E}_8 . The generator L_{-1} also appears in the E_9 generalised Lie derivative [13].

A novel feature of E_9 exceptional field theory, compared to E_n for $n \leq 8$, is that the scalar sector (1.2) of maximal supergravity is not sufficient to define the theory. This can be seen by extrapolating the generic field content of exceptional field theories down to two external dimensions and noting that the covariantly constrained additional p -forms mentioned above already start from $p = 0$ forms for $D = 2$ external dimensions. Therefore one has to enhance the scalar sector (1.2) by additional constrained fields χ_M transforming under E_9 and generalised diffeomorphisms. Specifically, these additional scalar fields obey the constraints

$$Y^{MN}{}_{PQ} \chi_M \otimes \partial_N = Y^{MN}{}_{PQ} \chi_M \otimes \chi_N = 0, \tag{1.4}$$

¹More precisely, $K(E_9)$ is the maximal E_9 subgroup acting unitarily on the representation of the extended space coordinates. We shall henceforth refer to it as the maximal unitary subgroup.

with the tensor $Y^{MN}{}_{PQ}$ from (1.1), and similar relations with other constrained objects. For the M-theory solution of the section constraint which makes all fields independent of all but nine of the extended directions Y^M , the field χ_M thus has at most nine truly independent components.

As is already the case for E_8 [24], the closure of the algebra of generalised diffeomorphisms requires to not only consider the generalised Lie derivative along a gauge parameter Λ^M in the basic representation, but to also include an additional set of gauge transformations with a covariantly constrained parameter $\Sigma^M{}_N$ [13]. The Lie derivative along the generalised vector Λ^M only involves E_9 , while the constrained parameter also induces a gauging of the \mathbb{R}_{L-1} symmetry when $\Sigma^M{}_M \neq 0$.

Our construction of the E_9 potential is guided by the following requirements. Firstly, the different terms must transform as scalar densities under rigid E_9 and rigid \mathbb{R}_{L-1} transformations. Secondly, the combination of these terms must be such that the potential is invariant under generalised diffeomorphisms, up to a total derivative. Finally, the potential must reproduce parts of the known Lagrangian of E_8 exceptional field theory upon truncation to a suitable subset of coordinates. These requirements allow us to uniquely pinpoint the E_9 potential.

The \mathbb{R}_{L-1} gauge transformations mentioned above can be gauge-fixed by setting $\tilde{\rho} = 0$ without breaking the rest of the generalised diffeomorphisms (i.e. those satisfying $\Sigma^M{}_M = 0$). This choice also breaks the rigid \mathbb{R}_{L-1} invariance. For simplicity, we now present the resulting potential for $\tilde{\rho} = 0$ such that \mathcal{M}_{MN} is valued in E_9 ,

$$V(\mathcal{M}, \chi) = \frac{1}{4} \rho^{-1} \eta^{\alpha\beta} \mathcal{M}^{MN} \mathcal{J}_{M,\alpha} \mathcal{J}_{N,\beta} - \frac{1}{2} \rho^{-1} \mathcal{M}^{PQ} (\mathcal{J}_M)^N{}_P (\mathcal{J}_N)^M{}_Q + \frac{1}{2} \rho \mathcal{M}^{PQ} (\mathcal{J}_M^-)^N{}_P (\mathcal{J}_N^-)^M{}_Q + \rho^{-2} \partial_M \rho \partial_N \mathcal{M}^{MN}. \quad (1.5)$$

The scalar current \mathcal{J}_M is defined as

$$\mathcal{M}^{PS} \partial_M \mathcal{M}_{SQ} = (\mathcal{J}_M)^P{}_Q = \mathcal{J}_{M,\alpha} (T^\alpha)^P{}_Q, \quad (1.6)$$

in terms of the generators T^α of \mathfrak{e}_9 written in the representation $R(\Lambda_0)_0$ that we define below. The (inverse) invariant bilinear form on \mathfrak{e}_9 is denoted $\eta^{\alpha\beta}$, and the *shifted* current \mathcal{J}_M^- is defined as

$$\mathcal{J}_M^- = \mathcal{S}_{-1}(\mathcal{J}_M) + \chi_M \mathbf{K}, \quad (1.7)$$

where the operator \mathcal{S}_{-1} defined in (2.18) shifts the mode number of the \hat{E}_8 loop generators and the Virasoro generators. The additional scalar field χ_M appears as the component along the \mathfrak{e}_9 central element \mathbf{K} , and is necessary to ensure covariance of the shifted current under rigid E_9 transformations. The first two terms in (1.5) are the E_9 version of the generic terms that appear in the potential of all exceptional field theories [14]. The third term contains the new constrained scalar field χ_M via (1.7) and generalises a structure which has so far only occurred in the potential of the E_8 exceptional field theory [24]. Each term in (1.5) is separately invariant under rigid \hat{E}_8 and scales with the same weight under rigid \mathbb{R}_d^+ , which is analogous to the homogeneous scaling of higher-dimensional exceptional field

theory Lagrangians under the trombone symmetry. The relative coefficients of the various terms in (1.5) are fixed by generalised diffeomorphism invariance.

In the bulk of the paper we shall also derive the potential away from $\tilde{\rho} = 0$ and in this way restore the full invariance under generalised diffeomorphisms and rigid $\mathbb{R}_{L_{-1}}$ transformations. All the terms in (1.5) then become functions of $\tilde{\rho}$ and its derivatives, such that they are invariant under rigid $\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ transformations up to homogeneous scaling under \mathbb{R}_d^+ . Moreover, the same relative combination as in (1.5) becomes invariant under all generalised diffeomorphisms as we shall demonstrate in detail.

The construction of E_9 exceptional field theory is interesting for several reasons. To begin with, it yields the first example of an exceptional field theory based on an infinite-dimensional duality group with fields and coordinates transforming in infinite-dimensional representations. As an immediate application, the E_9 potential can provide a prediction for the yet elusive scalar potential of gauged maximal $D = 2$ supergravity [31] by performing a generalised Scherk-Schwarz reduction. The $D = 2$ potential seems at present inaccessible by standard supersymmetry considerations because of the intricacies of $K(E_9)$ representation theory. As two-dimensional gauged supergravities generically involve a gauging of the $\mathbb{R}_{L_{-1}}$ symmetry [31], it is crucial to construct the E_9 potential at $\tilde{\rho} \neq 0$, which is invariant under all generalised diffeomorphisms. Another possible application is the study of non-geometric backgrounds [1, 2, 6, 8]. Moreover, $D = 2$ supergravity is the arena for exotic branes of co-dimension two (or lower) [32, 33] for which E_9 exceptional field theory may provide the appropriate framework [34–36].

Our construction does not depend on the details of the group E_8 and in fact the expressions we give will be valid for any simple group G and its affine extension \hat{G} . This provides the potential for extended field theories with coordinates in the basic representation of \hat{G} that are invariant under rigid $\hat{G} \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ and \hat{G} generalised diffeomorphisms.

The rest of this paper is organised as follows. In section 2 we review some basic facts and properties of the exceptional algebra \mathfrak{e}_9 and its representations. Section 3 introduces the building blocks for E_9 exceptional field theory by reviewing the field content of $D = 2$ maximal supergravity and the E_9 generalised Lie derivative from [13]. We furthermore introduce the covariantly constrained scalar fields χ_M . Section 4 presents the main result of this paper, the construction of the E_9 potential $V(\mathcal{M}, \chi)$ invariant under generalised diffeomorphisms. Finally, in section 5 we consider the branching under E_8 and show that the E_9 potential reproduces all the terms of the E_8 exceptional field theory that do not depend on the two-dimensional external derivatives. This shows that after solving the section condition, the E_9 potential $V(\mathcal{M}, \chi)$ reproduces $D = 11$ and type IIB supergravity for field configurations constant along the two-dimensional external spacetime. We finish with conclusions in section 6 and two appendices that contain some technical details and identities.

2 E_9 basic representation and Virasoro algebra

In this section, we introduce some notions we require from E_9 along with our notation to be used throughout the paper.

2.1 E_9 preliminaries

At the Lie algebra level, E_9 is an infinite-dimensional Kac-Moody algebra that we call \mathfrak{e}_9 . The core part of \mathfrak{e}_9 is the centrally extended loop algebra $\hat{\mathfrak{e}}_8$ over the Lie algebra \mathfrak{e}_8 and we only work with the split real forms. The Lie algebra \mathfrak{e}_8 has dimension 248 and we denote its generators by T^A with $A = 1, \dots, 248$ and \mathfrak{e}_8 structure constants

$$\mathfrak{e}_8 : \quad [T^A, T^B] = f^{AB}{}_C T^C. \quad (2.1)$$

The \mathfrak{e}_8 -invariant and non-degenerate metric is η^{AB} with inverse η_{AB} . The loop generators of $\hat{\mathfrak{e}}_8$ are denoted by T_m^A with mode number $m \in \mathbb{Z}$ and commutation relations

$$[T_m^A, T_n^B] = f^{AB}{}_C T_{m+n}^C + m \eta^{AB} \delta_{m,-n} \mathbf{K}, \quad (2.2)$$

where \mathbf{K} denotes the central extension of the loop algebra with $[\mathbf{K}, T_m^A] = 0$. In order to define the affine algebra \mathfrak{e}_9 one also has to adjoin the derivation operator \mathbf{d} that satisfies

$$[\mathbf{d}, T_m^A] = -m T_m^A, \quad [\mathbf{d}, \mathbf{K}] = 0. \quad (2.3)$$

As a vector space, $\mathfrak{e}_9 = \hat{\mathfrak{e}}_8 \oplus \langle \mathbf{d} \rangle$. There is an isomorphic copy of \mathfrak{e}_8 embedded in \mathfrak{e}_9 by considering the generators T_0^A at mode number zero. In terms of a loop parameter w , the loop generators can be realised in the adjoint representation as $T_m^A \sim w^m T^A$, $\mathbf{d} \sim -w \partial_w$ [37].

The above defines the adjoint representation of \mathfrak{e}_9 and we will also require other representations. Irreducible highest or lowest weight representations can be constructed in a Fock space manner as reviewed for example in [13, 37]. Here, we focus on the so-called basic representation that starts from an \mathfrak{e}_8 invariant ground state and we shall employ a Fock space notation. The ground state $|0\rangle$ satisfies

$$\begin{aligned} T_0^A |0\rangle = 0, \quad \mathbf{K} |0\rangle = |0\rangle, \quad \mathbf{d} |0\rangle = h |0\rangle \\ \text{and } T_n^A |0\rangle = 0 \quad \text{for } n > 0. \end{aligned} \quad (2.4)$$

While the eigenvalue of \mathbf{K} is fixed to one by unitarity, the eigenvalue of \mathbf{d} is a priori not determined. There is a one-parameter family of basic representations labelled by h that appears in $\mathbf{d} |0\rangle = h |0\rangle$, and we shall denote these representations by $R(\Lambda_0)_h$.² In general, the eigenvalue of the central element \mathbf{K} on an irreducible module is an integer called the affine level.

Note that \mathfrak{e}_9 is not simple as \mathbf{K} is central and \mathbf{d} never appears on the right-hand side of any commutator. Thus \mathfrak{e}_9 admits a one-dimensional representation ρ^h on which $\hat{\mathfrak{e}}_8$ acts trivially and \mathbf{d} acts with eigenvalue $-h/2$. The module $R(\Lambda_0)_h$ can therefore be defined as the product representation $R(\Lambda_0)_h = \rho^{-2h} R(\Lambda_0)_0$.

General elements in the basic representation will be denoted by ket-vectors $|V\rangle$ and can be expanded as

$$|V\rangle = \left(V^0 + \sum_{n=1}^{\infty} V_{A_1 \dots A_n} T_{-1}^{A_1} \dots T_{-1}^{A_n} \right) |0\rangle. \quad (2.5)$$

²We have changed conventions with respect to [13], where such representations were denoted $R(\Lambda_0)_{-h}$.

This representation of \mathfrak{e}_9 is made irreducible by removing all singular vectors (submodules) that arise when acting on $|0\rangle$. As a consequence, each coefficient $V_{A_1 \dots A_n}$ is constrained to live in the subset of irreducible \mathfrak{e}_8 representations contained in $\otimes^n \mathbf{248}$ according to the graded decomposition of $R(\Lambda_0)_h$ under \mathfrak{e}_8

$$R(\Lambda_0)_h = \mathbf{1}_h \oplus \mathbf{248}_{h+1} \oplus (\mathbf{1} \oplus \mathbf{248} \oplus \mathbf{3875})_{h+2} \oplus \dots \tag{2.6}$$

The subscript on the \mathfrak{e}_8 representations denotes their d eigenvalue. This representation of the Lie algebra \mathfrak{e}_9 is integrable and can be lifted to a representation of the affine Kac-Moody group $E_9 = \hat{E}_8 \rtimes \mathbb{R}_d^+$. Subtleties in defining this infinite-dimensional group will be discussed in section 4.3.

At some places we shall also utilise an index notation for elements of the basic representation rather than a Fock space notation. Choosing an infinite countable basis of the Fock space module $|e_M\rangle$ with $M \in \{0; A; \dots\}$ a collection of \mathfrak{e}_8 indices reproducing the decomposition (2.6), equation (2.5) becomes

$$|V\rangle = V^M |e_M\rangle, \tag{2.7}$$

so that components of vectors are V^M and we will refer to M as a ‘fundamental index’. We will use the bases $|e_M\rangle$ and indices M, N, \dots to label the components of $R(\Lambda_0)_h$ for all h , understanding that they characterise the \hat{E}_8 basic representation, whereas h labels the representation under \mathbb{R}_d^+ . This convention is defined such that all the E_9 group elements g are understood to be defined in the $R(\Lambda_0)_0$ representation, and the additional factor of $\rho(g)^{-2h}$ will be written explicitly. Note that for finite dimensional groups E_n with $n < 9$, the symmetry of exceptional field theories is $E_n \times \mathbb{R}^+$, and one writes various E_n tensors of different weight with respect to \mathbb{R}^+ . For E_9 the structure is very similar except that E_9 is only a semi-product $E_9 = \hat{E}_8 \rtimes \mathbb{R}_d^+$.

We shall also require the representation $\overline{R(\Lambda_0)_h}$ conjugate to $R(\Lambda_0)_h$. Elements of the conjugate representation will be denoted by bra-vectors. As we shall review below, coordinates of the E_9 exceptional geometry belong to the $R(\Lambda_0)$ representation and derivatives to its conjugate. To describe objects in $\overline{R(\Lambda_0)_h}$ in index notation, we introduce a basis $\langle e^M|$ dual to $|e_M\rangle$ so that

$$\langle W| = W_M \langle e^M|. \tag{2.8}$$

Again, we use the same notation for any value of h , which will be specified separately.

2.2 Virasoro algebra

For the Fock space representation of the basic module (at affine level 1) we define, following Sugawara [38], Virasoro generators in the enveloping algebra by

$$L_m = \frac{1}{2(1+g^\vee)} \sum_{n \in \mathbb{Z}} \eta_{AB} : T_n^A T_{m-n}^B :, \tag{2.9}$$

where the colon denotes normal ordering such that the positive modes appear on the right. For \mathfrak{e}_8 the dual Coxeter number $g^\vee = 30$. The Virasoro generators (2.9) acting on the lowest weight basic representation satisfy the Virasoro algebra

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}m(m^2 - 1)\delta_{m,-n}\mathbf{K} \tag{2.10}$$

with $c = \frac{\dim \mathfrak{e}_8}{1+g^\vee} = 8$ and $\mathbf{K} = 1$ is the identity operator. The central charge $c = 8$ comes from the fact that the module $R(\Lambda_0)_0$ can be realised as the Hilbert space of the two-dimensional conformal field theory of eight free chiral bosons parameterising the E_8 torus on which $\hat{\mathfrak{e}}_8$ acts as the current algebra.³ The Virasoro algebra is valid on $R(\Lambda_0)_h$ for any h .

We will denote the span of all Virasoro generators by

$$\mathfrak{vir} = \langle L_m \mid m \in \mathbb{Z} \rangle . \tag{2.11}$$

We note that this space is not closed under commutation (2.10) due to the central extension that we identify with \mathbf{K} and that is already contained in $\hat{\mathfrak{e}}_8$. The maximal algebra that we shall consider in this paper is

$$\mathfrak{f} = \hat{\mathfrak{e}}_8 \oplus \mathfrak{vir} , \tag{2.12}$$

which is the loop algebra extended by \mathbf{K} and *all* Virasoro generators L_m . The Virasoro generators (2.9) act on any lowest weight \mathfrak{e}_9 representation which therefore is automatically a representation of $\hat{\mathfrak{e}}_8 \oplus \mathfrak{vir}$ where the sum is semi-direct according to

$$[L_m, T_n^A] = -nT_{m+n}^A , \quad [L_m, \mathbf{K}] = 0 . \tag{2.13}$$

We shall use more generally the notation \oplus to denote an indecomposable \hat{E}_8 (or E_9) representation $X \oplus Y$ such that X and the quotient $X \oplus Y / \{X \sim 0\}$ are submodules of \hat{E}_8 but $Y \subset X \oplus Y$ is not, because $T_n^A Y \not\subset Y$.

In terms of a loop parameter w , the loop generators can be realised in the adjoint representation as $L_m = -w^{m+1}\partial_m$.

From (2.13) we see that \mathfrak{d} acting on the module $R(\Lambda_0)_h$ can be identified with $L_0 + h$. Both \mathfrak{d} and L_0 satisfy therefore the same commutation relations with the loop algebra, but $L_0|0\rangle = 0$ for any h . In the basic representation, \mathfrak{e}_9 is the span of the generators $\{T_M^A, \mathbf{K}\}$ of $\hat{\mathfrak{e}}_8$ and L_0 such that

$$\mathfrak{e}_9 = \langle T_m^A, \mathbf{K}, \mathfrak{d} \rangle = \langle T_m^A, \mathbf{K}, L_0 \rangle . \tag{2.14}$$

As is well-known, the Virasoro algebra (2.10) has an $\mathfrak{sl}(2)$ subalgebra given by $\langle L_{-1}, L_0, L_1 \rangle$. The group generated by $\hat{\mathfrak{e}}_8 \oplus \langle L_{-1}, L_0, L_1 \rangle$ through the exponential map is

$$\hat{E}_8 \rtimes \mathrm{SL}(2) . \tag{2.15}$$

³For general extended loop groups \hat{G} , c is the rank of the group G , corresponding to the fact that the basic representation of $\hat{\mathfrak{g}}$ can be realised as the Hilbert space of c chiral bosons on the torus $\mathbb{R}^c/\Lambda_{\mathfrak{g}}$ with $\Lambda_{\mathfrak{g}}$ the even lattice generated by the simple roots of the simply-laced Lie algebra \mathfrak{g} .

This group was identified in [39] as a symmetry of ungauged maximal supergravity in two dimensions. The symmetry group of the E_9 exceptional field theory will turn out to be its subgroup generated by $\mathfrak{e}_9 \oplus \langle L_{-1} \rangle$,

$$\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}) . \tag{2.16}$$

This group acts on \mathfrak{f} in an indecomposable representation.

We will denote collectively the generators of \mathfrak{f} in the basic representation $R(\Lambda_0)_h$ by $\langle T^\alpha \rangle = \langle T_n^A, \mathbf{K}, L_n \rangle$. By construction they do not depend on h , and $\mathbf{K} = 1$ on the module, although we choose to write it explicitly for clarity.

We can define a collection of \hat{E}_8 invariant symmetric bilinear forms η_m by

$$\eta_{m\alpha\beta} T^\alpha \otimes T^\beta = \sum_{n \in \mathbb{Z}} \eta_{AB} T_n^A \otimes T_{m-n}^B - L_m \otimes \mathbf{K} - \mathbf{K} \otimes L_m . \tag{2.17}$$

For the value $m = 0$ the range of the generators T^α is restricted to $\langle T_n^A, \mathbf{K}, L_0 \rangle$ and this form defined in the basic representation $R(\Lambda_0)_0$ (for which $\mathbf{d} = L_0$) is the inverse of the standard invariant Killing form $\eta^{\alpha\beta}$ on \mathfrak{e}_9 . We shall also denote this form by just $\eta_{\alpha\beta} \equiv \eta_{0\alpha\beta}$. Similarly for η_m the range of the generators T^α is restricted to $\langle T_n^A, \mathbf{K}, L_m \rangle$ (for all $n \in \mathbb{Z}$ but only one m), which also defines an algebra for which η_m is an invariant non-degenerate bilinear form.

It will be very convenient to also introduce shift operators \mathcal{S}_m (for $m \in \mathbb{Z}$), acting on $\mathfrak{f} = \hat{\mathfrak{e}}_8 \oplus \mathfrak{vir}$ according to

$$\mathcal{S}_m(\mathbf{K}) = \delta_{m,0} \mathbf{K}, \quad \mathcal{S}_m(L_n) = L_{m+n}, \quad \mathcal{S}_m(T_n^A) = T_{m+n}^A . \tag{2.18}$$

\mathcal{S}_0 is the identity. Combining this definition with (2.17) we find a useful identity for $m \neq 0$

$$\eta_{(n+m)\alpha\beta} T^\alpha \otimes T^\beta = \eta_{m\alpha\beta} T^\alpha \otimes \mathcal{S}_m(T^\beta) - L_{n+m} \otimes \mathbf{K} . \tag{2.19}$$

It follows that the shift operators are not invariant under \hat{E}_8 . Their transformation properties are discussed in appendix A.

Finally, the Hermitian conjugate $T^{\alpha\dagger}$ in the representation $R(\Lambda_0)_h$ is defined as

$$L_n^\dagger = L_{-n}, \quad \mathbf{K}^\dagger = \mathbf{K}, \quad \mathbf{d} = \mathbf{d}^\dagger, \quad T_n^{A\dagger} = \eta_{AB} T_{-n}^B . \tag{2.20}$$

It acts on the shift operators as $\mathcal{S}_m(T^\alpha)^\dagger = \mathcal{S}_{-m}(T^{\alpha\dagger})$. We use the Hermitian conjugate to also define the maximal unitary subgroup $K(E_9) \subset E_9$ which consists of those elements $k \in E_9$ satisfying $k^\dagger k = k k^\dagger = 1$ when acting on $R(\Lambda_0)_h$.⁴

The representation of \mathfrak{f} on $\overline{R(\Lambda_0)_h}$ in terms of the generator T^α on the Hilbert space of bra vectors can be realised equivalently in terms of the generators $T^{\alpha\dagger}$ on the Hilbert space of ket vectors, using

$$(\langle W|X \rangle)^\dagger = X^\dagger|W \rangle \tag{2.21}$$

⁴For finite-dimensional groups, the notion of maximal unitary subgroup coincides with that of maximal compact subgroup.

where we will write $\langle W |^\dagger = |W \rangle$ and it will be clear from the context that $|W \rangle \in \overline{R(\Lambda_0)_h}$. One consequence of this is that $\langle W | d = \langle W | (L_0 + h)$. Note that the representation of \mathfrak{f} on $R(\Lambda_0)_h$ in terms of the generator $-T^\alpha$ on the Hilbert space of ket vectors only agrees with the conjugate representation for anti-Hermitian elements. In particular, the two modules are isomorphic modules of the maximal unitary subgroup $K(E_9) \subset E_9$.

3 Building blocks of E_9 exceptional field theory

Many of the variables of E_9 exceptional field theory can be extracted from the knowledge of $D = 2$ maximal supergravity, which we review first in a reduction from $D = 3$. Then we discuss the additional structures that enter the exceptional field theory, in particular the generalised Lie derivative, the section constraint and the presence of constrained fields.

3.1 $D = 2$ maximal supergravity fields

An affine symmetry of $D = 2$ gravity coupled to matter arises on-shell when it is obtained by dimensional reduction of a $D = 3$ system with scalars taking values in a symmetric space [16, 40–42]. In principle, all propagating bosonic fields in $D = 3$ can be dualized to scalars coupled minimally to a non-dynamical three-dimensional metric. Assuming an additional space-like isometry with Killing vector $\partial_3 \equiv \partial_\varphi$ in the three-dimensional space-time leads to a metric of the form

$$ds_3^2 = e^{2\sigma}(-dt^2 + dx^2) + \rho^2(d\varphi + A_\mu^{(3)}dx^\mu)^2, \tag{3.1}$$

where coordinates have been chosen to make the two-dimensional metric conformally flat with scale factor $e^{2\sigma}$. The variable ρ measures the size of the Killing direction. The index $\mu = 0, 1$ labels the two coordinates t and x on which all the fields depend.

The field $A_\mu^{(3)}$ is the Kaluza-Klein vector arising in the reduction from 3 to 2 dimensions. Vector fields in $D = 2$ are not propagating and can be eliminated locally by a gauge transformation. In the usual formulation of ungauged $D = 2$ supergravity with rigid E_9 symmetry (and its associated linear systems), this vector field is set to zero [17, 41]. However, the investigation of gauged supergravity in various dimensions has shown the importance of the hierarchy of tensor fields including the non-propagating ones [20, 21]. In particular, the gauging of $D = 2$ supergravity requires the introduction of an infinity of vector fields (including $A_\mu^{(3)}$) transforming in the basic representation of E_9 [31]. A similar requirement is expected to hold in exceptional field theory, but our goal in the present paper is to construct only the potential of E_9 exceptional field theory, which does not depend on vectors. Hence, we will postpone their analysis (and possibly that of higher rank forms) to future work.

The propagating scalar fields descend from $D = 3$ and can be arranged in a representative V_0 of the coset space $E_8/(\text{Spin}(16)/\mathbb{Z}_2)$, where $\text{Spin}(16)/\mathbb{Z}_2$ is chosen to act on V_0 from the left, or alternatively in a Hermitian E_8 matrix $M_0 = V_0^\dagger V_0$. Written in the adjoint representation of E_8 , the matrix takes the form M_{0AB} with $A, B \in \{1, \dots, 248\}$ of \mathfrak{e}_8 . The (bosonic) dynamics of the maximal $D = 2$ supergravity theory is entirely described in terms of V_0 together with the scalars σ and ρ arising from the metric (3.1).

However, this does not make the infinite-dimensional affine symmetry and the associated integrability of the $D = 2$ theory manifest. In order to exhibit this symmetry one has to use an infinite set of dual scalar fields (a.k.a. cascade of dual potentials) that are related to the original scalar fields by non-linear duality relations that are consistent with the equations of motion. These are manifested in a so-called linear system for a generating function of a spectral parameter w that should be identified with the loop parameter of the loop algebra $\hat{\mathfrak{e}}_8$ discussed in section 2.1.⁵

The dualisation of the scalar fields V_0 leads to scalar fields that parameterise the coset space $\hat{E}_8/K(E_9)$ where $K(E_9)$ denotes the maximal unitary subgroup of E_9 [41, 43] which agrees with the maximal unitary subgroup of \hat{E}_8 in the basic representation $R(\Lambda_0)_0$. Expanding around $w = \infty$ we can parameterise a coset representative of the centrally extended loop group \hat{E}_8 in the representation $R(\Lambda_0)_0$ as

$$\hat{V} = e^{-\sigma K} V_0 \exp(\eta_{AB} Y_1^A T_{-1}^B) \exp(\eta_{AB} Y_2^A T_{-2}^B) \dots \quad (3.2)$$

V_0 here is the $E_8/(\text{Spin}(16)/\mathbb{Z}_2)$ coset representative containing the original E_8 scalars of the $D = 3$ theory while the Y_n^A are scalar fields corresponding to the $\hat{\mathfrak{e}}_8$ generators T_{-n}^A for $n > 0$, and η_{AB} is the Cartan-Killing form over \mathfrak{e}_8 . The local group $K(E_9)$ acts on \hat{V} from the left while the rigid \hat{E}_8 acts from the right. In writing the coset representative of the centrally extended loop group \hat{E}_8 we have fixed a Borel gauge, meaning that only the negative mode loop generators T_{-n}^A appear. This corresponds to fixing the action of $K(E_9)$.

The fields Y_n^A are on-shell dual to the propagating E_8 scalar fields parameterising V_0 . The first duality relation is

$$\partial_\mu Y_1^A \eta_{AB} T^B = \rho \varepsilon_{\mu\nu} V_0^{-1} P^\nu V_0, \quad (3.3)$$

where the integrability of this equation is guaranteed by the equations of motion for V_0 . In the above equation, $P^\nu = \frac{1}{2}(\partial^\nu V_0 V_0^{-1} + (\partial^\nu V_0 V_0^{-1})^\dagger)$ denotes the coset component of the Maurer-Cartan form, i.e., the projection of $\partial^\nu V_0 V_0^{-1}$ to the 128 non-compact generators of E_8 .

Equation (3.3) and similar equations for the other Y_n^A are summarised in the linear system of the $D = 2$ maximal supergravity [17] whose precise form we do not require here. These infinitely many fields are required to realise the on-shell \hat{E}_8 symmetry.

According to (2.14), the full E_9 symmetry also requires the inclusion of the generator \mathfrak{d} . The scalar field of $D = 2$ supergravity associated with this generator is the dilaton ρ [16, 39]. This scalar field satisfies the free Klein-Gordon equation and is dual to an axion $\tilde{\rho}$ via

$$\partial_\mu \rho = \varepsilon_{\mu\nu} \partial^\nu \tilde{\rho}. \quad (3.4)$$

⁵We note that there are two spectral parameters that are relevant in gravity reduced to $D = 2$; one that is called the ‘constant spectral parameter’ and that we denote by w and another one, often called ‘space-time dependent spectral parameter’ that also depends on the $D = 2$ coordinates and that we denote by γ . The two are related by $\gamma + \gamma^{-1} = 2(w + \tilde{\rho})/\rho$, so that γ is a double cover of w . The ‘axion’ $\tilde{\rho}$ in this relation is defined in equation (3.4). One can define $K(E_9)$ groups with respect to both choices of spectral parameter and the one that is commonly used in the linear system is γ . The one that we are using when writing the representative (3.2) is the constant spectral parameter w .

Unlike for the E_8 scalars V_0 , this duality relation is linear and does not give rise to an infinite sequence of dual scalar fields.

The dilaton ρ and the axion $\tilde{\rho}$ parameterise the group $\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}$. The full coset space of relevance is therefore

$$\frac{\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})}{K(E_9)}. \quad (3.5)$$

In the $R(\Lambda_0)_0$ representation, where we can identify d with L_0 , we write the $\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}$ group element as

$$v = \rho^{-L_0} e^{-\tilde{\rho}L_{-1}}, \quad (3.6)$$

and the full coset representative becomes

$$\mathcal{V} = v\hat{V}. \quad (3.7)$$

It transforms from the left under the denominator group $K(E_9)$ and from the right under the rigid symmetry group $\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$, i.e. as $\mathcal{V} \rightarrow k\mathcal{V}g$.

Using the fact that v can be embedded into the $SL(2)$ group generated by L_{-1} , L_0 and L_{+1} , we can conveniently work with the Hermitian element

$$\mathcal{M} = \mathcal{V}^\dagger \mathcal{V} = \hat{V}^\dagger m \hat{V} \in \hat{E}_8 \times SL(2) \quad (3.8)$$

with

$$m = v^\dagger v = e^{-\tilde{\rho}L_1} \rho^{-2L_0} e^{-\tilde{\rho}L_{-1}}, \quad (3.9)$$

so that $\mathcal{M} = \mathcal{M}^\dagger$. We shall also decompose \mathcal{M} as follows

$$\mathcal{M} = \hat{V}^\dagger m \hat{V} = m \hat{g}_{\mathcal{M}} = \hat{g}_{\mathcal{M}}^\dagger m. \quad (3.10)$$

Note that while $m \in SL(2)$ satisfies $m^\dagger = m$, the \hat{E}_8 element $\hat{g}_{\mathcal{M}}$ does not. We stress that \mathcal{M} is defined as a group element in the $R(\Lambda_0)_0$ representation, in which $d = L_0$.

In the fundamental representation $\mathbf{2}$ of $SL(2)$, m can be written as the 2×2 matrix

$$m_{\mathbf{2}} = \rho^{-1} \begin{pmatrix} \rho^2 - \tilde{\rho}^2 & -\tilde{\rho} \\ \tilde{\rho} & 1 \end{pmatrix}. \quad (3.11)$$

We note that the Hermitian conjugate (2.20) isolates a non-compact unitary $SO(1,1)$ subgroup of $SL(2)$. This implies that the finite-dimensional $m_{\mathbf{2}}$ cannot be a symmetric matrix but rather satisfies $m_{\mathbf{2}} = \sigma_3 m_{\mathbf{2}}^T \sigma_3$. Note that the whole $SL(2)$ is an on-shell symmetry of ungauged supergravity in two dimensions [39], but the exceptional field theory potential will only exhibit the parabolic subgroup $\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}$ as symmetry.

The advantage of working with \mathcal{M} instead of \mathcal{V} is that it only transforms under the rigid $\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ symmetry of the coset space as

$$\mathcal{M} \rightarrow g^\dagger \mathcal{M} g, \quad g \in \hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}}). \quad (3.12)$$

Formally, for $\tilde{\rho} = 0$, the element v in (3.6) is simply a dilatation \mathbb{R}_d^+ and \mathcal{M} becomes an element of the affine E_9 group only. As we shall see many formulæ simplify for $\tilde{\rho} = 0$. Even though most of our derivations use \mathcal{M} for simplicity, a proper evaluation of the potential and definition of the dual scalar fields requires descending to \mathcal{V} , a step we shall explain at the end in section 4.3.

In summary, all scalar fields appearing in the $D = 2$ maximal supergravity theory can be packaged into the operator \mathcal{M} . It contains the 128 propagating fields in the $E_8/(\text{Spin}(16)/\mathbb{Z}_2)$ coset representative V_0 along with all their dual potentials Y_n^A as well as the dilaton ρ , the axion $\tilde{\rho}$ and the scale factor σ .

To give a more concrete idea of how the E_8 scalar fields parameterising M_0 and the dual potentials Y_n^A are embedded in \mathcal{M} , we now display some components of the inverse matrix \mathcal{M}^{MN} . One can give formal definitions of the index-notation coefficients \mathcal{M}_{MN} and \mathcal{M}^{MN} using (2.7) and (2.8) as

$$\mathcal{M} = \mathcal{M}_{MN} \langle e^M |^\dagger \otimes \langle e^N |, \quad \mathcal{M}^{-1} = \mathcal{M}^{MN} |e_M\rangle \otimes |e_N\rangle^\dagger. \quad (3.13)$$

Taking $|e_0\rangle = |0\rangle$, $|e_A\rangle = T_{-1A}|0\rangle$, the first few matrix components of \mathcal{M}^{MN} are computed from (3.8) in the parameterisation (3.2) and (3.9) as follows

$$\mathcal{M}^{00} = \langle 0 | \mathcal{M}^{-1} | 0 \rangle = e^{2\sigma}, \quad (3.14a)$$

$$\mathcal{M}^{0A} = \langle 0 | \mathcal{M}^{-1} T_1^{A\dagger} | 0 \rangle = -e^{2\sigma} Y_1^A, \quad (3.14b)$$

$$\mathcal{M}^{AB} = \langle 0 | T_1^A \mathcal{M}^{-1} T_1^{B\dagger} | 0 \rangle = e^{2\sigma} (\rho^2 M_0^{AB} + Y_1^A Y_1^B), \quad (3.14c)$$

with M_0^{AB} the matrix components of the inverse E_8 matrix $M_0^{-1} = V_0^{-1} (V_0^\dagger)^{-1}$. We have also used $M_0^{-1} (T_m^A)^\dagger M_0 = M_0^{AB} T_{mB}$ and the fact that $|0\rangle$ is $\text{SL}(2)$ invariant. The other dual potentials Y_m^A , $m > 1$, as well as $\tilde{\rho}$ start appearing at higher levels. Performing a similar expansion for \mathcal{M}_{MN} would give infinite divergent series in Y_m^A at each level in the graded decomposition. However, the way \mathcal{M} enters in the potential $V(\mathcal{M}, \chi)$ is such that the latter is well-defined for \mathcal{V} in the Borel gauge (3.2) and only involves finite combinations of terms, as we shall exhibit in section 4.3 and also in section 5 when we discuss the reduction to E_8 .

3.2 Generalised diffeomorphisms and scalar fields

As usual in the construction of exceptional field theory, the supergravity fields are the basic building blocks and are promoted to fields depending on both the ‘external coordinates’ and the ‘internal coordinates’ of the exceptional geometry. The E_9 exceptional geometry is characterised by its generalised diffeomorphisms that we shall review first.

3.2.1 Generalised Lie derivative

As shown in [13], the correct representation for coordinates and canonical generalised vectors in E_9 exceptional geometry is the basic representation $R(\Lambda_0)_{-1}$ discussed above.

Writing $|V\rangle$ for a vector field in this representation, the action of a generalised diffeomorphism reads

$$\mathcal{L}_{\Lambda,\Sigma}|V\rangle = \langle\partial_V|\Lambda\rangle|V\rangle - \eta_{\alpha\beta}\langle\partial_\Lambda|T^\alpha|\Lambda\rangle T^\beta|V\rangle - \langle\partial_\Lambda|\Lambda\rangle|V\rangle - \eta_{-1\alpha\beta}\text{Tr}(T^\alpha\Sigma)T^\beta|V\rangle. \quad (3.15)$$

This very compact expression uses the Fock space notation for \mathfrak{e}_9 representations and the bilinear forms (2.17). As the coordinates are valued in the $R(\Lambda_0)_{-1}$ representation just like generalised vectors, derivatives are in the dual $\overline{R(\Lambda_0)_{-1}}$ representation with \mathfrak{d} eigenvalue of the vacuum state $\langle 0|\mathfrak{d} = (-1)\langle 0|$. Derivatives are represented as bra-vectors $\langle\partial|$ with the subscript indicating which object they act on.

The generalised Lie derivative (3.15) depends on two gauge parameters, Λ and Σ . The first parameter Λ is the usual generalised diffeomorphism parameter and is also valued in $R(\Lambda_0)_{-1}$. It is thus written as a ket vector. The second parameter Σ is an extra constrained parameter that generalises a similar constrained parameter in the E_8 exceptional field theory [24]. For E_9 , Σ belongs to $R(\Lambda_0)_0 \otimes \overline{R(\Lambda_0)_{-1}}$ with a constrained ‘bra index’. This constraint will be spelt out below. The trace $\text{Tr}(T^\alpha\Sigma)$ is guaranteed to be finite due to the constrained nature of Σ .

Since the Fock space notation in (3.15) is different from that used for finite-dimensional symmetry groups, we provide a short translation into index notation using (2.7) and (2.8). Vectors carry an upper fundamental index M and co-vectors have a lower fundamental index. In this component notation, the gauge parameters have index structure Λ^M and Σ^N_M . The generalised Lie derivative then takes the index form

$$\begin{aligned} \mathcal{L}_{\Lambda,\Sigma}V^M &= \Lambda^N\partial_N V^M - \eta_{\alpha\beta}(T^\alpha)^P_Q(T^\beta)^M_N\partial_P\Lambda^Q V^N - \partial_N\Lambda^N V^M \\ &\quad - \eta_{-1\alpha\beta}(T^\alpha)^P_Q(T^\beta)^M_N\Sigma^Q_P V^N. \end{aligned} \quad (3.16)$$

More examples of translating between the Fock space notation and the index notation were given in [13].

The definition (3.15) generalises to any field Φ admitting a well-defined action of $\mathfrak{e}_9 \oplus \langle L_{-1}\rangle$, not necessarily in a highest/lowest weight representation

$$\mathcal{L}_{\Lambda,\Sigma}\Phi = \langle\partial_\Phi|\Lambda\rangle\Phi + \eta_{\alpha\beta}\langle\partial_\Lambda|T^\alpha|\Lambda\rangle\delta^\beta\Phi + \eta_{-1\alpha\beta}\text{Tr}(T^\alpha\Sigma)\delta^\beta\Phi, \quad (3.17)$$

where $\delta^\alpha\Phi$ is the variation of the field with respect to the global symmetry algebra $\mathfrak{e}_9 \oplus \langle L_{-1}\rangle$. We stress that this includes the variation with respect to the derivation \mathfrak{d} and not L_0 , thus reproducing the third term in (3.15) with $\mathfrak{d}|V\rangle = (L_0 - 1)|V\rangle$.

From the comparison of (3.16) with the common form of generalised Lie derivatives [7] one can already anticipate the form of the section constraint to be

$$\eta_{\alpha\beta}\langle\partial_1|T^\alpha \otimes \langle\partial_2|T^\beta + \langle\partial_1| \otimes \langle\partial_2| - \langle\partial_2| \otimes \langle\partial_1| = 0. \quad (3.18a)$$

This constraint defines the tensor Y^{MN}_{PQ} in (1.1) in the introduction. The above constraint has for consequence the additional constraints

$$\eta_{-n\alpha\beta}\langle\partial_1|T^\alpha \otimes \langle\partial_2|T^\beta = 0 \quad \text{for all } n > 0, \quad (3.18b)$$

$$\eta_{1\alpha\beta}(\langle\partial_1|T^\alpha \otimes \langle\partial_2|T^\beta + \langle\partial_2|T^\alpha \otimes \langle\partial_1|T^\beta) = 0. \quad (3.18c)$$

The constraint on the gauge parameter Σ is more conveniently written using the notation introduced in [13]

$$\Sigma = \Sigma^M{}_N |e^N\rangle \langle e_M| = |\Sigma\rangle \langle \pi_\Sigma|. \quad (3.19)$$

The section constraint (3.18) is then also imposed when one derivative is replaced by $\langle \pi_\Sigma|$, and when both derivatives are replaced with $\langle \pi_{\Sigma_1}|$, $\langle \pi_{\Sigma_2}|$, where Σ_1 , Σ_2 can be the same or two different gauge parameters. Notice that Σ generally does not factorize into a tensor product of a bra-vector $\langle \pi_\Sigma|$ with a ket-vector $|\Sigma\rangle$.⁶

As was shown in [13], the generalised Lie derivative (3.15) provides a closed gauge algebra when the section constraints are imposed, satisfying

$$[\mathcal{L}_{\Lambda_1, \Sigma_1}, \mathcal{L}_{\Lambda_2, \Sigma_2}] \Phi = \mathcal{L}_{\Lambda_{12}, \Sigma_{12}} \Phi, \quad (3.20)$$

with

$$\begin{aligned} \Lambda_{12} &= \frac{1}{2} (\mathcal{L}_{\Lambda_1} \Lambda_2 - \mathcal{L}_{\Lambda_2} \Lambda_1), \\ \Sigma_{12} &= \mathcal{L}_{\Lambda_1} \Sigma_2 + \frac{1}{2} \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma_1) T^\beta \Sigma_2 - \frac{1}{2} \eta_{1\alpha\beta} \langle \partial_{\Lambda_2} | T^\alpha | \Lambda_{[1} \rangle T^\beta | \Lambda_{2]} \rangle \langle \partial_{\Lambda_2} | - (1 \leftrightarrow 2). \end{aligned} \quad (3.21)$$

The parameter Σ_{12} depends nontrivially on Λ_1 and Λ_2 (as well as on Σ_1 and Σ_2), compensating for the fact that the Λ part of E_9 generalised diffeomorphisms does not close onto itself. An observation that will be crucial in the following is that closure of the generalised Lie derivative is ensured already by restricting to traceless Σ parameters, i.e. Σ_{12} is traceless if Σ_1 and Σ_2 are. As is clear from (3.15) and (2.17), the trace component $\text{Tr}(\Sigma)$ is the only one generating L_{-1} transformations. We can then gauge-fix the trace component of Σ transformations acting on \mathcal{M} by setting $\tilde{\rho} = 0$ and then consistently restrict to arbitrary Λ and traceless Σ transformations. No compensating gauge transformations are then needed to keep $\tilde{\rho} = 0$ and E_9 covariance is preserved.

3.2.2 Unconstrained scalar fields and currents

The fields of the theory include the element $\mathcal{M} \in \hat{E}_8 \rtimes \text{SL}(2)$ introduced in (3.8) and it depends on the coordinates of the exceptional geometry that take values in the $R(\Lambda_0)_{-1}$ representation. We reiterate that we always take \mathcal{M} to be defined as a group element in the representation $R(\Lambda_0)_0$. As a side-remark we note that this discrepancy between the weight h of the coordinate representation and the representation of \mathcal{M} means that the ‘generalised metric’ of E_9 exceptional geometry is $\rho^2 \mathcal{M}$.

From \mathcal{M} in $R(\Lambda_0)_0$ we can, as usual, construct the current

$$\mathcal{J}_M = \mathcal{M}^{-1} \partial_M \mathcal{M} \quad (3.22)$$

which is valued in the Lie algebra $\hat{e}_8 \oplus \mathfrak{sl}(2)$ and carries a constrained derivative index M in the $\overline{R(\Lambda_0)_{-1}}$ representation. In index notation and using the bases (2.7) and (2.8) this

⁶To write the constraint (3.18a) for the operator Σ itself one must introduce an additional arbitrary vector $|V\rangle$, such that for any $|V\rangle$: $\eta_{\alpha\beta} \Sigma T^\alpha \langle V | \partial | T^\beta + \Sigma | V \rangle \langle \partial | - \Sigma \langle \partial | V \rangle = 0$ and $\eta_{\alpha\beta} \Sigma_1 T^\alpha | V_1 \rangle \otimes \Sigma_2 T^\beta | V_2 \rangle + \Sigma_1 | V_1 \rangle \otimes \Sigma_2 | V_2 \rangle - \Sigma_1 | V_2 \rangle \otimes \Sigma_2 | V_1 \rangle = 0$.

reads

$$\mathcal{M}^{PS} \partial_M \mathcal{M}_{SQ} = (\mathcal{J}_M)^P_Q = \mathcal{J}_{M,\alpha} (T^\alpha)^P_Q, \quad (3.23)$$

where $T^\alpha \in \mathfrak{f}$ but $\mathcal{J}_{M,\alpha}$ is only non-vanishing along $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$. It will be more convenient to use the Fock space notation, in which the current is defined such that

$$\langle \partial_M | \otimes \mathcal{M} = \langle \mathcal{J}_\alpha | \otimes \mathcal{M} T^\alpha. \quad (3.24)$$

To see the equivalence of this formula with the definition (3.23) above one may reintroduce indices as

$$\partial_M \mathcal{M}_{PQ} = \mathcal{J}_{M,\alpha} \mathcal{M}_{PS} (T^\alpha)^S_Q \quad (3.25)$$

and multiply the equation with \mathcal{M}^{-1} . The current satisfies the following useful identity

$$\langle \mathcal{J}_\alpha | \otimes \mathcal{M}^{-1} T^{\alpha\dagger} \mathcal{M} = \langle \mathcal{J}_\alpha | \otimes T^\alpha, \quad (3.26)$$

which follows from the Hermiticity of \mathcal{M} . It will also be convenient to introduce the matrix $H(\mathcal{M})^\alpha_\beta$ defined by

$$\mathcal{M}^{-1} T^{\alpha\dagger} \mathcal{M} = H(\mathcal{M})^\alpha_\beta T^\beta, \quad (3.27)$$

in terms of which the identity (3.26) reads

$$H(\mathcal{M})^\beta_\alpha \mathcal{J}_\beta = \mathcal{J}_\alpha. \quad (3.28)$$

We write out the Lie algebra valued current in its components more explicitly as follows

$$\langle \mathcal{J}_\alpha | \otimes T^\alpha = \sum_{n \in \mathbb{Z}} \langle J_A^n | \otimes T_n^A + \sum_{q = -1, 0, 1} \langle J_q | \otimes L_q + \langle J_K | \otimes K. \quad (3.29)$$

The above expression stresses once again that, while our conventions are such that T^α are the generators of $\mathfrak{f} = \hat{\mathfrak{e}}_8 \oplus \mathfrak{vir}$, the current has only components along $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ since it is constructed from an element \mathcal{M} in the group $\hat{E}_8 \times \text{SL}(2)$ (3.8). In other words, we have $\langle J_q | = 0$ for $|q| > 1$.

We also note that the $\mathfrak{sl}(2)$ -part of the current is identical to that constructed solely out of the $\text{SL}(2)$ element m defined in (3.9),

$$m^{-1} \partial_M m = J_{M,0} L_0 + J_{M,1} L_1 + J_{M,-1} L_{-1}, \quad (3.30)$$

where, due to the structure of m , one has in Fock space notation

$$\langle J_{-1} | + \tilde{\rho} \langle J_0 | + (\tilde{\rho}^2 - \rho^2) \langle J_1 | = 0. \quad (3.31)$$

This relation can be derived easily from the matrix representation (3.11) and can be used to solve for $\langle J_{-1} |$ in terms of the other components.

3.2.3 Constrained scalar fields and shifted currents

A crucial ingredient in exceptional field theory is the existence of constrained fields. For E_{11-D} exceptional field theory with a D -dimensional external spacetime, the constrained fields appear starting from the $(D-2)$ -form sector. For instance, in the E_8 exceptional field theory, corresponding to $D=3$, there are constrained vector fields that are crucial in the construction of the theory [24]. In the present case of $D=2$, the constrained fields appear already in the scalar sector and therefore are essential for the scalar potential.

As was argued in [13], the constrained scalar fields carry an index labelling the dual basic representation $\overline{R(\Lambda_0)_{-1}}$ and we write them as χ_M or $\langle\chi|$. The fact that they are constrained means that they satisfy a condition analogous to the section constraint (3.18), namely

$$\eta_{\alpha\beta}\langle\chi|T^\alpha \otimes \langle\partial|T^\beta + \langle\chi| \otimes \langle\partial| - \langle\partial| \otimes \langle\chi| = 0, \quad (3.32a)$$

$$\eta_{-n\alpha\beta}\langle\chi|T^\alpha \otimes \langle\partial|T^\beta = 0 \quad \text{for all } n > 0, \quad (3.32b)$$

$$\eta_{1\alpha\beta}(\langle\chi|T^\alpha \otimes \langle\partial|T^\beta + \langle\partial|T^\alpha \otimes \langle\chi|T^\beta) = 0, \quad (3.32c)$$

and the same identities bilinear in $\langle\chi|$. Here, $\langle\partial|$ can be any derivative, as for instance that carried by the current $\langle\mathcal{J}_\alpha|$, or also the constrained index of the generalised diffeomorphism parameter Σ .

As will become clearer when we discuss the transformation properties of the various fields, it is useful to also define a shifted version of the current $\langle\mathcal{J}_\alpha|$ defined in (3.24), by making use of the shift operators defined in (2.18)

$$\langle\mathcal{J}_\alpha^-| \otimes T^\alpha = \sum_{k=0}^{\infty} \hat{\rho}^k \langle\mathcal{J}_\alpha| \otimes \mathcal{S}_{-1-k}(T^\alpha) + \langle\chi| \otimes \mathbb{K}. \quad (3.33)$$

The constrained scalar field χ appears in the definition of the shifted current in order to ensure covariance under the rigid symmetries, as we shall explain in detail below. Unlike the unshifted current (3.29), the shifted current has also non-trivial components along the Virasoro generators L_q for all $q \leq 1$ because these are generated by the shift operators.

4 The potential

In this section, we present the E_9 exceptional field theory potential, depending on \mathcal{M} and the constrained scalar field χ , written as a sum of four terms as

$$V(M, \chi) = \frac{1}{4}\mathcal{L}_1 - \frac{1}{2}\mathcal{L}_2 + \frac{1}{2}\mathcal{L}_3 + \frac{1}{2}\mathcal{L}_4. \quad (4.1)$$

The four terms are independently invariant under rigid \hat{E}_8 and $\mathbb{R}_{L_{-1}}$ transformations, and transform with the expected homogeneous scaling under \mathbb{R}_d^+ . The symmetry \mathbb{R}_d^+ is known to be a symmetry of the equations of motion, whereas the potential itself is not invariant but transforms homogeneously with weight one. The relative coefficients between the four terms are fixed by requiring the potential to transform into a total derivative under generalised diffeomorphisms.

Let us briefly compare the general structure of the potential to that of other E_n exceptional field theories with $n \leq 8$. While the terms \mathcal{L}_1 , \mathcal{L}_2 and \mathcal{L}_4 have direct analogues in the other cases [14, 22–24], the term \mathcal{L}_3 is a vast generalisation of a term that only appears in E_8 exceptional field theory. The main new feature is that the E_9 scalar fields \mathcal{M} and χ form an *indecomposable* representation, meaning that they cannot be separated into a direct sum of irreducible E_9 representations. The term \mathcal{L}_3 contains crucially the shifted current (3.33) that comprises all these scalar fields. Another consequence of this indecomposability of the scalar fields is that the four individual terms of the potential are not all manifestly invariant under the rigid $\hat{E}_8 \rtimes \mathbb{R}_{L_{-1}}$ transformations and we shall therefore demonstrate this invariance explicitly.

For the sake of clarity of the presentation we shall first consider a partially gauge-fixed version of the potential in which the axion $\tilde{\rho} = 0$ and the rigid symmetry $\mathbb{R}_{L_{-1}}$ is broken. As mentioned in section 3.2.1, this gauge-fixing preserves the closed subalgebra of generalised diffeomorphisms (3.15) with $\text{Tr}(\Sigma) = 0$. In this case, the four terms are manifestly invariant under \hat{E}_8 . The term \mathcal{L}_3 still retains the indecomposable structure but now involves only the shift operator \mathcal{S}_{-1} (instead of all \mathcal{S}_n with $n < 0$). This shift operator and the associated shifted bilinear form η_{-1} are also expected on the basis of the structure of maximal gauged supergravity [31] where the embedding tensor couples through η_{-1} .

In a second step, we reintroduce the $\tilde{\rho}$ dependence and consequently the full generalised diffeomorphism invariance. Besides generalised diffeomorphism invariance, the full potential presented in this section is invariant under rigid $\hat{E}_8 \rtimes \mathbb{R}_{L_{-1}}$ and these two requirements uniquely fix the combination of the four individual terms. In the following section 5, we moreover demonstrate that our potential, upon choosing an appropriate solution to the section condition, reproduces all the terms in the E_8 exceptional field theory [24] that can contribute to the E_9 potential. This provides a final check on the E_9 potential.

4.1 The potential at $\tilde{\rho} = 0$

In this section we restrict ourselves to the case $\tilde{\rho} = 0$ in which \mathcal{M} , defined in (3.8), is an element of E_9 in the $R(\Lambda_0)_0$ representation. The construction of the potential is greatly simplified in this setting as one simply requires its invariance under rigid \hat{E}_8 transformations and Λ generalised diffeomorphisms. The various terms of the potential then read

$$\mathcal{L}_1 = \rho^{-1} \eta^{\alpha\beta} \langle \mathcal{J}_\alpha | \mathcal{M}^{-1} | \mathcal{J}_\beta \rangle, \tag{4.2a}$$

$$\mathcal{L}_2 = \rho^{-1} \langle \mathcal{J}_\alpha | T^\beta \mathcal{M}^{-1} T^{\alpha\dagger} | \mathcal{J}_\beta \rangle, \tag{4.2b}$$

$$\mathcal{L}_3 = \rho \langle \mathcal{J}_\alpha^- | T^\beta \mathcal{M}^{-1} T^{\alpha\dagger} | \mathcal{J}_\beta^- \rangle, \tag{4.2c}$$

$$\mathcal{L}_4 = \rho^{-1} \langle J_0 | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle. \tag{4.2d}$$

Their expression in an index notation was already given in the introduction (1.5). The currents $\langle \mathcal{J}_\alpha |$ and $\langle \mathcal{J}_\alpha^- |$ were defined in (3.24) and (3.33), and since here $\mathcal{M} \in E_9$, their only non-vanishing components are along \mathfrak{e}_9 and $\hat{\mathfrak{e}}_8 \oplus L_{-1}$, respectively. Since here $\tilde{\rho} = 0$, the current component along L_0 is simply given by

$$\langle J_0 | = -2 \rho^{-1} \langle \partial_\rho | \rho. \tag{4.3}$$

We also point out that, while writing some of the currents as kets in (4.2) might seem confusing at first, our notation should be clear from the discussion at the end of section 2.2.

All the terms in (4.2) are manifestly Hermitian and, as we shall see below, invariant under rigid \hat{E}_8 transformations. Moreover, note that in this case the new constrained scalar field $\langle \chi |$ only appears in the third term. As previously mentioned, \mathcal{L}_3 generalises a term that so far only appeared in the E_8 potential, where it involved two E_8 currents contracted directly [24].

4.1.1 Rigid E_9 symmetry

Under $g \in E_9$, we have the following rigid transformations

$$\mathcal{M} \rightarrow g^\dagger \mathcal{M} g, \tag{4.4}$$

$$\langle \partial | \rightarrow \rho(g)^2 \langle \partial | g, \tag{4.5}$$

$$|Y\rangle \rightarrow \rho(g)^{-2} g^{-1} |Y\rangle, \tag{4.6}$$

where here and in the following, g will always be a group element in the $R(\Lambda_0)_0$ representation. It acts on the derivative bra in the $\overline{R(\Lambda_0)_{-1}}$ representation by multiplication with g from the right and on the coordinate ket in the $R(\Lambda_0)_{-1}$ by multiplication with g^{-1} from the left. The \mathbb{R}^+ -valued function $\rho(g)$ appearing in the transformation laws is the multiplicative character defined in (A.4). It occurs for instance in the above second variation to account for the fact that the derivative transforms in the $\overline{R(\Lambda_0)_{-1}}$ representation while g is a group element in the $R(\Lambda_0)_0$ representation.⁷ In particular, we have $\rho(\mathcal{M}) = \rho$.

The variation of the \mathfrak{e}_9 -valued current (3.24) then reads

$$\langle \mathcal{J}_\alpha | \otimes T^\alpha \rightarrow \rho(g)^2 \langle \mathcal{J}_\alpha | g \otimes g^{-1} T^\alpha g, \tag{4.7}$$

while its components $\langle \mathcal{J}_\alpha |$ transform as

$$\langle \mathcal{J}_\alpha | \rightarrow \rho(g)^2 R(g)^\beta{}_\alpha \langle \mathcal{J}_\beta | g, \tag{4.8}$$

where $R(g)^\alpha{}_\beta$ denotes the representation matrix of g acting by conjugation, which is defined by

$$R(g)^\alpha{}_\beta T^\beta = g^{-1} T^\alpha g. \tag{4.9}$$

To further clarify our notation for the current written as a ket, we also provide explicitly its transformation under E_9 ,

$$\begin{aligned} T^\alpha \otimes |\mathcal{J}_\alpha\rangle &\rightarrow g^{-1} T^\alpha g \otimes g^\dagger |\mathcal{J}_\alpha\rangle \rho(g)^2, \\ |\mathcal{J}_\alpha\rangle &\rightarrow g^\dagger |\mathcal{J}_\beta\rangle R(g)^\beta{}_\alpha \rho(g)^2, \end{aligned} \tag{4.10}$$

which follows from (2.21). The variation of the scalar $\rho = \rho(\mathcal{M})$ is by definition

$$\rho \rightarrow \rho(g)^2 \rho. \tag{4.11}$$

⁷We note that \mathcal{M} as a group element is in $R(\Lambda_0)_0$ which naturally multiplies ket vectors in $R(\Lambda_0)_{-h}$ from the left and produces bra vectors in $R(\Lambda_0)_{+h}$. It thus acts as an intertwiner of representations.

From (4.3), one then simply finds that $\langle J_0 | \rightarrow \rho(g)^2 \langle J_0 | g$. Using the above transformations it is straightforward to verify the rigid \hat{E}_8 invariance of (4.2b) and (4.2d). The \hat{E}_8 invariance of (4.2a) is ensured by the presence of the \mathfrak{e}_9 invariant bilinear form $\eta^{\alpha\beta}$, which satisfies $R(g)^\alpha_\gamma R(g)^\beta_\delta \eta^{\gamma\delta} = \eta^{\alpha\beta}$.

The invariance of (4.2c) is a bit more subtle. Start by considering the variation of the current which has been acted upon by the shift operator defined in (2.18),

$$\langle \mathcal{J}_\alpha | \otimes \mathcal{S}_{-1}(T^\alpha) \rightarrow \rho(g)^2 \langle \mathcal{J}_\alpha | g \otimes (\rho(g)^{-2} g^{-1} \mathcal{S}_{-1}(T^\alpha) g - \omega_1^\alpha(g^{-1}) \mathbf{K}). \quad (4.12)$$

This result follows directly from using the relation (A.13). It involves an \mathbb{R} -valued function $\omega_1^\alpha(g)$, which is a group 1-cocycle defined in (A.13). As explained in detail in appendix A, this cocycle and the character $\rho(g)$ define an extension of the adjoint representation of E_9 by the generator L_{-1} . The new scalar field $\langle \chi |$ must be chosen to transform as part of the dual of this extended representation (A.11), i.e. as

$$\langle \chi | \rightarrow \langle \chi | g + \rho(g)^2 \omega_1^\alpha(g^{-1}) \langle \mathcal{J}_\alpha | g, \quad (4.13)$$

in order for the shifted current to transform covariantly under E_9 . Indeed, using (4.12) and (4.13) we find that the shifted current (3.33) and its components transform as

$$\langle \mathcal{J}_\alpha^- | \otimes T^\alpha \rightarrow \langle \mathcal{J}_\alpha^- | g \otimes g^{-1} T^\alpha g, \quad (4.14)$$

$$\langle \mathcal{J}_\alpha^- | \rightarrow R(g)^\beta_\alpha \langle \mathcal{J}_\beta^- | g. \quad (4.15)$$

It is then straightforward to verify that the third term in the potential is invariant under \hat{E}_8 . Note that the appearance of fields transforming in indecomposable representations, such as $\langle \chi |$ in (4.13), is a new feature in E_9 exceptional field theory. In higher-dimensions, all the fields have to transform individually in irreducible representations of the duality group since E_n is then a finite-dimensional reductive group.

Let us finally remark that the potential scales uniformly under transformations generated by \mathfrak{d} ,

$$V \rightarrow \rho(g)^2 V. \quad (4.16)$$

As mentioned previously, the generator \mathfrak{d} is associated to a symmetry of the equations of motion and not of the Lagrangian itself, as is the so-called trombone symmetry in higher dimensions [44]. According to its original definition, the trombone symmetry in two dimensions shifts the conformal factor σ of the metric and is the symmetry of the action generated by the central charge \mathbf{K} . It is instead the symmetry generated by \mathfrak{d} that rescales the dilaton field ρ which is not a symmetry of the action.

4.1.2 Invariance under generalised diffeomorphisms

We denote an infinitesimal variation under generalised diffeomorphisms by $\delta_{\Lambda, \Sigma}$. By definition it splits into

$$\delta_{\Lambda, \Sigma} = \mathcal{L}_{\Lambda, \Sigma} + \Delta_{\Lambda, \Sigma}, \quad (4.17)$$

where the action of the generalised Lie derivative $\mathcal{L}_{\Lambda, \Sigma}$ on an arbitrary field was defined in (3.17), and where $\Delta_{\Lambda, \Sigma}$ explicitly collects all the non-covariant pieces in the variation. The latter are those terms involving second derivatives of the gauge parameter Λ or a single derivative of the gauge parameter Σ . In this section, we exclusively focus on variations under Λ generalised diffeomorphisms, which are simply denoted by δ_{Λ} . The reason is that in the expression of the generalised Lie derivative, the trace of Σ appears as the gauge parameter of an infinitesimal \mathbb{R}_{L-1} transformation, which can only be considered at $\tilde{\rho} \neq 0$ when $\mathcal{M} \in \hat{E}_8 \times \text{SL}(2) \supset \mathbb{R}_{L-1}$. While we could already consider traceless Σ variations in this section, we postpone this discussion to section 4.2 where we will prove the invariance of the full potential under arbitrary Σ variations.

We start with \mathcal{M} which transforms covariantly under generalised diffeomorphisms, i.e. as

$$\delta_{\Lambda} \mathcal{M} = \mathcal{L}_{\Lambda} \mathcal{M} = \langle \partial_{\mathcal{M}} | \Lambda \rangle \mathcal{M} + \eta_{\alpha\beta} \langle \partial_{\Lambda} | T^{\alpha} | \Lambda \rangle (\mathcal{M} T^{\beta} + T^{\beta\dagger} \mathcal{M}). \quad (4.18)$$

The expression of the rotation term follows from the action (4.4) of E_9 on \mathcal{M} . In particular, there is no density term as \mathcal{M} transforms (from the right and the left) in the $R(\Lambda_0)_0$ representation and therefore carries no weight. According to (4.11), the field ρ is an E_9 scalar density of weight one and thus transforms as a total derivative

$$\delta_{\Lambda} \rho = \mathcal{L}_{\Lambda} \rho = \langle \partial_{\rho} | \Lambda \rangle \rho + \langle \partial_{\Lambda} | \Lambda \rangle \rho = \langle \partial | (| \Lambda \rangle \rho). \quad (4.19)$$

The variation of the current (3.24) follows from that of \mathcal{M} and takes the form

$$\delta_{\Lambda} \langle \mathcal{J}_{\alpha} | \otimes T^{\alpha} = \mathcal{L}_{\Lambda} \langle \mathcal{J}_{\alpha} | \otimes T^{\alpha} + \eta_{\alpha\beta} \langle \partial_{\Lambda} | T^{\alpha} | \Lambda \rangle \langle \partial_{\Lambda} | \otimes (T^{\beta} + \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}), \quad (4.20)$$

where its Lie derivative is given by

$$\mathcal{L}_{\Lambda} \langle \mathcal{J}_{\alpha} | \otimes T^{\alpha} = (\langle \partial_{\mathcal{J}} | \Lambda \rangle \langle \mathcal{J}_{\alpha} | + \langle \mathcal{J}_{\alpha} | \Lambda \rangle \langle \partial_{\Lambda} |) \otimes T^{\alpha} - \eta_{\alpha\beta} \langle \partial_{\Lambda} | T^{\alpha} | \Lambda \rangle \langle \mathcal{J}_{\gamma} | \otimes [T^{\beta}, T^{\gamma}]. \quad (4.21)$$

The variation of the current components then reads

$$\delta_{\Lambda} \langle \mathcal{J}_{\alpha} | = \mathcal{L}_{\Lambda} \langle \mathcal{J}_{\alpha} | + \eta_{\alpha\beta} \langle \partial_{\Lambda} | (T^{\beta} + \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}) | \Lambda \rangle \langle \partial_{\Lambda} |, \quad (4.22)$$

with

$$\mathcal{L}_{\Lambda} \langle \mathcal{J}_{\alpha} | = \langle \partial_{\mathcal{J}} | \Lambda \rangle \langle \mathcal{J}_{\alpha} | + \langle \mathcal{J}_{\alpha} | \Lambda \rangle \langle \partial_{\Lambda} | - \eta_{\gamma\delta} \langle \partial_{\Lambda} | T^{\gamma} | \Lambda \rangle f^{\delta\beta}{}_{\alpha} \langle \mathcal{J}_{\beta} |, \quad (4.23)$$

where $f^{\alpha\beta}{}_{\gamma}$ denotes the structure constants of \mathfrak{f} . To write the non-covariant terms we also used the identity

$$\eta_{-n\alpha\beta} \mathcal{M}^{-1} T^{\alpha\dagger} \mathcal{M} \otimes T^{\beta} = \rho^{-2n} \eta_{n\alpha\beta} T^{\alpha} \otimes \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \quad (4.24)$$

which only holds here as $\mathcal{M} \in E_9$, and follows from the covariance of $\eta_{n\alpha\beta}$ under E_9 . From the non-covariant variation in (4.22), one gets in particular

$$\Delta_{\Lambda} \langle J_0 | = -2 \langle \partial_{\Lambda} | \Lambda \rangle \langle \partial_{\Lambda} |. \quad (4.25)$$

Let us now discuss the variation of the shifted current. Acting with the shift operator \mathcal{S}_{-1} on (4.20) gives

$$\begin{aligned}
 \delta_\Lambda \langle \mathcal{J}_\alpha | \otimes \mathcal{S}_{-1}(T^\alpha) &= (\langle \partial_{\mathcal{J}} | \Lambda \rangle \langle \mathcal{J}_\alpha | + \langle \mathcal{J}_\alpha | \Lambda \rangle \langle \partial_\Lambda | - \langle \partial_\Lambda | \Lambda \rangle \langle \mathcal{J}_\alpha |) \otimes \mathcal{S}_{-1}(T^\alpha) \\
 &\quad - \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \mathcal{J}_\gamma | \otimes [T^\beta, \mathcal{S}_{-1}(T^\gamma)] - \sum_{n \in \mathbb{Z}} (n-1) \langle \partial_\Lambda | T_{n-1}^A | \Lambda \rangle \langle J_A^n | \otimes \mathbb{K} \\
 &\quad + \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \otimes (\rho^{-2} \mathcal{M}^{-1} \mathcal{S}_1(T^\beta)^\dagger \mathcal{M} - \omega_{-1}^\beta(\mathcal{M}) \mathbb{K} + \mathcal{S}_{-1}(T^\beta)). \quad (4.26)
 \end{aligned}$$

The density term and the term involving the explicit sum over the loop algebra are generated by pulling the shift operator inside of the commutator in the second term of the second line, while the appearance of the cocycle is a consequence of using the identity (A.13). The variation of the constrained field $\langle \chi |$ is chosen to be

$$\begin{aligned}
 \delta_\Lambda \langle \chi | &= \mathcal{L}_\Lambda \langle \chi | - \langle \partial_\Lambda | (L_{-1} + \rho^{-2} L_1 - \eta_{\alpha\beta} \omega_{-1}^\alpha(\mathcal{M}) T^\beta) | \Lambda \rangle \langle \partial_\Lambda | \\
 &= \mathcal{L}_\Lambda \langle \chi | - \langle \partial_\Lambda | (L_{-1} + \mathcal{M}^{-1} L_{-1}^\dagger \mathcal{M}) | \Lambda \rangle \langle \partial_\Lambda |, \quad (4.27)
 \end{aligned}$$

with the Lie derivative

$$\mathcal{L}_\Lambda \langle \chi | = \langle \partial_\chi | \Lambda \rangle \langle \chi | + \langle \chi | \Lambda \rangle \langle \partial_\Lambda | - \langle \partial_\Lambda | \Lambda \rangle \langle \chi | + \sum_{n \in \mathbb{Z}} (n-1) \langle \partial_\Lambda | T_{n-1}^A | \Lambda \rangle \langle J_A^n |. \quad (4.28)$$

The Lie derivative is determined according to (3.17) and the linearisation of the E_9 action (4.13) on the field $\langle \chi |$,

$$\delta_X \langle \chi | \equiv X_\alpha \delta^\alpha \langle \chi | = \langle \chi | X_\alpha T^\alpha + \eta^{AB} \sum_{n \in \mathbb{Z}} (n-1) X_A^{1-n} \langle J_B^n | \quad (X \in \mathfrak{e}_9 \text{ only}), \quad (4.29)$$

which follows from using the section constraint and the linearisation of the cocycle (A.14). Note furthermore that the non-covariant variations in (4.27) are consistent with the property that $\langle \chi |$ transforms as part of the dual of the extended representation (A.11) which includes L_{-1} . Combining (4.26) and (4.27) yields for the shifted current

$$\begin{aligned}
 \delta_\Lambda \langle \mathcal{J}_\alpha^- | \otimes T^\alpha &= \mathcal{L}_\Lambda \langle \mathcal{J}_\alpha^- | \otimes T^\alpha + \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \otimes (\eta_{-1\alpha\beta} T^\beta + \eta_{1\alpha\beta} \rho^{-2} \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}), \\
 \delta_\Lambda \langle \mathcal{J}_\alpha^- | &= \mathcal{L}_\Lambda \langle \mathcal{J}_\alpha^- | + \eta_{-1\alpha\beta} \langle \partial_\Lambda | (T^\beta + \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}) | \Lambda \rangle \langle \partial_\Lambda |, \quad (4.30)
 \end{aligned}$$

where the non-covariant terms were recombined using (2.19) and (4.24) in the first and second line, respectively. Due to the E_9 covariance (4.14) of the shifted current, its Lie derivative simply reads

$$\begin{aligned}
 \mathcal{L}_\Lambda \langle \mathcal{J}_\alpha^- | \otimes T^\alpha &= (\langle \partial_{\mathcal{J}} | \Lambda \rangle \langle \mathcal{J}_\alpha^- | + \langle \mathcal{J}_\alpha^- | \Lambda \rangle \langle \partial_\Lambda | - \langle \partial_\Lambda | \Lambda \rangle \langle \mathcal{J}_\alpha^- |) \otimes T^\alpha \\
 &\quad - \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \mathcal{J}_\gamma^- | \otimes [T^\beta, T^\gamma], \quad (4.31)
 \end{aligned}$$

and matches that of the regular current up to a density term.

Having established the necessary transformation rules of the various fields under generalised diffeomorphisms, we now move on to proving the invariance of the potential (4.1) at $\tilde{\rho} = 0$. Its variation takes the form

$$\delta_\Lambda V = \mathcal{L}_\Lambda V + \Delta_\Lambda V = \langle \partial | (|\Lambda\rangle V) + \Delta_\Lambda V. \quad (4.32)$$

As was shown in (4.16), the potential is an E_9 scalar of weight one. As a result, we immediately deduce that the generalised Lie derivative of the potential is a total derivative. In the following we then exclusively focus on the non-covariant variations $\Delta_\Lambda V$. From (4.20), (4.22) and (4.25) one computes that

$$\Delta_\Lambda \mathcal{L}_1 = 4\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle, \quad (4.33)$$

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_4 = & -2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\ & - \rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle J_0 | \mathcal{M}^{-1} | \partial_\Lambda \rangle + \rho^{-1} \langle J_0 | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \partial_\Lambda \rangle, \end{aligned} \quad (4.34)$$

where we used (3.26) in the variation of \mathcal{L}_1 and the section constraint (3.18a) in the variation of \mathcal{L}_4 . For the second term in the potential, we find

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_2 = & 2\rho^{-1} \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\gamma \mathcal{M}^{-1} (T^{\beta\dagger} + \mathcal{M} T^\beta \mathcal{M}^{-1}) | \mathcal{J}_\gamma \rangle \\ = & 2\rho^{-1} \langle \mathcal{J}_\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \partial_\Lambda \rangle - 2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\ & + 2\rho^{-1} \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle. \end{aligned} \quad (4.35)$$

The section constraint (3.18a) was used on both terms to simplify the first line. Using (4.30), the variation of \mathcal{L}_3 reads

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_3 = & 2\rho^{-1} \eta_{1\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\gamma T^\beta \mathcal{M}^{-1} | \mathcal{J}_\gamma^- \rangle \\ = & 2\rho^{-1} \eta_{1\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma^- \rangle \\ = & 2\rho^{-1} \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [\mathcal{S}_{-1}(T^\gamma), \mathcal{S}_1(T^\beta)] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle \\ = & 2\rho^{-1} \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle \\ & - 2\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle + 2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle. \end{aligned} \quad (4.36)$$

The section constraints (3.18b) and (3.18c) were used in the first and second line, respectively, and in the last line we used that for α and β restricted to \mathfrak{e}_9 one has

$$[\mathcal{S}_{-1}(T^\alpha), \mathcal{S}_1(T^\beta)] = [T^\alpha, T^\beta] - \eta^{\alpha\beta} \mathbf{K} - \delta_0^\alpha T^\beta - \delta_0^\beta T^\alpha, \quad (4.37)$$

with the Kronecker symbol defined such that $\delta_0^\alpha \langle \mathcal{J}_\alpha | = \langle J_0 |$, as well as the section constraint (3.18a).

Combining the above non-covariant variations, we find

$$\begin{aligned} \Delta_\Lambda V = & \frac{1}{2} \rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | (2T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle - \mathcal{M}^{-1} | J_0 \rangle) \\ & - \rho^{-1} \langle \mathcal{J}_\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \partial_\Lambda \rangle + \frac{1}{2} \rho^{-1} \langle J_0 | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \partial_\Lambda \rangle \end{aligned} \quad (4.38)$$

which, upon using (3.24) and (4.3), reduces to a total derivative

$$\Delta_\Lambda V = \langle \partial | (\rho^{-1} |\Lambda\rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \partial_\Lambda\rangle - \rho^{-1} \mathcal{M}^{-1} | \partial_\Lambda\rangle \langle \partial_\Lambda | \Lambda) \rangle. \quad (4.39)$$

This proves that the potential (4.1) is invariant, at $\tilde{\rho} = 0$, under generalised diffeomorphisms up to total derivatives.

4.2 The potential at $\tilde{\rho} \neq 0$

We shall now present the general expression of the potential (4.1) at $\tilde{\rho} \neq 0$. In this case, $\mathcal{M} \in \hat{E}_8 \rtimes \text{SL}(2)$ and the various terms read

$$\begin{aligned} \mathcal{L}_1 = \rho^{-1} \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | \mathcal{M}^{-1} | J_B^n \rangle \eta^{AB} - 2 \rho^{-1} \langle J_0 + 2 \tilde{\rho} J_1 | \mathcal{M}^{-1} | J_K \rangle \\ - 2 \rho^{-1} \langle J_1 | \mathcal{M}^{-1} | 2 \rho^2 \chi - \rho^2 J_1 + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle, \end{aligned} \quad (4.40a)$$

$$\mathcal{L}_2 = \rho^{-1} \langle \mathcal{J}_\alpha | T^\beta \mathcal{M}^{-1} T^{\alpha \dagger} | \mathcal{J}_\beta \rangle, \quad (4.40b)$$

$$\mathcal{L}_3 = \rho \langle \mathcal{J}_\alpha^- | T^\beta \mathcal{M}^{-1} T^{\alpha \dagger} | \mathcal{J}_\beta^- \rangle, \quad (4.40c)$$

$$\mathcal{L}_4 = \rho^{-1} \langle J_0 + 2 \tilde{\rho} J_1 | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle. \quad (4.40d)$$

The currents components $\langle \mathcal{J}_\alpha |$ defined in (3.24) are now non-vanishing along $\mathfrak{e}_8 \oplus \mathfrak{sl}(2)$, while $\langle \mathcal{J}_\alpha^- |$ defined in (3.33) has non-vanishing components along all Virasoro generators L_n with $n \leq 1$ in \mathfrak{f} . The $\mathfrak{sl}(2)$ components of the current are the same as for $\text{SL}(2)/\text{SO}(1,1)$

$$\langle J_0 | = -2 \rho^{-1} \langle \partial_\rho | \rho + 2 \rho^{-2} \tilde{\rho} \langle \partial_{\tilde{\rho}} | \tilde{\rho}, \quad (4.41a)$$

$$\langle J_{-1} | = 2 \rho^{-1} \tilde{\rho} \langle \partial_\rho | \rho - (1 + \rho^{-2} \tilde{\rho}^2) \langle \partial_{\tilde{\rho}} | \tilde{\rho}, \quad (4.41b)$$

$$\langle J_1 | = -\rho^{-2} \langle \partial_{\tilde{\rho}} | \tilde{\rho}. \quad (4.41c)$$

It is straightforward to check that these components satisfy the identity (3.31). The function $\Omega^\alpha(\mathcal{M})$ defined in (A.28) is a combination of E_9 group cocycles that reduces to the cocycle $\omega_1^\alpha(\mathcal{M})$ when $\tilde{\rho}$ is set to zero. In the following, we will show that each term is invariant under rigid $\hat{E}_8 \rtimes \mathbb{R}_{L_{-1}}$ and scales with weight one under rigid \mathbb{R}_d^+ . Then we will show that the combination (4.1) is invariant under Λ and Σ generalised diffeomorphisms. At this point, the most striking difference with the expression of the potential at $\tilde{\rho} = 0$ is perhaps the complexity of the expression of \mathcal{L}_1 , and the fact that its rigid \hat{E}_8 invariance is not manifest. This is due to the absence of an invariant bilinear form over $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$.

4.2.1 Rigid $\hat{E}_8 \rtimes (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ symmetry

For clarity, we will treat separately the transformations under E_9 and $\mathbb{R}_{L_{-1}}$. Under $g \in E_9$, the derivatives, \mathcal{M} and the field ρ still transform as in (4.5), (4.4) and (4.8), respectively. From the parameterisation of the coset element (3.7), one finds that

$$\tilde{\rho} \rightarrow \rho(g)^2 \tilde{\rho}. \quad (4.42)$$

The transformation of the current and its components still take the same form as in the $\tilde{\rho} = 0$ case

$$\langle \mathcal{J}_\alpha | \otimes T^\alpha \rightarrow \rho(g)^2 \langle \mathcal{J}_\alpha | g \otimes g^{-1} T^\alpha g, \quad (4.43)$$

$$\langle \mathcal{J}_\alpha | \rightarrow \rho(g)^2 R(g)^\beta_\alpha \langle \mathcal{J}_\beta | g, \quad (4.44)$$

but now the current is $\mathfrak{e}_8 \oplus \mathfrak{sl}(2)$ -valued. The adjoint representation matrix $R(g)^\alpha_\beta$ is still defined from the generators of \mathfrak{f} by (4.9). Note also that, using (4.41), the combination appearing in \mathcal{L}_4 reads

$$\langle J_0 + 2\tilde{\rho} J_1 | = -2\rho^{-1} \langle \partial_\rho | \rho, \quad (4.45)$$

and therefore simply transforms as $\langle J_0 + 2\tilde{\rho} J_1 | \rightarrow \rho(g)^2 \langle J_0 + 2\tilde{\rho} J_1 | g$ under E_9 .

The computation of the variation of the shifted current relies on a similar reasoning as for $\tilde{\rho} = 0$. We start with the infinite series of shift operators in the expression of the shifted current (3.33), that transforms under E_9 as

$$\langle \mathcal{J}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha) \rightarrow \rho(g)^2 \langle \mathcal{J}_\alpha | g \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k (\rho(g)^{-2} g^{-1} \mathcal{S}_{-1-k}(T^\alpha) g - \rho(g)^{2k} \omega_{1+k}^\alpha(g^{-1}) \mathbf{K}), \quad (4.46)$$

where we used (A.13). Each cocycle $\omega_k^\alpha(g)$, together with the character $\rho(g)$, defines an extension of the adjoint representation of E_9 by the generator L_{-k} . Once again we choose the variation of the field $\langle \chi |$ such that the shifted current transforms covariantly under E_9 ,

$$\langle \chi | \rightarrow \langle \chi | g + \sum_{k=0}^{\infty} \tilde{\rho}^k \rho(g)^{2k+2} \omega_{1+k}^\alpha(g^{-1}) \langle \mathcal{J}_\alpha | g. \quad (4.47)$$

Indeed, it follows from (4.46) and (4.47) that the shifted current transforms as

$$\langle \mathcal{J}_\alpha^- | \otimes T^\alpha \rightarrow \langle \mathcal{J}_\alpha^- | g \otimes g^{-1} T^\alpha g, \quad (4.48)$$

and admits non-zero components along all Virasoro generators L_n for $n \leq 1$. Using the above results, it is straightforward to check that under E_9 , the terms \mathcal{L}_2 , \mathcal{L}_3 and \mathcal{L}_4 only scale by a factor $\rho(g)^2$. The term \mathcal{L}_1 is more complicated, and we will only prove its invariance under infinitesimal $\hat{E}_8 \times \mathbb{R}_{L_{-1}}$ transformations. We postpone this task to first prove the invariance of \mathcal{L}_2 , \mathcal{L}_3 and \mathcal{L}_4 under $\mathbb{R}_{L_{-1}}$ and come back to \mathcal{L}_1 afterwards.

Under $e^{X-L_{-1}} \in \mathbb{R}_{L_{-1}}$, we have the following transformations

$$\begin{aligned} \mathcal{M} &\rightarrow e^{X-L_1} \mathcal{M} e^{X-L_{-1}}, \\ \langle \partial | &\rightarrow \langle \partial | e^{X-L_{-1}}, \\ \rho &\rightarrow \rho, \\ \tilde{\rho} &\rightarrow \tilde{\rho} - X_- \\ \langle \chi | &\rightarrow \langle \chi | e^{X-L_{-1}}. \end{aligned} \quad (4.49)$$

This implies

$$\langle \mathcal{J}_\alpha | \otimes T^\alpha \rightarrow \langle \mathcal{J}_\alpha | e^{X-L-1} \otimes e^{-X-L-1} T^\alpha e^{X-L-1}, \quad (4.50a)$$

$$\langle \mathcal{J}_\alpha^- | \otimes T^\alpha \rightarrow \langle \mathcal{J}_\alpha^- | e^{X-L-1} \otimes e^{-X-L-1} T^\alpha e^{X-L-1}. \quad (4.50b)$$

The covariance of the term involving the infinite series of shift operators in the expression of the shifted current $\langle \mathcal{J}_\alpha^- |$ can be verified using

$$\langle \mathcal{J}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha) = \langle \mathcal{J}_\alpha | \otimes e^{\tilde{\rho}L-1} \mathcal{S}_{-1}(e^{-\tilde{\rho}L-1} T^\alpha e^{\tilde{\rho}L-1}) e^{-\tilde{\rho}L-1}. \quad (4.51)$$

which follows from (A.30). The \mathbb{R}_{L-1} invariance of \mathcal{L}_2 , \mathcal{L}_3 and \mathcal{L}_4 is then a direct consequence of (4.49), (4.50a) and (4.50b).

Let us finally consider the transformation of \mathcal{L}_1 under E_9 and \mathbb{R}_{L-1} . As mentioned previously, this is more easily tackled by considering the infinitesimal variations of the current components. Under the infinitesimal variation with parameter X

$$X = X_K \mathbf{K} + X_0 \mathbf{d} + \sum_{n \in \mathbb{Z}} X_A^n T_n^A + X_{-L-1}, \quad (4.52)$$

one has

$$\delta_X \mathcal{M}^{-1} = -X_\alpha T^\alpha \mathcal{M}^{-1} - \mathcal{M}^{-1} X_\alpha T^{\alpha\dagger}, \quad \delta_X \langle \partial | = \langle \partial | (X_\alpha T^\alpha - X_0), \quad (4.53)$$

with by definition

$$X_\alpha T^\alpha = X_K \mathbf{K} + X_0 L_0 + \sum_{n \in \mathbb{Z}} X_A^n T_n^A + X_{-L-1}. \quad (4.54)$$

One obtains for the current components

$$\begin{aligned} \delta_X \langle J_A^n | &= \sum_{m \in \mathbb{Z}} X_B^{n-m} \langle J_C^m | f^{BC}{}_A - n X_A^n \langle J_0 | + (n-1) X_0 \langle J_A^n | \\ &\quad - (n+1) X_A^{n+1} \langle J_{-1} | - (n-1) X_A^{n-1} \langle J_1 | + (n+1) X_- \langle J_A^{n+1} | + \langle J_A^n | X_\alpha T^\alpha, \\ \delta_X \langle J_K | &= \sum_{m \in \mathbb{Z}} m \eta^{AB} X_A^{-m} \langle J_B^m | - X_0 \langle J_K | + \langle J_K | X_\alpha T^\alpha, \\ \delta_X \langle J_0 | &= -X_0 \langle J_0 | + 2 X_- \langle J_1 | + \langle J_0 | X_\alpha T^\alpha, \\ \delta_X \langle J_{-1} | &= -2 X_0 \langle J_{-1} | + X_- \langle J_0 | + \langle J_{-1} | X_\alpha T^\alpha, \\ \delta_X \langle J_1 | &= \langle J_1 | X_\alpha T^\alpha. \end{aligned} \quad (4.55)$$

The last term in $X_\alpha T^\alpha$ of each expression comes from the expansion of $g \in E_9$ and e^{X-L-1} acting on the derivative bra of $\langle \mathcal{J}_\alpha |$ in the $\overline{R(\Lambda_0)_0}$ representation as in (4.44) and (4.50a). These contributions trivially cancel the variation (4.53) of \mathcal{M}^{-1} . All the other terms follow from the linearisation of the character $\rho(g)$ and the adjoint representation matrix $R(g)^{\alpha\beta}$

defined in (A.22a) and (4.9), respectively. For the infinitesimal variation of the constrained field $\langle \chi |$ under E_9 and $\mathbb{R}_{L_{-1}}$, we obtain from (4.47) and (4.49),

$$\delta_X \langle \chi | = \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k (n-1-k) \eta^{AB} X_A^{1+k-n} \langle J_B^n | + \langle \chi | X_\alpha T^\alpha, \quad (4.56)$$

using the linearisation (A.14) of the cocycles $\omega_n^\alpha(g)$. With (4.55), we find that the infinitesimal variation of the first two terms of \mathcal{L}_1 gives

$$\begin{aligned} \delta_X \left(\rho^{-1} \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | \mathcal{M}^{-1} | J_B^n \rangle \eta^{AB} - 2 \rho^{-1} \langle J_0 + 2 \tilde{\rho} J_1 | \mathcal{M}^{-1} | J_K \rangle \right) \\ = -2 \rho^{-1} \sum_{n \in \mathbb{Z}} \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \rho^2 (k-n+1) X_A^{k+1-n} - (n+1) X_A^{-n-1} + n \tilde{\rho} X_A^{-n} \right) \langle J_B^n | \mathcal{M}^{-1} | J_1 \rangle \eta^{AB} \\ - X_0 \left(\rho^{-1} \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | \mathcal{M}^{-1} | J_B^n \rangle \eta^{AB} - 2 \rho^{-1} \langle J_0 + 2 \tilde{\rho} J_1 | \mathcal{M}^{-1} | J_K \rangle \right), \end{aligned} \quad (4.57)$$

where we used (3.31) to eliminate all the dependence on the components $\langle J_0 |$ and $\langle J_{-1} |$ in the second line. Note in particular the invariance of the above combination under $\mathbb{R}_{L_{-1}}$. For the remaining term in \mathcal{L}_1 , we need to consider the variation of $\Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha |$. We start from the expression

$$\Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | \otimes \mathbb{K} = \langle \mathcal{J}_\alpha | \otimes \left(\rho^2 \mathcal{M}^{-1} \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha)^\dagger \mathcal{M} + \tilde{\rho} T^\alpha - \mathcal{S}_1(T^\alpha) - \tilde{\rho} \delta_{\mathbb{K}}^\alpha \mathbb{K} \right), \quad (4.58)$$

which is obtain by using (A.28) and (3.26). With (4.51), one can show that the above combination transforms as

$$\Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | \rightarrow \Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | e^{X_{-L_{-1}}}, \quad (4.59)$$

under $\mathbb{R}_{L_{-1}}$, while under $g \in E_9$ one finds

$$\begin{aligned} \Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | \rightarrow \rho(g)^4 \Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | g - \sum_{k=0}^{\infty} \tilde{\rho}^k \rho(g)^{2k+6} \omega_{1+k}^\alpha(g^{-1}) \langle \mathcal{J}_\alpha | g \\ + \rho(g)^2 \omega_{-1}(g^{-1}) \langle \mathcal{J}_\alpha | g - \tilde{\rho} \rho(g)^4 R(g)^\alpha \mathbb{K} \langle \mathcal{J}_\alpha | g, \end{aligned} \quad (4.60)$$

using (A.13). By linearising (4.59) and (4.60) using (A.14), we find the following infinitesimal variation

$$\begin{aligned} \delta_X \left(\Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | \right) = - \sum_{n \in \mathbb{Z}} \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \rho^2 (n-1-k) X_B^{k+1-n} - (n+1) X_B^{-n-1} + n \tilde{\rho} X_B^{-n} \right) \eta^{AB} \langle \mathcal{J}_\alpha | \\ - 2 X_0 \Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | + \Omega^\alpha(\mathcal{M}) \langle \mathcal{J}_\alpha | X_\beta T^\beta. \end{aligned} \quad (4.61)$$

With this result and (4.56), it is easy to verify that the last term of \mathcal{L}_1 is $\mathbb{R}_{L_{-1}}$ invariant, while its \hat{E}_8 variation cancels out that of the first two terms in (4.57). We are then left with

$$\delta_X \mathcal{L}_1 = -X_0 \mathcal{L}_1, \quad (4.62)$$

which is the action of \mathfrak{d} on \mathcal{L}_1 . The full potential is then $\hat{E}_8 \times \mathbb{R}_{L_{-1}}$ invariant and, as in the $\tilde{\rho} = 0$ case (4.16), uniformly scales by a factor $\rho(g)^2$ under finite transformations generated by \mathfrak{d} .

4.2.2 Invariance under generalised diffeomorphisms

The fields $\mathcal{M} \in \hat{E}_8 \times \text{SL}(2)$ and ρ still transform covariantly under generalised diffeomorphisms, i.e. as in (4.18) and (4.19), respectively. According to (4.42), the field $\tilde{\rho}$ is an E_9 scalar density of weight one and thus transforms as a total derivative,

$$\delta_\Lambda \tilde{\rho} = \mathcal{L}_\Lambda \tilde{\rho} = \langle \partial | (|\Lambda\rangle \tilde{\rho}) \rangle. \quad (4.63)$$

The $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ -valued current $\langle \mathcal{J}_\alpha | \otimes T^\alpha$ still transforms as in (4.20) and (4.21). In particular, its non-covariant variation still reads

$$\Delta_\Lambda \langle \mathcal{J}_\alpha | \otimes T^\alpha = \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \otimes (T^\beta + \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}). \quad (4.64)$$

However, the bilinear form $\eta_{\alpha\beta}$ is not invariant under the extended group $\hat{E}_8 \times \text{SL}(2)$, so that the non-covariant variation of the current components, according to (A.32), is then

$$\Delta_\Lambda \langle \mathcal{J}_\alpha | = \langle \partial_\Lambda | \left(\eta_{\alpha\beta} T^\beta + \left[\frac{\tilde{\rho}}{\rho^2} \eta_{1\alpha\beta} + \left(1 - 2 \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{\alpha\beta} - \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{-1\alpha\beta} \right] \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_\Lambda |. \quad (4.65)$$

The Lie derivative of the current components is still given by (4.23), although the structure constant indices now take values over $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$.

Let us now turn to the variation of the shifted current. Using (2.19) and (4.65), one obtains that the non-covariant variation of the shifted current $\langle \mathcal{J}_\alpha^- |$ defined in (3.33) for $\alpha \neq \mathbf{K}$, transforms as

$$\begin{aligned} \Delta_\Lambda \langle \mathcal{J}_\alpha^- |_{\alpha \neq \mathbf{K}} &= \sum_{k=0}^{\infty} \tilde{\rho}^k \langle \partial_\Lambda | \left(\eta_{-1-k\alpha\beta} T^\beta \right. \\ &\quad \left. + \left[\frac{\tilde{\rho}}{\rho^2} \eta_{-k\alpha\beta} + \left(1 - 2 \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{-1-k\alpha\beta} - \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{-2-k\alpha\beta} \right] \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_\Lambda | \\ &= \langle \partial_\Lambda | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{-1-k\alpha\beta} T^\beta + \left[\frac{\tilde{\rho}}{\rho^2} \eta_{\alpha\beta} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{-1\alpha\beta} \right] \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_\Lambda |. \end{aligned} \quad (4.66)$$

It is therefore natural to define the non-covariant variation of $\langle \chi |$ such that this transformation rule also applies for $\alpha = \mathbf{K}$. We then choose

$$\Delta_\Lambda \langle \chi | = - \langle \partial_\Lambda | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k L_{-1-k} + \mathcal{M}^{-1} \left[\frac{\tilde{\rho}}{\rho^2} L_0 + \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_1 \right] \mathcal{M} \right) | \Lambda \rangle \langle \partial_\Lambda | - \frac{\tilde{\rho}}{\rho^2} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda |. \quad (4.67)$$

The presence of the last term does not follow from the previous argument, but we observe that it could be removed by redefining $\langle \chi | \rightarrow \langle \chi | - \langle J_1 |$, since $\Delta_\Lambda \langle J_1 | = - \frac{\tilde{\rho}}{\rho^2} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda |$. Such a redefinition would not modify the rigid transformation

rule of $\langle \chi |$ under $\hat{E}_8 \times (\mathbb{R}_d^+ \times \mathbb{R}_{L-1})$, but would lead to a slightly less compact expression for the potential. The Lie derivative of $\langle \chi |$ follows from its infinitesimal variation (4.56) under E_9 and reads

$$\mathcal{L}_\Lambda \langle \chi | = \langle \partial_\chi | \Lambda \rangle \langle \chi | + \langle \chi | \Lambda \rangle \langle \partial_\Lambda | - \langle \partial_\Lambda | \Lambda \rangle \langle \chi | + \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} (n-1-k) \langle \partial_\Lambda | T_{n-1-k}^A | \Lambda \rangle \langle J_A^n |. \quad (4.68)$$

Combining (4.66) and (4.67) and using (A.32), one finds for the shifted current

$$\begin{aligned} \Delta_\Lambda \langle \mathcal{J}_\alpha^- | &= \sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{-1-k} \alpha\beta \langle \partial_\Lambda | T^\beta | \Lambda \rangle \langle \partial_\Lambda | - \delta_\alpha^\kappa \frac{\tilde{\rho}}{\rho^2} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \\ &\quad + \left(\frac{\tilde{\rho}}{\rho^2} \eta_{\alpha\beta} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) \eta_{-1} \alpha\beta \right) \langle \partial_\Lambda | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} | \Lambda \rangle \langle \partial_\Lambda | \\ \Delta_\Lambda \langle \mathcal{J}_\alpha^- | \otimes T^\alpha &= \sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{-1-k} \alpha\beta \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \otimes T^\beta - \frac{\tilde{\rho}}{\rho^2} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \otimes \mathbb{K} \\ &\quad + \frac{1}{\rho^2} (\eta_{1\alpha\beta} - \tilde{\rho} \eta_{\alpha\beta}) \langle \partial_\Lambda | T^\beta | \Lambda \rangle \langle \partial_\Lambda | \otimes \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}, \end{aligned} \quad (4.69)$$

while its Lie derivative is still given by the expression (4.31).

Let us now consider the variation of each term in the potential. Just as in the $\tilde{\rho} = 0$ case, we will only focus on the non-covariant variations $\Delta_\Lambda V$ as the Lie derivative of the potential reduces to a total derivative. This is ensured by the fact that, as proved in section 4.2.1, the potential transforms as an E_9 scalar of weight one. For \mathcal{L}_2 and \mathcal{L}_4 , the computation is the same as in the $\tilde{\rho} = 0$ case and one finds

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_2 &= 2\rho^{-1} \langle \mathcal{J}_\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \partial_\Lambda \rangle - 2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\ &\quad + 2\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle \eta_{\alpha\beta}, \end{aligned} \quad (4.70)$$

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_4 &= -2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\ &\quad - \rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle J_0 + 2\tilde{\rho} J_1 | \mathcal{M}^{-1} | \partial_\Lambda \rangle + \rho^{-1} \langle J_0 + 2\tilde{\rho} J_1 | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \partial_\Lambda \rangle. \end{aligned} \quad (4.71)$$

For \mathcal{L}_3 , one obtains that the first infinite sum in (4.69) gives terms that all vanish according to the section constraints, while the others give

$$\begin{aligned} \Delta_\Lambda \mathcal{L}_3 &= 2\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | T^\gamma T^\beta \mathcal{M}^{-1} | \mathcal{J}_\gamma^- \rangle (\eta_{1\alpha\beta} - \tilde{\rho} \eta_{\alpha\beta}) \\ &\quad - 2\rho^{-1} \tilde{\rho} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle \\ &= 2\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma^- \rangle (\eta_{1\alpha\beta} - \tilde{\rho} \eta_{\alpha\beta}) \\ &\quad - 2\rho^{-1} \sum_{k=1}^{\infty} \tilde{\rho}^k \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \mathcal{S}_{-k}(T^\alpha) \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle - 2\rho^{-1} \tilde{\rho} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \chi \rangle \\ &= 2\rho^{-1} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle \eta_{\alpha\beta} \\ &\quad - 2\rho^{-1} \sum_{k=0}^{\infty} \tilde{\rho}^k \langle \partial_\Lambda | \mathcal{S}_{-k}(T^\alpha) | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\ &\quad + 2\rho^{-1} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle - 2\rho^{-1} \tilde{\rho} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \chi \rangle, \end{aligned} \quad (4.72)$$

where we used the following identity in the last step

$$\begin{aligned}
 & \sum_{k=0}^{\infty} \tilde{\rho}^k \langle \partial_{\Lambda} | T^{\alpha} | \Lambda \rangle \langle \partial_{\Lambda} | [\mathcal{S}_{-1-k}(T^{\gamma}), T^{\beta}] \mathcal{M}^{-1} | \mathcal{J}_{\gamma} \rangle (\eta_{1\alpha\beta} - \tilde{\rho} \eta_{\alpha\beta}) \\
 &= \sum_{k=0}^{\infty} \tilde{\rho}^k \left(\langle \partial_{\Lambda} | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{S}_{-k}(T^{\alpha}) \mathcal{M}^{-1} | \mathcal{J}_{\alpha} \rangle - \langle \partial_{\Lambda} | \mathcal{S}_{-k}(T^{\alpha}) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | \mathcal{J}_{\alpha} \rangle \right) \\
 & \quad + \langle \partial_{\Lambda} | T^{\alpha} | \Lambda \rangle \langle \partial_{\Lambda} | [T^{\gamma}, T^{\beta}] \mathcal{M}^{-1} | \mathcal{J}_{\gamma} \rangle \eta_{\alpha\beta}. \tag{4.73}
 \end{aligned}$$

Let us finally consider the variation of \mathcal{L}_1 . Using (4.65) one computes that

$$\Delta_{\Lambda} \sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | = \eta_{AB} \langle \partial_{\Lambda} | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k T_{n-k}^B + \frac{\tilde{\rho}}{\rho^2} \mathcal{M}^{-1} T_{1+n}^{B\dagger} \mathcal{M} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right) \mathcal{M}^{-1} T_n^{B\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_{\Lambda} |, \tag{4.74}$$

which allows computing the variation of the first term in $\rho\mathcal{L}_1$ as

$$\begin{aligned}
 & \Delta_{\Lambda} \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | \mathcal{M}^{-1} | J_B^n \rangle \eta^{AB} \right) \\
 &= 2 \langle \partial_{\Lambda} | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-k}(T^{\alpha}) + \frac{\tilde{\rho}}{\rho^2} \mathcal{M}^{-1} \mathcal{S}_1(T^{\alpha})^{\dagger} \mathcal{M} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right) \mathcal{M}^{-1} T^{\alpha\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | \mathcal{J}_{\alpha} \rangle \\
 & \quad - 2 \sum_{q=-1,0,1} \langle \partial_{\Lambda} | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-k}(L_q) + \frac{\tilde{\rho}}{\rho^2} \mathcal{M}^{-1} \mathcal{S}_1(L_q)^{\dagger} \mathcal{M} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right) \mathcal{M}^{-1} L_q^{\dagger} \mathcal{M} \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_q \rangle \\
 & \quad - 2 \langle \partial_{\Lambda} | \left(2 - \frac{\tilde{\rho}^2}{\rho^2} \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_K \rangle \\
 &= 2 \langle \partial_{\Lambda} | \left(2 \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-k}(T^{\alpha}) - \frac{\tilde{\rho}}{\rho^2} \Omega^{\alpha}(\mathcal{M}) \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | \mathcal{J}_{\alpha} \rangle - 4 \langle \partial_{\Lambda} | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_K \rangle \\
 & \quad - 2 \sum_{q=-1,0,1} \langle \partial_{\Lambda} | \left(\sum_{k=0}^{\infty} \tilde{\rho}^k L_{q-k} + \mathcal{M}^{-1} \left[\frac{\tilde{\rho}}{\rho^2} L_{-1-q} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right) L_{-q} \right] \mathcal{M} \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_q \rangle. \tag{4.75}
 \end{aligned}$$

In the first step, the loop valued currents were completed to the full $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ currents by adding and subtracting explicitly the missing components. In the second step, we used (3.28) and substituted $\mathcal{S}_1(T^{\alpha})$ by its expression following from (4.58). Using once again (4.65), the variation of the second term in $\rho\mathcal{L}_1$ gives

$$\begin{aligned}
 & \Delta_{\Lambda} \left(-2 \langle J_0 + 2\tilde{\rho} J_1 | \mathcal{M}^{-1} | J_K \rangle \right) \\
 &= 2 \langle \partial_{\Lambda} | \left(L_0 + \mathcal{M}^{-1} \left[\frac{\tilde{\rho}}{\rho^2} L_{-1} + \left(1 - 2\frac{\tilde{\rho}^2}{\rho^2}\right) L_0 - \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right) L_1 \right] \mathcal{M} \right) | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_0 + 2\tilde{\rho} J_1 \rangle \\
 & \quad + 4 \langle \partial_{\Lambda} | \Lambda \rangle \langle \partial_{\Lambda} | \mathcal{M}^{-1} | J_K \rangle. \tag{4.76}
 \end{aligned}$$

For the variation of the last term in \mathcal{L}_1 , we need that

$$\Delta_{\Lambda} \Omega^{\alpha}(\mathcal{M}) = 0, \tag{4.77}$$

which follows from the fact $\Omega^\alpha(\mathcal{M})$ is a function of \mathcal{M} and not of its derivative. Using (A.40), one can also show that (4.67) can be rewritten as

$$\begin{aligned} \Delta_\Lambda \langle \chi | &= - \sum_{k=0}^{\infty} \tilde{\rho}^k \langle \partial_\Lambda | L_{-1-k} | \Lambda \rangle \langle \partial_\Lambda | \\ &\quad - \frac{1}{\rho^2} \langle \partial_\Lambda | \left(L_1 + \eta_{\alpha\beta} H(\mathcal{M})^\alpha{}_\gamma (\Omega^\gamma(\mathcal{M}) + \tilde{\rho} \delta_K^\gamma) T^\beta + \tilde{\rho} \mathbf{K} \right) | \Lambda \rangle \langle \partial_\Lambda | . \end{aligned} \quad (4.78)$$

Together with (4.65), this yields

$$\begin{aligned} &\Delta_\Lambda \left(-2 \langle J_1 | \mathcal{M}^{-1} | 2\rho^2 \chi - \rho^2 J_1 + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle \right) \\ &= 2 \frac{\tilde{\rho}}{\rho^2} \langle \partial_\Lambda | \Lambda \rangle \langle J_1 | \mathcal{M}^{-1} | 2\rho^2 \chi + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle \\ &\quad + 2 \langle \partial_\Lambda | \left(2\rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k L_{-1-k} + 2L_1 + \eta_{\alpha\beta} T^\beta (H(\mathcal{M})^\alpha{}_\gamma \Omega^\gamma(\mathcal{M}) - \Omega^\alpha(\mathcal{M})) \right) | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | J_1 \rangle \\ &\quad - 4\tilde{\rho} \langle \partial_\Lambda | \mathcal{M}^{-1} \left(\frac{\tilde{\rho}}{\rho^2} L_{-1} + \left(1 - 2\frac{\tilde{\rho}^2}{\rho^2} \right) L_0 - \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_1 \right) \mathcal{M} | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | J_1 \rangle \end{aligned} \quad (4.79)$$

By combining the contributions (4.75), (4.76) and (4.79), and using the relation (3.31) to eliminate $\langle J_0 |$ and $\langle J_{-1} |$, we find

$$\begin{aligned} &\Delta_\Lambda (\rho \mathcal{L}_1) \quad (4.80) \\ &= 4 \langle \partial_\Lambda | \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-k}(T^\alpha) | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle + 4\tilde{\rho} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \chi \rangle \\ &\quad + 2 \langle \partial_\Lambda | \left(L_1 + \tilde{\rho} L_0 + \rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k L_{-1-k} + \eta_{\alpha\beta} T^\beta (H(\mathcal{M})^\alpha{}_\gamma \Omega^\gamma(\mathcal{M}) - \Omega^\alpha(\mathcal{M})) \right. \\ &\quad \left. - \mathcal{M}^{-1} \left[\frac{\tilde{\rho}}{\rho^2} L_{-2} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_{-1} + \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_0 + \rho^2 \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right)^2 L_1 \right] \mathcal{M} \right) | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | J_1 \rangle \end{aligned}$$

Writing $\mathcal{M} = m \hat{g}_\mathcal{M}$, with $\hat{g}_\mathcal{M} \in \hat{E}_8$ and $m \in \text{SL}(2)$ as in (3.9), one computes that

$$\begin{aligned} &\mathcal{M}^{-1} \left(\frac{\tilde{\rho}}{\rho^2} L_{-2} + \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_{-1} + \tilde{\rho} \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right) L_0 + \rho^2 \left(1 - \frac{\tilde{\rho}^2}{\rho^2} \right)^2 L_1 \right) \mathcal{M} \\ &= \hat{g}_\mathcal{M}^{-1} \left(L_1 + \tilde{\rho} L_0 + \rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k L_{-1-k} \right) \hat{g}_\mathcal{M} . \end{aligned} \quad (4.81)$$

Together with the identities (A.35) and (A.37), this implies that the variation of \mathcal{L}_1 reduces to

$$\Delta_\Lambda \mathcal{L}_1 = 4\rho^{-1} \langle \partial_\Lambda | \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-k}(T^\alpha) | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle + 4\frac{\tilde{\rho}}{\rho} \langle \partial_\Lambda | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} | \chi \rangle . \quad (4.82)$$

With (4.70)–(4.72) and (4.82), one can easily check that just as in the $\tilde{\rho} = 0$ case, the non-covariant variations of (4.1) exactly recombine into the expression (4.38), such that the full potential at $\tilde{\rho} \neq 0$ is invariant under generalised diffeomorphisms up to total derivatives.

4.2.3 Invariance under Σ transformations

We conclude our proof of the gauge invariance of the potential by considering its variation under Σ generalised diffeomorphisms. We denote such variations simply by δ_Σ . The fields \mathcal{M} , ρ and $\tilde{\rho}$ transform covariantly

$$\delta_\Sigma \mathcal{M} = \mathcal{L}_\Sigma \mathcal{M} = \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) (\mathcal{M} T^\beta + T^{\beta\dagger} \mathcal{M}), \quad (4.83a)$$

$$\delta_\Sigma \rho = \mathcal{L}_\Sigma \rho = 0, \quad (4.83b)$$

$$\delta_\Sigma \tilde{\rho} = \mathcal{L}_\Sigma \tilde{\rho} = \text{Tr}(\Sigma). \quad (4.83c)$$

These expressions follow from the general definition (3.17) of the Lie derivative and the transformation properties (4.49) of the fields under rigid \mathbb{R}_{L-1} . We then deduce that

$$\delta_\Sigma \langle \mathcal{J}_\alpha | \otimes T^\alpha = \mathcal{L}_\Sigma \langle \mathcal{J}_\alpha | \otimes T^\alpha + \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \otimes (T^\beta + \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M}), \quad (4.84)$$

with

$$\mathcal{L}_\Sigma \langle \mathcal{J}_\alpha | \otimes T^\alpha = -\eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma | \otimes [T^\beta, T^\gamma]. \quad (4.85)$$

This implies the following variation for the current components

$$\delta_\Sigma \langle \mathcal{J}_\alpha | = \mathcal{L}_\Sigma \langle \mathcal{J}_\alpha | + \left(\eta_{-1\alpha\beta} \text{Tr}(T^\beta \Sigma) + \frac{1}{\rho^2} (\eta_{1\alpha\beta} - 2\tilde{\rho} \eta_{\alpha\beta} + \tilde{\rho}^2 \eta_{-1\alpha\beta}) \text{Tr}(\mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \Sigma) \right) \langle \partial_\Sigma |, \quad (4.86)$$

where we used the first identity in (A.32). Note that this implies $\Delta_\Sigma \langle J_0 + 2\tilde{\rho} J_1 | = 0$. The Lie derivative simply reads

$$\mathcal{L}_\Sigma \langle \mathcal{J}_\alpha | = -\eta_{-1\gamma\delta} \text{Tr}(T^\gamma \Sigma) f^{\delta\beta}{}_\alpha \langle \mathcal{J}_\beta |. \quad (4.87)$$

To derive the variation of the shifted current, we start by considering the Lie derivative of the term involving the infinite series of shift operators

$$\begin{aligned} \mathcal{L}_\Sigma \langle \mathcal{J}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha) &= -\eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k [T^\beta, \mathcal{S}_{-1-k}(T^\gamma)] \\ &\quad - \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k (n-1-k) \text{Tr}(T_{n-2-k}^A \Sigma) \langle J_A^n | \otimes \mathbb{K}. \end{aligned} \quad (4.88)$$

The Lie derivative of the shifted current follows from its covariance (4.50b) under rigid \mathbb{R}_{L-1} transformations

$$\mathcal{L}_\Sigma \langle \mathcal{J}_\alpha^- | \otimes T^\alpha = -\eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma^- | \otimes [T^\beta, T^\gamma]. \quad (4.89)$$

Together with (4.88), this implies that the Lie derivative of $\langle \chi |$ must be given by

$$\mathcal{L}_\Sigma \langle \chi | = \sum_{n \in \mathbb{Z}} \sum_{k=0}^{\infty} \tilde{\rho}^k (n-1-k) \text{Tr}(T_{n-2-k}^A \Sigma) \langle J_A^n |, \quad (4.90)$$

and which is indeed consistent with the infinitesimal action (4.56) of $\mathbb{R}_{L_{-1}}$ on $\langle \chi |$. In order to determine the non-covariant variation of $\langle \chi |$ we proceed as in the preceding section, and compute from (4.86) the non-covariant variation of the shifted current $\langle \mathcal{J}_\alpha^- |$ for $\alpha \neq \mathbb{K}$,

$$\Delta_\Sigma \langle \mathcal{J}_\alpha^- |_{\alpha \neq \mathbb{K}} = \sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{-2-k \alpha \beta} \text{Tr}(T^\beta \Sigma) \langle \partial_\Sigma | + \frac{1}{\rho^2} (\eta_{\alpha \beta} - \tilde{\rho} \eta_{-1 \alpha \beta}) \text{Tr}(\mathcal{M}^{-1} T^{\beta \dagger} \mathcal{M} \Sigma) \langle \partial_\Sigma |. \quad (4.91)$$

We define the non-covariant variation of the field $\langle \chi |$ as

$$\Delta_\Sigma \langle \chi | = -\frac{1}{\rho^2} \left(\rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k \text{Tr}(L_{-2-k} \Sigma) + \text{Tr}(\mathcal{M}^{-1} (L_0 - \tilde{\rho} L_1) \mathcal{M} \Sigma) \right) \langle \partial_\Sigma | - \frac{1}{\rho^2} \langle \partial_\Sigma | \Sigma, \quad (4.92)$$

such that the first two terms precisely reproduce the expression of the non-covariant variation (4.91), but for $\alpha = \mathbb{K}$. The last term turns out to be necessary for the closure of the algebra of generalised diffeomorphisms on $\langle \chi |$. Note that this expression is consistent with the section constraint since the parameter Σ is covariantly constrained on its right. It is important to mention, that by using (4.67) and (4.92), one can verify the closure of the gauge algebra on $\langle \chi |$. This is proven explicitly in appendix B. With the above results, we find that the non-covariant variation of the shifted current reads

$$\begin{aligned} \Delta_\Sigma \langle \mathcal{J}_\alpha^- | \otimes T^\alpha &= \sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{-2-k \alpha \beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \otimes T^\beta - \frac{1}{\rho^2} \langle \partial_\Sigma | \Sigma \otimes \mathbb{K} \\ &+ \frac{1}{\rho^2} (\eta_{\alpha \beta} - \tilde{\rho} \eta_{-1 \alpha \beta}) \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \otimes \mathcal{M}^{-1} T^{\beta \dagger} \mathcal{M}, \end{aligned} \quad (4.93)$$

where we used (A.33) for the last term.

With the various transformation properties derived above, we are now equipped to discuss the Σ variation of each term in the potential. The rigid $\hat{E}_8 \times \mathbb{R}_{L_{-1}}$ invariance of the potential, proven in section 4.2.1, directly implies that the variation generated by the Lie derivative vanishes, such that

$$\delta_\Sigma V = \mathcal{L}_\Sigma V + \Delta_\Sigma V = \Delta_\Sigma V. \quad (4.94)$$

In the following we therefore exclusively focus on the non-covariant variations of the various terms. For \mathcal{L}_2 and \mathcal{L}_4 , we obtain

$$\Delta_\Sigma \mathcal{L}_2 = 2 \rho^{-1} \eta_{-1 \alpha \beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle, \quad (4.95)$$

$$\Delta_\Sigma \mathcal{L}_4 = 0, \quad (4.96)$$

using the section constraint (3.18b). For the variation of \mathcal{L}_3 , we first compute that

$$\begin{aligned} &\eta_{\alpha \beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma | \otimes [\mathcal{S}_{-1-k}(T^\gamma), T^\beta] \\ &= \eta_{-1 \alpha \beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma | \otimes [\mathcal{S}_{-k}(T^\gamma), T^\beta] - \text{Tr}(\mathcal{S}_{-1-k}(T^\alpha) \Sigma) \langle \mathcal{J}_\alpha | \otimes \mathbb{K} \\ &+ \text{Tr}(\Sigma) \langle \mathcal{J}_\alpha | \otimes \mathcal{S}_{-1-k}(T^\alpha) - \sum_{q=-1,0,1} \eta_{q-1-k \alpha \beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_q | \otimes T^\beta, \end{aligned} \quad (4.97)$$

such that

$$\begin{aligned}
 & \eta_{\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma^- | \otimes [T^\gamma, T^\beta] \\
 &= \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma | \otimes [T^\gamma, T^\beta] + \tilde{\rho} \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\gamma^- | \otimes [T^\gamma, T^\beta] \\
 & \quad - \text{Tr}(T^\alpha \Sigma) \langle \mathcal{J}_\alpha^- | \otimes \mathbf{K} + \text{Tr}(\Sigma) \langle \mathcal{J}_\alpha^- | \otimes T^\alpha - \sum_{q=-1,0,1} \sum_{k=0}^{\infty} \tilde{\rho}^k \eta_{q-1-k} \alpha\beta \text{Tr}(T^\alpha \Sigma) \langle J_q | \otimes T^\beta .
 \end{aligned} \tag{4.98}$$

Using this formula one finds with (4.93),

$$\begin{aligned}
 \Delta_\Sigma \mathcal{L}_3 &= 2\rho^{-1} (\eta_{\alpha\beta} - \tilde{\rho} \eta_{-1\alpha\beta}) \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma^- \rangle \\
 & \quad - 2\rho^{-1} \text{Tr}(\Sigma) \langle \partial_\Sigma | T^\alpha \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle \\
 &= 2\rho^{-1} \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | [T^\gamma, T^\beta] \mathcal{M}^{-1} | \mathcal{J}_\gamma \rangle - 2\rho^{-1} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle \\
 & \quad - 2\rho^{-1} \langle \partial_\Sigma | \Sigma \mathcal{M}^{-1} | J_1 \rangle + 2\rho^{-1} \text{Tr}(\Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle ,
 \end{aligned} \tag{4.99}$$

where we used the section constraint on Σ and in particular

$$\eta_{\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | T^\beta = \langle \partial_\Sigma | \Sigma - \text{Tr}(\Sigma) \langle \partial_\Sigma | . \tag{4.100}$$

Let us finally consider the variation of $\rho \mathcal{L}_1$. For its first term, we obtain

$$\begin{aligned}
 & \Delta_\Sigma \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \langle J_A^{k-n} | \mathcal{M}^{-1} | J_B^n \rangle \eta^{AB} \right) \\
 &= 2 \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \text{Tr}(\mathcal{S}_{-1-k}(T^\alpha) \Sigma) + \frac{1}{\rho^2} \text{Tr} \left(\mathcal{M}^{-1} [\mathcal{S}_1(T^\alpha)^\dagger - \tilde{\rho} T^{\alpha\dagger} + \tilde{\rho} \delta_K^\alpha] \mathcal{M} \Sigma \right) \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | \mathcal{J}_\alpha \rangle \\
 & \quad - 2 \sum_{q=-1,0,1} \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \text{Tr}(L_{q-1-k} \Sigma) + \frac{1}{\rho^2} \text{Tr} \left(\mathcal{M}^{-1} [L_{-1-q} - \tilde{\rho} L_{-q}] \mathcal{M}^{-1} \Sigma \right) \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_q \rangle \\
 &= 4 \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle - \frac{2}{\rho^2} \langle \partial_\Sigma | \mathcal{M}^{-1} | 2\rho^2 \chi + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle \\
 & \quad - 2 \sum_{q=-1,0,1} \left(\sum_{k=0}^{\infty} \tilde{\rho}^k \text{Tr}(L_{q-1-k} \Sigma) + \frac{1}{\rho^2} \text{Tr} \left(\mathcal{M}^{-1} [L_{-1-q} - \tilde{\rho} L_{-q}] \mathcal{M} \Sigma \right) \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_q \rangle .
 \end{aligned} \tag{4.101}$$

In the first step we used (4.86) and subsequently completed the loop valued currents to their full $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ -valued expressions. In the second step, we used (4.58) to eliminate the term containing $\mathcal{S}_1(T^\alpha)$. Note that we wrote the result explicitly in terms of the shifted current and that the $\langle \chi |$ contributions in fact cancel.

For the variation of the second term in $\rho \mathcal{L}_1$, we get

$$\begin{aligned}
 & \Delta_\Sigma \left(-2 \langle J_0 + 2\tilde{\rho} J_1 | \mathcal{M}^{-1} | J_K \rangle \right) \\
 &= 2 \left(\text{Tr}(L_{-1} \Sigma) + \frac{1}{\rho^2} \text{Tr} \left(\mathcal{M}^{-1} [L_1 - 2\tilde{\rho} L_0 + \tilde{\rho}^2 L_1] \mathcal{M} \Sigma \right) \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_0 + 2\tilde{\rho} J_1 \rangle ,
 \end{aligned} \tag{4.102}$$

using (4.86). For computing the variation of the last term in $\rho\mathcal{L}_1$, we note that

$$\Delta_\Sigma \Omega^\alpha(\mathcal{M}) = 0, \quad (4.103)$$

since $\Omega^\alpha(\mathcal{M})$ does not depend on derivatives of \mathcal{M} . Together with (4.92), this leads to

$$\begin{aligned} & \Delta_\Sigma \left(-2 \langle J_1 | \mathcal{M}^{-1} | 2\rho^2 \chi - \rho^2 J_1 + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle \right) \\ &= \frac{2}{\rho^2} \text{Tr}(\Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | 2\rho^2(\chi - J_1) + \Omega^\alpha(\mathcal{M}) \mathcal{J}_\alpha \rangle + 4 \langle \partial_\Sigma | \Sigma \mathcal{M}^{-1} | J_1 \rangle \\ &+ 4 \left(\rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k \text{Tr}(L_{-2-k} \Sigma) + \text{Tr}(\mathcal{M}^{-1} [L_0 - \tilde{\rho} L_1] \mathcal{M} \Sigma) \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle \\ &- 2 \eta_{-1 \alpha \beta} (\Omega^\alpha(\mathcal{M}) + \Omega^\gamma(\mathcal{M}) H(\mathcal{M})^\alpha_\gamma) \text{Tr}(T^\beta \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle. \end{aligned} \quad (4.104)$$

Recombining the results (4.101), (4.102), (4.104) and eliminating the components $\langle J_0 |$ and $\langle J_{-1} |$ using (3.31), we reach

$$\begin{aligned} & \Delta_\Sigma(\rho\mathcal{L}_1) \quad (4.105) \\ &= 4 \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle + 4 \langle \partial_\Sigma | \Sigma \mathcal{M}^{-1} | J_1 \rangle - 4 \text{Tr}(\Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle \\ &- 2 \eta_{-1 \alpha \beta} (\Omega^\alpha(\mathcal{M}) + \Omega^\gamma(\mathcal{M}) H(\mathcal{M})^\alpha_\gamma) \text{Tr}(T^\beta \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle \\ &+ 2 \text{Tr} \left(\left[\rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k L_{-2-k} + \tilde{\rho} L_{-1} - L_0 \right. \right. \\ &\quad \left. \left. - \frac{1}{\rho^2} \mathcal{M}^{-1} (L_{-2} - 3\tilde{\rho} L_{-1} - (\rho^2 - 3\tilde{\rho}^2) L_0 + \tilde{\rho}(\rho^2 - \tilde{\rho}^2) L_1) \mathcal{M} \right] \Sigma \right) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle. \end{aligned}$$

Using once again the decomposition $\mathcal{M} = m \hat{g}_\mathcal{M}$, with $m \in \text{SL}(2)$ as in (3.9), one finds the following intermediary result

$$\begin{aligned} & - \frac{1}{\rho^2} \mathcal{M}^{-1} (L_{-2} - 3\tilde{\rho} L_{-1} - (\rho^2 - 3\tilde{\rho}^2) L_0 + \tilde{\rho}(\rho^2 - \tilde{\rho}^2) L_1) \mathcal{M} \\ &= \hat{g}_\mathcal{M}^{-1} \left(L_0 - \tilde{\rho} L_{-1} - \rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k L_{-2-k} \right) \hat{g}_\mathcal{M}, \end{aligned}$$

which, together with (A.41) and (A.42), allow reducing the expression (4.105) to

$$\Delta_\Sigma \mathcal{L}_1 = 4\rho^{-1} \left(\text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | \mathcal{J}_\alpha^- \rangle + \langle \partial_\Sigma | \Sigma \mathcal{M}^{-1} | J_1 \rangle - \text{Tr}(\Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} | J_1 \rangle \right) \quad (4.106)$$

With the above result and the variations (4.95), (4.96) and (4.99), it is straightforward to verify that the non-covariant variation of the potential (4.1) vanishes, thus proving its invariance under Σ generalised diffeomorphisms.

4.3 The potential in the unendlichbein formalism

In contrast to finite-dimensional Lie groups, care has to be taken when defining the Lie group from the algebra in the affine case. As the Lie algebra has infinitely many generators,

the formal exponential of Lie algebra elements does not manifestly form a group or it may introduce formal infinite sum expressions whose well-definedness needs to be established. For instance, the current $\mathcal{J}_M = \mathcal{M}^{-1}\partial_M\mathcal{M}$ suffers from this problem as \mathcal{M} depends on an infinity of dual potentials Y_n^A and applying Baker-Campbell-Hausdorff-type identities for evaluating the algebra-valued \mathcal{J}_M leads naively to infinite sums within each component along the loop algebra.

In this section, we explain how to make sense of this infinity and that our potential (4.1) is well-defined. As a preliminary step, we recall different definitions of an affine Kac-Moody group. The ‘minimal Kac-Moody group’ is defined as the closure of the group generated by the one-parameter subgroups of the real roots [45] that can also be interpreted using the Tits group functor [46]. This corresponds to the definition of the loop group \hat{E}_8 as the group of rational functions in E_8 of the spectral parameter w , that are meromorphic on $P^1(\mathbb{C})$. This group can be completed with respect to a topology of an associated building [47]. This ‘completed Kac-Moody group’ is then defined as the group of meromorphic functions in E_8 of the spectral parameter $w \in \mathbb{C}$. It corresponds to choosing one standard Borel subalgebra (in our case the negative Borel associated with negative powers of w) and allowing infinite analytic power series of w in that direction while keeping only a finite number of powers for the other direction (for us positive powers of w).

One can then write a representative \hat{V} of the coset space $\hat{E}_8/K(\hat{E}_8)$ for the completed group \hat{E}_8 in Borel gauge, using the Iwasawa decomposition. Putting explicit coordinates on the affine Borel group is subtle and is best done using building theory [48]. Here, we will choose coordinates formally through exponentiation as we did in (3.2) and we can also extend the coset representative to include the $\mathbb{R}_d^+ \times \mathbb{R}_{L-1}$ part for the axio-dilaton to have a group element $\mathcal{V} = v\hat{V}$, which we consider in the $R(\Lambda_0)_0$ representation as usual. In this way of writing \mathcal{V} one can see that when acting on an element of a lowest weight module, only finite expressions arise.

The Hermitian conjugation \mathcal{V}^\dagger does not preserve the Kac-Moody group completed in one direction as it interchanges the two standard Borel subgroups, so $\mathcal{M} = \mathcal{V}^\dagger\mathcal{V}$ is only well-defined when \mathcal{V} is in the minimal group. This is another way of seeing that the definition of \mathcal{J}_M requires qualification.

The advantage of working in the Borel gauge (3.2) is that the Maurer-Cartan derivative $\partial_M\mathcal{V}\mathcal{V}^{-1}$ is well-defined as it requires only finitely many commutators to determine the $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ element at each (negative) power of the spectral parameter. Similarly, the coset component of the Maurer-Cartan form

$$\mathcal{P}_M = \frac{1}{2}(\partial_M\mathcal{V}\mathcal{V}^{-1} + (\partial_M\mathcal{V}\mathcal{V}^{-1})^\dagger), \tag{4.107}$$

is well-defined. By construction, one can write the $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ current as

$$\mathcal{J}_M = 2\mathcal{V}^{-1}\mathcal{P}_M\mathcal{V}, \tag{4.108}$$

and this expression makes sense in the completed group. In the Fock space notation, this definition of the current takes the form

$$\langle \mathcal{J}_\alpha | \otimes T^\alpha = 2 \langle \mathcal{P}_\alpha | \otimes \mathcal{V}^{-1}T^\alpha\mathcal{V}, \quad \langle \mathcal{J}_\alpha | = 2 R(\mathcal{V})^\beta_\alpha \langle \mathcal{P}_\beta | = 2 R(v)^\beta_\gamma R(\hat{V})^\gamma_\alpha \langle \mathcal{P}_\beta |, \tag{4.109}$$

where we have define $R(\mathcal{V})^\alpha_\beta T^\beta = \mathcal{V}^{-1} T^\alpha \mathcal{V}$ in analogy with (A.7) in the appendix. In particular one has

$$\langle J_0 | = 2 \langle P_0 | - 4 \frac{\tilde{\rho}}{\rho} \langle P_1 |, \quad \langle J_1 | = 2 \rho^{-1} \langle P_1 |. \quad (4.110)$$

For the shifted current, one computes that

$$\begin{aligned} \langle \mathcal{J}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha) &= 2 \langle \mathcal{P}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(\hat{V}^{-1} v^{-1} T^\alpha v \hat{V}) \\ &= 2 \langle \mathcal{P}_\alpha | \otimes \sum_{k=0}^{\infty} \tilde{\rho}^k R(v)^\alpha_\beta (\hat{V}^{-1} \mathcal{S}_{-1-k}(T^\beta) \hat{V} - \omega_{1+k}^\beta(\hat{V}^{-1}) \mathbf{K}) \\ &= 2 \langle \mathcal{P}_\alpha | \otimes \left(\hat{V}^{-1} e^{\tilde{\rho} L_{-1}} \mathcal{S}_{-1}(\rho^{L_0} T^\alpha \rho^{-L_0}) e^{-\tilde{\rho} L_{-1}} \hat{V} - \sum_{k=0}^{\infty} \tilde{\rho}^k R(v)^\alpha_\beta \omega_{1+k}^\beta(\hat{V}^{-1}) \mathbf{K} \right) \\ &= 2 \langle \mathcal{P}_\alpha | \otimes \left(\rho^{-1} \mathcal{V}^{-1} \mathcal{S}_{-1}(T^\alpha) \mathcal{V} - \left(\omega_1^\alpha(\rho^{L_0}) + \sum_{k=0}^{\infty} \tilde{\rho}^k R(v)^\alpha_\beta \omega_{1+k}^\beta(\hat{V}^{-1}) \right) \mathbf{K} \right), \end{aligned} \quad (4.111)$$

using (A.13) and (A.30). This motivates the change of variable

$$\langle \chi | = \rho^{-1} \langle \tilde{\chi} | + 2 \left(\omega_1^\alpha(\rho^{L_0}) + \sum_{k=0}^{\infty} \tilde{\rho}^k R(v)^\alpha_\beta \omega_{1+k}^\beta(\hat{V}^{-1}) \right) \langle \mathcal{P}_\alpha |. \quad (4.112)$$

Note in particular that $\langle \tilde{\chi} |$ is a constrained field, since both $\langle \chi |$ and $\langle \mathcal{P}_\alpha |$ are. Using this formula one obtains by construction that

$$\rho \langle \mathcal{J}_\alpha^- | \otimes T^\alpha = 2 \langle \mathcal{P}_\alpha | \otimes \mathcal{V}^{-1} \mathcal{S}_{-1}(T^\alpha) \mathcal{V} + \langle \tilde{\chi} | \otimes \mathbf{K}. \quad (4.113)$$

With the above results, it becomes straightforward to re-express \mathcal{L}_2 , \mathcal{L}_3 and \mathcal{L}_4 in terms of the Maurer-Cartan form. We find

$$\begin{aligned} \mathcal{L}_2 &= 4 \rho^{-1} \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} T^\beta T^\alpha \mathcal{V}^{-1 \dagger} | \mathcal{P}_\beta \rangle, \\ \mathcal{L}_3 &= 4 \rho^{-1} \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_{-1}(T^\beta) \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1 \dagger} | \mathcal{P}_\beta \rangle + 4 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1 \dagger} | \tilde{\chi} \rangle + \langle \tilde{\chi} | \mathcal{V}^{-1} \mathcal{V}^{-1 \dagger} | \tilde{\chi} \rangle, \\ \mathcal{L}_4 &= 4 \rho^{-1} \langle P_0 | \mathcal{V}^{-1} T^\alpha \mathcal{V}^{-1 \dagger} | \mathcal{P}_\alpha \rangle. \end{aligned} \quad (4.114)$$

Instead of working out explicitly the expression of \mathcal{L}_1 step by step, we argue what the result should be based on the computation of the rigid $\hat{E}_8 \rtimes (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ invariance of \mathcal{L}_1 . To this purpose, let us first observe that the changes of variable (4.109), (4.112) and (4.113) essentially take the same form as the rigid transformations of the currents $\langle \mathcal{J}_\alpha |$, $\langle \mathcal{J}_\alpha^- |$ and the field $\langle \chi |$ under $\hat{E}_8 \rtimes (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$ that were presented in section 4.2.1, but now with \mathcal{V} playing the rôle of the group element associated to the transformation. An important point is that this comparison only holds if one disregards the constrained $\overline{R(\Lambda_0)_{-1}}$ vector index of the currents and $\langle \chi |$, whose associated transformation cancels that of \mathcal{M}^{-1} when considering the variation of \mathcal{L}_1 . The analogy drawn above and the rigid invariance of \mathcal{L}_1 imply that the explicit dependence on \mathcal{V} , apart from the \mathcal{M}^{-1} contracting the derivatives, is eliminated by the substitution (4.109) and (4.112) that induces $\mathcal{M} \rightarrow \mathcal{V}^{-1 \dagger} \mathcal{M} \mathcal{V}^{-1} = \mathbf{1}$. In the end, this change of variables simply amounts to cancel $\tilde{\rho}$ and $\Omega^\alpha(\mathcal{M})$ through

$\Omega^\alpha(\mathbf{1}) = 0$, such that the new expression of \mathcal{L}_1 in terms of $\langle \mathcal{P}_\alpha |$ and $\langle \tilde{\chi} |$ only depends implicitly on ρ , $\tilde{\rho}$ and \hat{V} through $\langle \mathcal{P}_\alpha |$ and the operator $\rho^{-\frac{1}{2}}\mathcal{V}^{-1}$

$$\mathcal{L}_1 = 4\rho^{-1} \sum_{n \in \mathbb{Z}} \langle P_A^{-n} | \mathcal{M}^{-1} | P_B^n \rangle \eta^{AB} - 8\rho^{-1} \langle P_0 | \mathcal{M}^{-1} | P_K \rangle - 8\rho^{-1} \langle P_1 | \mathcal{M}^{-1} | \tilde{\chi} - P_1 \rangle. \quad (4.115)$$

Combining the various terms one obtains the potential (4.1) in the following simple form

$$\begin{aligned} \rho V = & \eta^{\alpha\beta} \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle + \frac{1}{2} \langle \tilde{\chi} - 2P_1 | \mathcal{V}^{-1} \mathcal{V}^{-1\dagger} | \tilde{\chi} - 2P_1 \rangle - 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} T^\beta T^\alpha \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle \\ & + 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_{-1}(T^\beta) \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle + 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1\dagger} | \tilde{\chi} \rangle \\ & + 2 \langle P_0 | \mathcal{V}^{-1} T^\alpha \mathcal{V}^{-1\dagger} | \mathcal{P}_\alpha \rangle. \end{aligned} \quad (4.116)$$

Any solution to the section constraint only has a finite number of non-trivial components for the derivative $\langle \partial |$. This means that $\langle \partial | \mathcal{V}^{-1}$ only involves finite sums and is regular in the Borel gauge. Moreover, all potentially infinite sums of the dual potentials Y_n^A cancel in the potential. This cancellation can be associated with the invariance under δ_Σ generalised diffeomorphisms as these can be used to gauge away almost all Y_n^A . We shall see these facts more explicitly in the following section where we work out the potential in an E_8 decomposition. In summary, using the completed Kac-Moody group and a Borel gauge representative, the E_9 exceptional field theory potential (4.1) is completely well-defined.

5 Reduction to E_8 and consistency with supergravity

An inherent property of E_n exceptional field theories for $n \leq 8$, is that they reduce to eleven-dimensional supergravity or type IIB supergravity upon choosing the appropriate solution to the section condition. In this section, we shall partially demonstrate this property for the part of the E_9 exceptional field theory dynamics encoded in the E_9 scalar potential. Our strategy will consist in proving that the E_8 exceptional field theory with two external isometries is embedded in the scalar potential of E_9 exceptional field theory. In other words, we will show explicitly that when the infinite number of E_9 internal generalised coordinates are truncated to those of E_8 exceptional field theory, the potential (4.2) reproduces all the terms of the E_8 exceptional field theory Lagrangian for field configurations that do not depend on the two external coordinates. As a corollary, this implies that our potential encodes the dynamics of eleven-dimensional supergravity and type IIB supergravity with two external isometries.

5.1 E_8 section and exceptional field theory

In the present section, we are interested in relating the E_9 exceptional field theory to E_8 exceptional field theory [24] in $3 + 248 = 2 + (1 + 248)$ dimensions where the 248 directions are subject to an analogous E_8 section constraint. As indicated in the decomposition, we require $1 + 248$ directions to emerge from the coordinates $|Y\rangle$ in $R(\Lambda_0)_{-1}$ of E_9 . Similar

to (2.5) we have a decomposition of the coordinates according to

$$|Y\rangle = \left(\varphi + \eta_{AB} y^A T_{-1}^B + \sum_{n=2}^{\infty} y_{A_1 \dots A_n} T_{-1}^{A_1} \dots T_{-1}^{A_n} \right) |0\rangle. \quad (5.1)$$

It was shown in [13] that for any hyperplane solution to the E_9 section constraint (3.18) there exists an E_9 element that rotates this hyperplane to one lying completely along the directions φ and y^A , i.e., the lowest two pieces in the E_8 graded decomposition of $R(\Lambda_0)_{-1}$ corresponding to $1 + 248$ directions. Moreover, the remnant of the E_9 section constraint (3.18) implies that fields depend on the 248 directions y^A in such a way that they satisfy the E_8 section constraint [24, eq. (1.1)]. In practice, this solution to the section constraint is implemented by restricting $\langle \partial |$ to be of the form

$$\langle \partial | = \langle 0 | (\partial_\varphi + T_1^A \partial_A), \quad (5.2)$$

where $\partial_A = \frac{\partial}{\partial y^A}$ satisfies the E_8 section constraint.

The direction φ has an interpretation as the third external coordinate and y^A as the internal coordinates in the E_8 exceptional field theory. In this section, we will show that the potential (4.1) introduced in this paper is indeed consistent with the action of E_8 exceptional field theory in that it reproduces all its terms with no external derivatives with respect to the two directions $x^\mu = (t, x)$. The Lagrangian of E_8 exceptional field theory is of the schematic form

$$\mathcal{L}_{E_8} = \sqrt{-g} \hat{R} + \frac{1}{240} \sqrt{-g} g^{mn} \eta^{AB} J_{mA} J_{nB} - \sqrt{-g} V(M_0, g) + \mathcal{L}_{CS}, \quad (5.3)$$

where \hat{R} denotes the (improved) Ricci scalar of the 3 external directions with metric g_{mn} , where $m = 0, 1, 2$. M_0 are the scalar fields parameterising E_8 (see section 3.1) and J_{mA} are the components of the E_8 covariant current $J_m = M_0^{-1} D_m M_0$ where the covariant derivative D_m featuring in J_m and \hat{R} is covariantised with respect to the two gauge fields A_m^A and B_{mA} of the theory, where $A = 1, \dots, 248$ labels the adjoint of E_8 , and the gauge fields also appear in the Chern-Simons term \mathcal{L}_{CS} . The vector field B_{mA} is constrained in its E_8 index. The potential term $V(M_0, g)$ can be expressed through the E_8 internal current $M_0^{-1} \partial_A M_0$ and is a combination of terms similar to (4.2). The exact form of the various terms of (5.3) is given in [24].

As we shall show in detail in section 5.3, all terms except for the topological Chern-Simons term give contributions when restricting to derivatives along φ and the 248 internal coordinates y^A according to (5.2). First we parameterise the E_8 fields in a way that facilitates the comparison. For the metric g_{mn} on the three-dimensional external space we shall consider the (static) ansatz

$$ds^2 = e^{2\sigma} (-dt^2 + dx^2) + \rho^2 d\varphi^2. \quad (5.4)$$

Compared to (3.1) there is no Kaluza-Klein vector $A_\mu^{(3)}$ since we disregard all external form fields in this paper. In other words, the only components of the two vector fields that will appear are those along φ . For simplicity we shall write the remaining components of the vector fields $A^A = A_\varphi^A$ and $B_A = B_{\varphi A}$, without writing explicitly their φ index.

5.2 Parameterising \mathcal{M} and decomposition of the potential

We shall now decompose the potential (4.1) of E_9 exceptional field theory in the E_8 solution (5.2) to the section constraint. Moreover, we shall see explicitly that even though $R(\Lambda_0)_0$ is an infinite-dimensional representation, the total potential only gives rise to finitely many terms as we explained in section 4.3. In particular, we use the formulation in terms of the coset representative \mathcal{V} rather than \mathcal{M} .

First, as explained earlier, we can work at $\tilde{\rho} = 0$ without loss of generality by gauge-fixing partially the invariance of the potential under Σ transformations and this simplifies the analysis in this section as we only have to analyse (4.2). We now demonstrate this gauge-fixing explicitly using (5.2) as a solution to the section condition. On (5.2), the constrained parameter $\Sigma \sim |\Sigma\rangle\langle\pi_\Sigma|$ can be parameterised as

$$\begin{aligned} \Sigma = & \left(\sigma_0 - \eta^{AB} \Sigma_{A,B} + \sum_{n=1}^{\infty} \sigma_{A_1 \dots A_n} T_{-1}^{A_1} \dots T_{-1}^{A_n} \right) |0\rangle\langle 0| \\ & + \left(\Sigma_A + \Sigma_{A,B} T_{-1}^B + \sum_{n=2}^{\infty} \Sigma_{A,A_1 \dots A_n} T_{-1}^{A_1} \dots T_{-1}^{A_n} \right) |0\rangle\langle 0| T_1^A, \end{aligned} \quad (5.5)$$

where the coefficients have to be projected to the irreducible representations appearing at level n in (2.6). Moreover, it is clear from the structure of the generalised Lie derivative (3.15) that many of the components of Σ have a trivial action on \mathcal{M} .

The trace of Σ is a finite expression and given by

$$\text{Tr } \Sigma = \sigma_0. \quad (5.6)$$

Considering for simplicity a Σ gauge transformation with parameter $\Sigma = \sigma_0 |0\rangle\langle 0|$, one obtains according to (4.83a), (4.83c) and (4.92) the variations

$$\begin{aligned} \delta_{\sigma_0} \tilde{\rho} &= \sigma_0, \\ \delta_{\sigma_0} \mathcal{M} &= -\sigma_0 (\mathcal{M} L_{-1} + L_1 \mathcal{M}), \\ \delta_{\sigma_0} \langle \chi | &= -\frac{1}{\rho^2} \left(1 + \langle 0 | \mathcal{M}^{-1} (L_0 - \tilde{\rho} L_1) \mathcal{M} | 0 \rangle \right) \langle 0 | (\partial_\varphi + T_1^A \partial_A) \sigma_0. \end{aligned} \quad (5.7)$$

The first transformation shows that we can shift the field $\tilde{\rho}$ by a gauge parameter when exponentiated. Therefore, we can use a finite gauge transformation to set $\tilde{\rho} = 0$. As is evident from the other two equations, this will have a non-trivial effect on the dual potentials and $\langle \chi |$. Moreover, setting $\tilde{\rho} = 0$ can be done while preserving a residual gauge invariance under traceless Σ transformations satisfying $\sigma_0 = 0$. By a similar reasoning one can consider a more general class of Σ parameters and find that all the higher level potentials Y_n^A for $n \geq 2$ are also pure gauge, as follows from η_{-1} in (4.83a), see also [13, eq. (4.36)]. By contrast, Y_1^A cannot be completely gauged away but transforms as it should in $D = 3$ under gauge transformations [24].

For the rest of this section, we shall then work with the potential (4.2) at $\tilde{\rho} = 0$. For vanishing axion $\tilde{\rho}$, the matrix \mathcal{M}_{MN} belongs to E_9 and can be parameterised by a suitable $E_9/K(E_9)$ coset space representative in the Borel gauge. The latter follows from (3.2), and reads

$$\mathcal{V} = \rho^{-L_0} e^{-\sigma} V_0 \prod_{n=1}^{\infty} \exp(Y_n^A \eta_{AB} T_{-n}^B). \quad (5.8)$$

The potential at $\tilde{\rho} = 0$ can be expressed in terms of the \mathfrak{e}_9 -valued Maurer-Cartan form associated to the coset representative (5.8). This is most easily obtained by taking the expression (4.116) and setting $\tilde{\rho} = 0$. This gives

$$\begin{aligned} \rho V(M, \tilde{\chi}) &= \eta^{\alpha\beta} \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle - 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} T^\beta T^\alpha \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle \\ &\quad + 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_{-1}(T^\beta) \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1\dagger} | \mathcal{P}_\beta \rangle + 2 \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} \mathcal{S}_1(T^\alpha) \mathcal{V}^{-1\dagger} | \tilde{\chi} \rangle \\ &\quad + \frac{1}{2} \langle \tilde{\chi} | \mathcal{V}^{-1} \mathcal{V}^{-1\dagger} | \tilde{\chi} \rangle + 2 \langle \mathcal{P}_0 | \mathcal{V}^{-1} T^\alpha \mathcal{V}^{-1\dagger} | \mathcal{P}_\alpha \rangle, \end{aligned} \quad (5.9)$$

where $\langle \mathcal{P}_\alpha |$ and $\langle \tilde{\chi} |$ are given by (4.107) and (4.112) for $\tilde{\rho} = 0$, respectively. The (negative) Borel gauge representative (5.8) was chosen such that $\langle \mathcal{P}_\alpha | \mathcal{V}^{-1}$ gives rise to a finite expansion. To see this explicitly, we first note that the solution (5.2) of the section constraint implies for \mathcal{P}_M that one has the decomposition $\langle \mathcal{P}_\alpha | = \langle 0 | (\mathcal{P}_{\varphi,\alpha} + \mathcal{P}_{A,\alpha} T_1^A)$. Multiplying then by \mathcal{V}^{-1} from the right and using (5.8) one obtains the finite expression

$$\begin{aligned} \langle \mathcal{P}_\alpha | \mathcal{V}^{-1} &= \langle 0 | (\mathcal{P}_{\varphi,\alpha} + \mathcal{P}_{A,\alpha} T_1^A) \prod_{n=1}^{\infty} \exp(-Y_{nB} T_{-n}^B) \rho^{L_0} e^\sigma V_0^{-1} \\ &= \langle 0 | (\mathcal{P}_{\varphi,\alpha} - \eta^{AB} Y_{1A} \mathcal{P}_{B,\alpha} + \mathcal{P}_{A,\alpha} T_1^A) \rho^{L_0} e^\sigma V_0^{-1} \\ &= e^\sigma \langle 0 | (\mathcal{P}_{\varphi,\alpha} - \eta^{AB} Y_{1A} \mathcal{P}_{B,\alpha} + \rho \mathcal{P}_{A,\alpha} V_0 T_1^A V_0^{-1}). \end{aligned} \quad (5.10)$$

Similarly, the scalar field $\langle \tilde{\chi} |$ from (4.112) satisfies the section constraint and can thus be parameterised as

$$\langle \tilde{\chi} | = \langle 0 | (\tilde{\chi}_\varphi + \tilde{\chi}_A T_1^A). \quad (5.11)$$

As another preparatory step we need to introduce indices for the local $K(E_8) = \text{Spin}(16)/\mathbb{Z}_2$ subgroup appearing in the coset space $E_8/K(E_8)$ represented by V_0 . We do this by writing \underline{A} for the adjoint of E_8 transforming under the local $K(E_8)$ subgroup and define

$$\begin{aligned} \hat{\mathcal{P}}_{\varphi,\alpha} &\equiv \mathcal{P}_{\varphi,\alpha} - \eta^{AB} Y_{1A} \mathcal{P}_{B,\alpha}, & \hat{\mathcal{P}}_{\underline{A},\alpha} T_n^{\underline{A}} &\equiv \mathcal{P}_{A,\alpha} V_0 T_n^A V_0^{-1}, \\ \hat{\chi}_\varphi &\equiv \tilde{\chi}_\varphi - \eta^{AB} Y_{1A} \tilde{\chi}_B, & \hat{\chi}_{\underline{A}} T_n^{\underline{A}} &\equiv \tilde{\chi}_A V_0 T_n^A V_0^{-1}. \end{aligned} \quad (5.12)$$

More generally, we shall consider the notation that an underlined index is related to a

normal one through $X_{\underline{A}}T^{\underline{A}} = X_A V_0 T^A V_0^{-1}$ and in particular

$$\begin{aligned} \mathcal{P}_{A,\alpha}\mathcal{P}_{B,\beta}\langle 0|V_0T_n^AV_0^{-1}(V_0T_n^BV_0^{-1})^\dagger|0\rangle &= nM_0^{AB}\mathcal{P}_{A,\alpha}\mathcal{P}_{B,\beta} \\ &= \hat{\mathcal{P}}_{\underline{A},\alpha}\hat{\mathcal{P}}_{\underline{B},\beta}\langle 0|T_n^{\underline{A}}(T_n^{\underline{B}})^\dagger|0\rangle = n\delta^{\underline{A}\underline{B}}\hat{\mathcal{P}}_{\underline{A},\alpha}\hat{\mathcal{P}}_{\underline{B},\beta}. \end{aligned} \quad (5.13)$$

Since $\mathcal{P}_{M,\alpha}T^\alpha$ belongs to the coset component it satisfies the Hermiticity property $\mathcal{P}_{M,\alpha}T^\alpha = (\mathcal{P}_{M,\alpha}T^\alpha)^\dagger$ and thus the components (5.12) can be decomposed as

$$\begin{aligned} \hat{\mathcal{P}}_{\varphi,\alpha}T^\alpha &= P_{\varphi,0}L_0 + P_{\varphi,\mathbb{K}}\mathbb{K} + P_{\varphi,\underline{A}}T^{\underline{A}} + \frac{1}{2}\sum_{n=1}^{\infty}P_{\varphi,\underline{A}}^n(T_{-n}^{\underline{A}} + (T_{-n}^{\underline{A}})^\dagger), \\ \hat{\mathcal{P}}_{\underline{A},\alpha}T^\alpha &= P_{\underline{A},0}L_0 + P_{\underline{A},\mathbb{K}}\mathbb{K} + P_{\underline{A},\underline{B}}T^{\underline{B}} + \frac{1}{2}\sum_{n=1}^{\infty}P_{\underline{A},\underline{B}}^n(T_{-n}^{\underline{B}} + (T_{-n}^{\underline{B}})^\dagger). \end{aligned} \quad (5.14)$$

Since $(P_{\varphi,\underline{A}}T^{\underline{A}})^\dagger = P_{\varphi,\underline{A}}T^{\underline{A}}$ we did not explicitly symmetrise this \mathfrak{e}_8 -valued component of $\hat{\mathcal{P}}$.

Using this notation, one obtains for the \mathcal{L}_1 part of (5.9)

$$\eta^{\alpha\beta}\langle \mathcal{P}_\alpha|\mathcal{V}^{-1}\mathcal{V}^{-1\dagger}|\mathcal{P}_\beta\rangle = e^{2\sigma}\eta^{\alpha\beta}\left(\hat{\mathcal{P}}_{\varphi,\alpha}\hat{\mathcal{P}}_{\varphi,\beta} + \rho^2M_0^{AB}\mathcal{P}_{A,\alpha}\mathcal{P}_{B,\beta}\right), \quad (5.15)$$

for the \mathcal{L}_2 part

$$\begin{aligned} \langle \mathcal{P}_\alpha|\mathcal{V}^{-1}T^\beta T^\alpha \mathcal{V}^{-1\dagger}|\mathcal{P}_\beta\rangle &= e^{2\sigma}\langle 0|\left(\hat{\mathcal{P}}_{\varphi,\beta}T^\beta\hat{\mathcal{P}}_{\varphi,\alpha}T^\alpha + \rho^2T_1^{\underline{C}}\hat{\mathcal{P}}_{\underline{D},\beta}T^\beta\hat{\mathcal{P}}_{\underline{C},\alpha}T^\alpha(T_1^{\underline{D}})^\dagger\right)|0\rangle \\ &\quad + 2\rho e^{2\sigma}\langle 0|T_1^{\underline{C}}\hat{\mathcal{P}}_{\varphi,\beta}T^\beta\hat{\mathcal{P}}_{\underline{C},\alpha}T^\alpha|0\rangle, \end{aligned} \quad (5.16)$$

and similarly for \mathcal{L}_3

$$\begin{aligned} &2\langle \mathcal{P}_\alpha|\mathcal{V}^{-1}\mathcal{S}_{-1}(T^\beta)\mathcal{S}_1(T^\alpha)\mathcal{V}^{-1\dagger}|\mathcal{P}_\beta\rangle + 2\langle \mathcal{P}_\alpha|\mathcal{V}^{-1}\mathcal{S}_1(T^\alpha)\mathcal{V}^{-1\dagger}|\tilde{\chi}\rangle + \frac{1}{2}\langle \tilde{\chi}|\mathcal{V}^{-1}\mathcal{V}^{-1\dagger}|\tilde{\chi}\rangle \\ &= \frac{1}{2}e^{2\sigma}\langle 0|(2\hat{\mathcal{P}}_{\varphi,\beta}\mathcal{S}_{-1}(T^\beta) + \hat{\chi}_\varphi)(2\hat{\mathcal{P}}_{\varphi,\alpha}\mathcal{S}_{-1}(T^\alpha)^\dagger + \hat{\chi}_\varphi)|0\rangle \\ &\quad + \frac{1}{2}e^{2\sigma}\rho^2\langle 0|T_1^{\underline{C}}(\hat{\mathcal{P}}_{\underline{D},\beta}\mathcal{S}_{-1}(2T^\beta) + \hat{\chi}_{\underline{D}})(2\hat{\mathcal{P}}_{\underline{C},\alpha}\mathcal{S}_{-1}(T^\alpha)^\dagger + \hat{\chi}_{\underline{C}})(T_1^{\underline{D}})^\dagger|0\rangle \\ &\quad + e^{2\sigma}\rho\langle 0|T_1^{\underline{C}}(2\hat{\mathcal{P}}_{\varphi,\beta}\mathcal{S}_{-1}(T^\beta) + \hat{\chi}_\varphi)(2\hat{\mathcal{P}}_{\underline{C},\alpha}\mathcal{S}_{-1}(T^\alpha)^\dagger + \hat{\chi}_{\underline{C}})|0\rangle. \end{aligned} \quad (5.17)$$

We now start to collect the different pieces in the potential term $\frac{1}{4}\mathcal{L}_1 - \frac{1}{2}\mathcal{L}_2 + \frac{1}{2}\mathcal{L}_3$ in order to match them with the corresponding terms in the E_8 exceptional field theory

action. We begin with the terms bilinear in $\mathcal{P}_{\varphi,\alpha}$. These are, after removing the overall $e^{2\sigma}$ factor,

$$\begin{aligned}
 & \eta^{\alpha\beta}\hat{\mathcal{P}}_{\varphi,\alpha}\hat{\mathcal{P}}_{\varphi,\beta}-2\langle 0|\hat{\mathcal{P}}_{\varphi,\beta}T^\beta\hat{\mathcal{P}}_{\varphi,\alpha}T^\alpha|0\rangle+\frac{1}{2}\langle 0|(2\hat{\mathcal{P}}_{\varphi,\beta}\mathcal{S}_{-1}(T^\beta)+\hat{\chi}_\varphi)(2\hat{\mathcal{P}}_{\varphi,\alpha}\mathcal{S}_{-1}(T^\alpha)^\dagger+\hat{\chi}_\varphi)|0\rangle \\
 & =\delta^{AB}\hat{\mathcal{P}}_{\varphi,A}\hat{\mathcal{P}}_{\varphi,B}-2\hat{\mathcal{P}}_{\varphi,0}\hat{\mathcal{P}}_{\varphi,K}+\frac{1}{2}\sum_{n=1}^{\infty}\delta^{AB}\hat{\mathcal{P}}_{\varphi,A}^n\hat{\mathcal{P}}_{\varphi,B}^n \\
 & \quad -2\hat{\mathcal{P}}_{\varphi,K}\hat{\mathcal{P}}_{\varphi,K}-\frac{1}{2}\sum_{n=1}^{\infty}n\delta^{AB}\hat{\mathcal{P}}_{\varphi,A}^n\hat{\mathcal{P}}_{\varphi,B}^n+\frac{1}{2}\hat{\chi}_\varphi\hat{\chi}_\varphi+\frac{1}{2}\sum_{n=1}^{\infty}n\delta^{AB}\hat{\mathcal{P}}_{\varphi,A}^{n+1}\hat{\mathcal{P}}_{\varphi,B}^{n+1} \\
 & =\delta^{AB}\hat{\mathcal{P}}_{\varphi,A}\hat{\mathcal{P}}_{\varphi,B}-2(\hat{\mathcal{P}}_{\varphi,0}+\hat{\mathcal{P}}_{\varphi,K})\hat{\mathcal{P}}_{\varphi,K}+\frac{1}{2}\hat{\chi}_\varphi\hat{\chi}_\varphi. \tag{5.18}
 \end{aligned}$$

Note that all the higher potential field strengths $\hat{\mathcal{P}}_{\varphi,A}^n$ cancel.

Next we consider all terms bilinear in $\mathcal{P}_{\underline{C},\alpha}$. The \mathcal{L}_1 terms are simply $2\eta^{\alpha\beta}M_0^{CD}\mathcal{P}_{\underline{C},\alpha}\mathcal{P}_{\underline{D},\beta}$, while the ones from \mathcal{L}_2 are

$$\begin{aligned}
 & 4\langle 0|T_1^C\mathcal{P}_{\underline{D},\beta}T^\beta\mathcal{P}_{\underline{C},\alpha}T^\alpha(T_1^D)^\dagger|0\rangle \\
 & =\langle 0|\left[2(P_{\underline{D},0}+P_{\underline{D},K})T_1^C+2P_{\underline{D},B}f_{\underline{E}}^{CB}T_1^E+P_{\underline{D},B}\eta^{CB}+\sum_{n=1}^{\infty}P_{\underline{D},B}^nT_1^C(T_{-n}^B)^\dagger\right] \\
 & \quad \left[2(P_{\underline{C},0}+P_{\underline{C},K})(T_1^D)^\dagger+2P_{\underline{C},A}f_{\underline{E}}^{DA}T_1^E+P_{\underline{C},A}\eta^{DA}+\sum_{n=1}^{\infty}P_{\underline{C},A}^nT_{-n}^A(T_1^D)^\dagger\right]|0\rangle \\
 & =4\delta^{CD}(P_{\underline{D},0}+P_{\underline{D},K})(P_{\underline{C},0}+P_{\underline{C},K})+4P_{\underline{D},B}P_{\underline{C},A}\delta^{EF}f_{\underline{E}}^{CB}f_{\underline{E}}^{DA} \\
 & \quad +8(P_{\underline{D},0}+P_{\underline{D},K})P_{\underline{C},A}\delta^{CE}f_{\underline{E}}^{DA}+(\eta^{CB}\eta^{DA}+\eta^{BD}\eta^{AC})P_{\underline{D},B}P_{\underline{C},A} \\
 & \quad +\sum_{n=1}^{\infty}(n\delta^{AB}\delta^{CD}+\delta^{BE}\delta^{DF}f_{\underline{G}\underline{E}}^A f_{\underline{G}\underline{F}}^{CG})P_{\underline{D},B}^n P_{\underline{C},A}^n. \tag{5.19}
 \end{aligned}$$

The terms quadratic in $\mathcal{P}_{\underline{C},\alpha}$ coming from \mathcal{L}_3 are

$$\begin{aligned}
 & \langle 0|T_1^C(2\hat{\mathcal{P}}_{\underline{D},\beta}\mathcal{S}_{-1}(T^\beta)+\hat{\chi}_{\underline{D}})(2\hat{\mathcal{P}}_{\underline{C},\alpha}\mathcal{S}_{-1}(T^\alpha)^\dagger+\hat{\chi}_{\underline{C}})(T_1^D)^\dagger|0\rangle \\
 & =\langle 0|\left[\hat{\chi}_{\underline{D}}T_1^C+P_{\underline{D},B}\delta^{BF}f_{\underline{E}}^C T_1^E+2P_{\underline{D},B}\eta^{CB}+\sum_{n=1}^{\infty}P_{\underline{D},B}^{n+1}T_1^C(T_{-n}^B)^\dagger\right] \\
 & \quad \left[\hat{\chi}_{\underline{C}}(T_1^D)^\dagger+P_{\underline{C},A}\delta^{AG}f_{\underline{G}\underline{H}}^D(T_1^H)^\dagger+2P_{\underline{C},A}\eta^{DA}+\sum_{n=1}^{\infty}P_{\underline{C},A}^{n+1}T_{-n}^A(T_1^D)^\dagger\right]|0\rangle \\
 & =\delta^{CD}\hat{\chi}_{\underline{C}}\hat{\chi}_{\underline{D}}+4\eta^{CB}\eta^{DA}P_{\underline{D},B}P_{\underline{C},A}+2f_{\underline{B}}^{CA}\delta^{BD}\hat{\chi}_{\underline{D}}P_{\underline{C},A}+\delta^{EF}f_{\underline{E}}^{DB}f_{\underline{E}}^{CA}P_{\underline{D},B}P_{\underline{C},A} \\
 & \quad +\eta^{BD}\eta^{AC}P_{\underline{D},B}P_{\underline{C},A}+\sum_{n=1}^{\infty}((n-1)\delta^{AB}\delta^{CD}+\delta^{BE}\delta^{DF}f_{\underline{G}\underline{E}}^A f_{\underline{G}\underline{F}}^{CG})P_{\underline{D},B}^n P_{\underline{C},A}^n. \tag{5.20}
 \end{aligned}$$

In rewriting the final expression we have used,⁸

$$\delta^{BE}\delta^{DF}f_{\underline{G}\underline{E}}^C f_{\underline{G}\underline{F}}^{AG}=\delta^{BE}\delta^{DF}f_{\underline{G}\underline{E}}^A f_{\underline{G}\underline{F}}^{CG}+\delta^{EF}f_{\underline{E}}^{DB}f_{\underline{E}}^{CA}. \tag{5.21}$$

⁸The structure constants $f_{\underline{A}\underline{B}}^C$ are given by $f_{\underline{A}\underline{B}}^C=\eta_{\underline{A}\underline{D}}\eta_{\underline{B}\underline{E}}\eta^{CE}f^{DE}_{\underline{F}}=-\delta_{\underline{A}\underline{D}}\delta_{\underline{B}\underline{E}}\delta^{CE}f^{DE}_{\underline{F}}$ and one uses the Jacobi identity to derive this identity.

Combining the terms bilinear in $\mathcal{P}_{\underline{C},\alpha}$ in $\frac{1}{4}\mathcal{L}_1 - \frac{1}{2}\mathcal{L}_2 + \frac{1}{2}\mathcal{L}_3$ determined above then gives the following somewhat lengthy expression

$$\begin{aligned}
 & \eta^{\alpha\beta} M_0^{CD} \mathcal{P}_{C,\alpha} \mathcal{P}_{D,\beta} - 2 \langle 0 | T_1^{\underline{C}} \hat{\mathcal{P}}_{\underline{D},\beta} T^\beta \hat{\mathcal{P}}_{\underline{C},\alpha} T^\alpha (T_1^{\underline{D}})^\dagger | 0 \rangle \\
 & \quad + \frac{1}{2} \langle 0 | T_1^{\underline{C}} (2\hat{\mathcal{P}}_{\underline{D},\beta} \mathcal{S}_{-1}(T^\beta) + \hat{\chi}_{\underline{D}}) (2\hat{\mathcal{P}}_{\underline{C},\alpha} \mathcal{S}_{-1}(T^\alpha)^\dagger + \hat{\chi}_{\underline{C}}) (T_1^{\underline{D}})^\dagger | 0 \rangle \\
 = & M_0^{CD} \delta^{AB} P_{C,\underline{A}} P_{D,\underline{B}} - 2M_0^{CD} P_{C,0} P_{D,\kappa} + \frac{1}{2} \delta^{CD} \sum_{n=1}^{\infty} \delta^{AB} P_{C,\underline{A}}^n P_{D,\underline{B}}^n \\
 & - 2\delta^{CD} (P_{\underline{D},0} + P_{\underline{D},\kappa}) (P_{\underline{C},0} + P_{\underline{C},\kappa}) - 2P_{\underline{D},\underline{B}} P_{C,\underline{A}} \delta^{EF} f^{CB}{}_E f^{DA}{}_F \\
 & - 4(P_{\underline{D},0} + P_{\underline{D},\kappa}) P_{C,\underline{A}} \delta^{CE} f^{DA}{}_E - \frac{1}{2} (\eta^{CB} \eta^{DA} + \eta^{BD} \eta^{AC}) P_{\underline{D},\underline{B}}^1 P_{C,\underline{A}}^1 \\
 & \quad - \frac{1}{2} \sum_{n=1}^{\infty} (n \delta^{AB} \delta^{CD} + \delta^{BE} \delta^{DF} f_{GE}{}^A f^{CG}{}_F) P_{\underline{D},\underline{B}}^n P_{C,\underline{A}}^n \\
 & + \frac{1}{2} \delta^{CD} \hat{\chi}_{\underline{C}} \hat{\chi}_{\underline{D}} + 2\eta^{CB} \eta^{DA} P_{\underline{D},\underline{B}} P_{C,\underline{A}} + f^{CA}{}_B \delta^{BD} \hat{\chi}_{\underline{D}} P_{C,\underline{A}}^1 + \frac{1}{2} \delta^{EF} f^{DB}{}_E f^{CA}{}_F P_{\underline{D},\underline{B}}^1 P_{C,\underline{A}}^1 \\
 & + \frac{1}{2} \eta^{BD} \eta^{AC} P_{\underline{D},\underline{B}}^2 P_{C,\underline{A}}^2 + \frac{1}{2} \sum_{n=1}^{\infty} ((n-1) \delta^{AB} \delta^{CD} + \delta^{BE} \delta^{DF} f_{GE}{}^A f^{CG}{}_F) P_{\underline{D},\underline{B}}^n P_{C,\underline{A}}^n \\
 = & M_0^{CD} \eta^{AB} P_{C,\underline{A}} P_{D,\underline{B}} - 2M_0^{EF} f^{CB}{}_E f^{DA}{}_F P_{D,\underline{B}} P_{C,\underline{A}} + 2\eta^{CB} \eta^{DA} P_{D,\underline{B}} P_{C,\underline{A}} \\
 & - \frac{1}{2} M_0^{CD} (4P_{C,0} P_{D,0} + 12P_{C,0} P_{D,\kappa} + 4P_{C,\kappa} P_{D,\kappa} - \hat{\chi}_C \hat{\chi}_D) \\
 & - \frac{1}{2} (\eta^{CB} \eta^{DA} + \eta^{BD} \eta^{AC}) P_{D,\underline{B}}^1 P_{C,\underline{A}}^1 + \frac{1}{2} M_0^{EF} f^{DB}{}_E f^{CA}{}_F P_{D,\underline{B}}^1 P_{C,\underline{A}}^1 \\
 & - 4M_0^{AB} f^{CD}{}_A P_{C,D} (P_{B,0} + P_{B,\kappa}) + f^{AB}{}_D M_0^{DC} P_{A,B}^1 \hat{\chi}_C + \frac{1}{2} \eta^{BD} \eta^{AC} P_{D,\underline{B}}^2 P_{C,\underline{A}}^2. \quad (5.22)
 \end{aligned}$$

Because the structure constant $f^{AB}{}_C$ and the Killing form η^{AB} are E_8 invariant, trading local $K(E_8)$ indices $\underline{A}, \underline{B}, \dots$ for E_8 indices A, B, \dots by conjugation with V_0 amounts in practice to simply replacing δ^{AB} by M_0^{AB} according to (5.13). Once again all the higher level scalar field strengths $P_{A,\underline{B}}^n$ cancel out for $n > 2$ for the final expression involving all bilinears in $\mathcal{P}_{\underline{C},\alpha}$.

Finally, we collect all the terms in $\mathcal{P}_{\varphi,\beta} \mathcal{P}_{\underline{C},\alpha}$. These have no contribution from \mathcal{L}_1 and the terms arising in $-\frac{1}{2}\mathcal{L}_2 + \frac{1}{2}\mathcal{L}_3$ are

$$\begin{aligned}
 & -2 \langle 0 | T_1^{\underline{C}} \hat{\mathcal{P}}_{\varphi,\beta} T^\beta \hat{\mathcal{P}}_{\underline{C},\alpha} T^\alpha | 0 \rangle + \frac{1}{2} \langle 0 | T_1^{\underline{C}} (2\hat{\mathcal{P}}_{\varphi,\beta} \mathcal{S}_{-1}(T^\beta) + \hat{\chi}_\varphi) (2\hat{\mathcal{P}}_{\underline{C},\alpha} \mathcal{S}_{-1}(T^\alpha)^\dagger + \hat{\chi}_{\underline{C}}) | 0 \rangle \\
 = & -\eta^{AB} P_{\varphi,\underline{A}}^1 P_{\underline{B},\kappa} + 2f^{ABC} P_{\varphi,\underline{A}} P_{\underline{B},\underline{C}}^1 - (P_{\varphi,0} + P_{\varphi,\kappa}) \eta^{AB} P_{A,\underline{B}}^1 + \frac{1}{2} \sum_{n=1}^{\infty} \delta^{CD} f^{AB}{}_D P_{\varphi,\underline{C}}^n P_{A,\underline{B}}^{n+1} \\
 & + \eta^{AB} P_{\varphi,\underline{A}} \hat{\chi}_{\underline{B}} + \frac{1}{2} \hat{\chi}_\varphi \eta^{AB} P_{A,\underline{B}}^2 - \frac{1}{2} \sum_{n=1}^{\infty} \delta^{CD} f^{AB}{}_D P_{\varphi,\underline{C}}^n P_{A,\underline{B}}^{n+1} \\
 = & -\eta^{AB} P_{\varphi,\underline{A}}^1 P_{\underline{B},\kappa} + f^{ABC} P_{\varphi,\underline{A}} P_{\underline{B},\underline{C}}^1 - (P_{\varphi,0} + P_{\varphi,\kappa}) \eta^{AB} P_{A,\underline{B}}^1 + \eta^{AB} P_{\varphi,\underline{A}} \hat{\chi}_{\underline{B}} + \frac{1}{2} \hat{\chi}_\varphi \eta^{AB} P_{A,\underline{B}}^2. \quad (5.23)
 \end{aligned}$$

Finally we compute the expression of $\frac{1}{2}\mathcal{L}_4$ that gives

$$2\langle\mathcal{P}_0|\mathcal{V}^{-1}T^\alpha\mathcal{V}^{-1\dagger}|\mathcal{P}_\alpha\rangle = e^{2\sigma}\left(2P_{\varphi,0}P_{\varphi,\kappa} + \rho\eta^{AB}P_{A,0}P_{\varphi,B}^1 + \rho P_{\varphi,0}\eta^{AB}P_{A,B}^1 \quad (5.24)\right. \\ \left.+ 2\rho^2M_0^{AB}P_{A,0}(P_{B,0} + P_{B,\kappa}) + 2\rho^2M_0^{AB}f^{CD}{}_B P_{A,0}P_{C,D}\right).$$

Having collected and simplified all the terms appearing in the potential, we now need to explain how the various components relate to the quantities of E_8 exceptional field theory. First, we identify the dual potential Y_1^A with the three-dimensional vector field along the φ direction $Y_1^A = A^A$. This is natural as the vector fields in $D = 3$ are dual to the scalar fields and after reduction to two dimensions the relevant part of this duality equation becomes exactly (3.3). Similarly, the φ component of the E_8 constrained vector field reduces to a constrained scalar in two dimensions, that is we impose $\tilde{\chi}_A = \rho^{-1}B_A$. Evaluating the components of \mathcal{P}_φ one obtains

$$P_{\varphi,0} = -\rho^{-1}(\partial_\varphi\rho - A^A\partial_A\rho) \equiv -\rho^{-1}D\rho, \\ P_{\varphi,\kappa} = -(\partial_\varphi\sigma - A^A\partial_A\sigma) \equiv -D\sigma, \quad (5.25) \\ P_{\varphi,A}^1 = \rho^{-1}\eta_{AB}(\partial_\varphi A^B - A^C\partial_C A^B) \equiv \rho^{-1}\eta_{AB}DA^B,$$

where we have introduced the notation D for $\partial_\varphi - A^A\partial_A$. Note that D as introduced here is *not* the full covariant derivative D_φ that defines J_{mA} and \hat{R} in (5.3) [24], but only includes the transport term.

The components of \mathcal{P}_A become similarly

$$P_{A,0} = -\rho^{-1}\partial_A\rho, \quad P_{A,\kappa} = -\partial_A\sigma, \\ 2P_{A,B} = \eta_{BC}J_A{}^C, \quad P_{A,B}^1 = \rho^{-1}\eta_{BC}\partial_A A^C, \quad (5.26) \\ P_{A,B}^2 = \rho^{-2}\eta_{BC}(\partial_A Y_2^C + \frac{1}{2}f_{DE}{}^C A^D\partial_A A^E).$$

Here, $J_A{}^C$ denotes the internal component of the \mathfrak{e}_8 current defined from M_0 as

$$M_0^{-1CD}\partial_A M_{0BD} = J_A{}^D f_{DB}{}^C. \quad (5.27)$$

With these identifications we can now rewrite the full potential (5.9) using also the rearrangements (5.18), (5.20), (5.23) and (5.24) for the various bilinears. The result is the following long expression

$$e^{-2\sigma}V = -2\rho^{-1}(D\sigma)^2 + DA^A(\partial_A\rho^{-1} + 2\rho^{-1}\partial_A\sigma) + (-D\rho^{-1} + 2\rho^{-1}D\sigma)\partial_A A^A \\ + \rho^{-1}\eta^{AB}P_{\varphi,A}P_{\varphi,B} - 2\rho^{-1}f_{AB}{}^C\eta^{AD}P_{\varphi,D}\partial_C A^B + \frac{1}{2}\rho^{-3}(\rho\tilde{\chi}_\varphi + \partial_A Y_2^A + \frac{1}{2}f_{AB}{}^C A^A\partial_C A^B)^2 \\ - \frac{1}{2}\rho^{-1}(\partial_A A^B\partial_B A^A + \partial_A A^A\partial_B A^B) + \frac{1}{2}\rho^{-1}M_0^{AB}f_{AC}{}^D\partial_D A^C f_{BE}{}^F\partial_F A^E \\ + \frac{1}{4}\rho\left(M_0^{CD}\eta_{AB}J_C{}^A J_{D,B}{}^B - 2M_0^{EF}f_{BE}{}^C f_{AF}{}^D J_{D,B}{}^B J_C{}^A + 2J_{A,B}{}^B J_B{}^A\right) \\ - 2M_0^{AB}\partial_A\sigma(\rho\partial_B\sigma + 2\partial_B\rho) - M_0^{AB}f_{AD}{}^C J_C{}^D(\partial_B\rho + 2\rho\partial_B\sigma) \\ + 2\rho^{-1}\eta^{AB}P_{\varphi,B}P_{A,B} + \frac{1}{2}\rho^{-1}M_0^{AB}B_A B_B - \rho^{-1}M_0^{AB}f_{AC}{}^D\partial_D A^C B_B. \quad (5.28)$$

5.3 Comparison with E_8 exceptional theory

The above form of the potential still does not look very similar to the standard action of E_8 exceptional field theory sketched in (5.3), and in particular it still contains the dual potential Y_2^A and the constrained scalar $\hat{\chi}_\varphi$ that were not considered in [24]. In order to recognise the standard terms we now expand them out.

The kinetic term for the scalar fields reduces to an expression in terms of the covariant current along the φ direction

$$J^A = 2\eta^{AB}P_{\varphi,B} - (M_0^{AB} + \eta^{AB})(f_{BC}{}^D\partial_D A^C - B_B), \quad (5.29)$$

with $2P_{\varphi,A}T^A = M_0^{-1}(\partial_\varphi - A^A\partial_A)M_0 = M_0^{-1}DM_0$. This is the only non-trivial surviving part of the kinetic term and becomes explicitly

$$\begin{aligned} \frac{1}{2}\eta_{AB}J^AJ^B &= 2\eta^{AB}P_{\varphi,A}P_{\varphi,B} - 4f_{AB}{}^C\eta^{AD}P_{\varphi,D}\partial_C A^B + \partial_A A^B\partial_B A^A + (\partial_A A^A)^2 + M_0^{AB}B_AB_B \\ &\quad + 4B_A\eta^{AB}P_{\varphi,B} - 2M_0^{AB}f_{AC}{}^D\partial_D A^C B_B + M_0^{AB}f_{AC}{}^D\partial_D A^C f_{BE}{}^F\partial_F A^E, \end{aligned} \quad (5.30)$$

where we can already anticipate how several of the terms in (5.28) above simplify.

With the metric (5.4), one computes that

$$\begin{aligned} g^{-1}\partial_A g &= 2\rho^{-1}(2\rho\partial_A\sigma + \partial_A\rho), \\ g^{-1}\partial_A g g^{-1}\partial_B g + \partial_A g^{\mu\nu}\partial_B g_{\mu\nu} &= 8\rho^{-1}\partial_A(\sigma(\rho\partial_B)\sigma + 2\partial_B\rho), \end{aligned} \quad (5.31)$$

and dropping the dependence in t and x one obtains for the improved Ricci scalar that

$$e^{2\sigma}\rho\hat{R} - d(\varepsilon_{abc}e^a \wedge \omega^{bc}) + \partial_A(A^A e^a \wedge \omega^{bd}) = -2\rho^{-1}e^{2\sigma}(D\sigma - \partial_A A^A)^2. \quad (5.32)$$

The two total derivatives were introduced to write the Einstein-Hilbert Lagrangian in terms of the generalised anholonomies. Using integration by part one obtains from this

$$\begin{aligned} &e^{2\sigma}\left(2DA^A(\partial_A\rho^{-1} + 2\rho^{-1}\partial_A\sigma) + 2(-D\rho^{-1} + 2\rho^{-1}D\sigma)\partial_A A^A\right) \\ &= 2\rho^{-1}e^{2\sigma}\left(4D\sigma\partial_A A^A + \partial_A A^B\partial_B A^A - (\partial_A A^A)^2\right) \\ &\quad + 2\partial_A\left(\rho^{-1}e^{2\sigma}(DA^A + A^A\partial_B A^B)\right) - 2\partial_\varphi\left(\rho^{-1}e^{2\sigma}\partial_A A^A\right). \end{aligned} \quad (5.33)$$

Using (5.30) and (5.33) to rewrite (5.28) one obtains that

$$\begin{aligned} -e^{-2\sigma}V &= \rho\hat{R} - \frac{1}{4}\rho^{-1}\eta_{AB}J^AJ^B + \frac{1}{2}\partial_A\mathcal{B}^A + \frac{1}{2}\partial_\varphi\mathcal{B}^0 \\ &\quad - \rho\left(\frac{1}{4}M_0^{CD}\eta_{AB}J_{C,A}J_{D,B} - \frac{1}{2}M_0^{EF}f_{BE}{}^C f_{AF}{}^D J_{D,B}J_{C,A} + \frac{1}{2}J_{A,B}J_{B,A}\right) \\ &\quad + \frac{1}{2}M_0^{AB}f_{AD}{}^C J_{C,D}g^{-1}\partial_B g + \frac{1}{4}M_0^{AB}(g^{-1}\partial_A g g^{-1}\partial_B g + \partial_A g^{\mu\nu}\partial_B g_{\mu\nu}) \\ &\quad - \frac{1}{2}\rho^{-1}\left(\hat{\chi}_\varphi + \rho^{-1}\left(\partial_A Y_2^A + \frac{1}{2}f_{AB}{}^C A^A\partial_C A^B\right)\right)^2. \end{aligned} \quad (5.34)$$

Here, \mathcal{B}^A and \mathcal{B}^0 are boundary terms introduced by the partial integrations. The three first lines of (5.34) reproduce indeed the Lagrangian (5.3), when neglecting the dependence

in the two-dimensional external coordinates t and x , such that the topological term does not contribute. In particular, the second and third line correspond to the potential of E_8 exceptional field theory written in terms of currents and is structurally the same as (4.2). The last line exhibits that $\hat{\chi}_\varphi$ is an auxiliary field that can be integrated out without affecting the other fields, and after its elimination the Lagrangian does not depend on the higher level potential Y_2^A . The fields $\langle\chi\rangle$ is eventually fixed to

$$\langle\chi\rangle = \rho^{-2}\langle 0|\left(-\partial_A Y_2^A - \frac{1}{2}f_{AB}{}^C A^A \partial_C A^B + B_A A^A + B_A T_1^A\right) + 2\omega_1^\alpha(\mathcal{V}^{-1})\langle P_\alpha\rangle. \quad (5.35)$$

This removes all dependence on the unwanted fields so that we obtain a perfect match with all terms of E_8 exceptional field theory that can be reproduced from the E_9 potential constructed in this paper.

It was shown in [49, 50], and more generally in [51], that the only two inequivalent maximal hyperplane solutions to the section constraint of E_n exceptional field theories for $n \leq 8$, correspond to eleven-dimensional supergravity and type IIB supergravity. For E_8 exceptional field theory, a partial dictionary with eleven-dimensional supergravity was provided explicitly in [24, 52]. The above results then imply that, after appropriately solving the section constraint, our E_9 exceptional field theory potential reduces to the eleven-dimensional or type IIB supergravity Lagrangians for field configurations that do not depend on the two external coordinates t and x .

6 Conclusions

In this paper, we have constructed the potential of E_9 exceptional field theory as an invariant under E_9 generalised diffeomorphisms. This potential is constructed out of (at most) two internal derivatives acting on the scalar fields and is the first example of such an invariant potential for an infinite-dimensional duality group and an infinite-dimensional coordinate representation. The potential consists of four terms, separately invariant under rigid $\hat{E}_8 \times \mathbb{R}_{L-1}$ transformations and transforming homogeneously under \mathbb{R}_d^+ , where $E_9 = \hat{E}_8 \times \mathbb{R}_d^+$. Invariance under generalised diffeomorphisms (up to a total derivative) is only achieved by conspiring cancellations among the variations of the different terms. Another key new feature of E_9 exceptional field theory is the appearance of a covariantly constrained field χ_M already in the scalar sector. This constrained scalar field also enters crucially in the potential by forming an indecomposable representation together with the (non-central) components of the \mathfrak{e}_9 current.

Because of the complicated representation theory of E_9 and of its extension by \mathbb{R}_{L-1} , which admit indecomposable (but not irreducible) representations, it is not known whether there are only a finite number of terms invariant under the rigid symmetries of the theory that could in principle contribute to the potential. It is therefore difficult to state whether our result could be uniquely determined by requiring invariance under generalised diffeomorphisms. This is however not necessary for our purpose, as we also require that the dynamics of $D = 11$ and type IIB supergravity are reproduced upon solving the section constraint, and have proved that this is the case by mapping our expression to the potential of E_8 exceptional field theory. This is sufficient to guarantee uniqueness of our result.

The result of this paper is the first building block for the full E_9 exceptional field theory. Specifically, it represents the truncation of E_9 exceptional field theory to scalar fields and vanishing external derivatives. The full theory will combine the scalar fields introduced in this paper with gauge fields $\{A_\mu^M, B_\mu^M{}_N\}$, transforming in the representations of the gauge parameters of the generalised diffeomorphisms (3.15). These gauge fields will covariantise external derivatives but also couple separately via a topological (Wess-Zumino-like) term. As customary in all even dimensions, the full theory will presumably admit its most compact formulation in terms of a pseudo-action supplemented by certain first-order duality equations, in this case for the scalar fields. The latter would define the extension of the linear system underlying two-dimensional maximal supergravity to the full exceptional field theory — after solving the section constraints thus to full $D = 11$ and type IIB supergravity. In particular, these equations should provide first order equations for the constrained scalar field χ_M , confirming that this is not an additional propagating degree of freedom, but rather is determined by the physical fields of the theory. The precise match with two-dimensional supergravity will require the identification of the dictionary among the components of our matrix \mathcal{M}^{MN} and the infinite tower of dual scalar potentials encoded in the various formulations of the linear system [17, 39, 53–55].

As already discussed in the introduction, an immediate application of E_9 exceptional field theory will be its reduction by means of a generalised Scherk-Schwarz ansatz [13], which together with the dictionary to supergravity fields would exhibit the structure of the yet elusive scalar potential of gauged maximal $D = 2$ supergravity [31] without the need to resort to the fermionic sector and supersymmetry of the theory. A notable aspect of the gauged maximal $D = 2$ supergravities studied in [31] is the ubiquity of the gauging of the L_{-1} generator that also featured in our construction and the generalised Lie derivative. The constrained scalar field χ_M is also indispensable in the generalised Scherk-Schwarz ansatz for such gaugings.

It would also be very desirable to reformulate our potential in terms of the manifestly covariant components of a suitably defined internal Ricci tensor, analogous to the structures identified for the lower-rank groups [6, 56, 57]. However, such a formulation would first require the identification of (the unambiguous components of) an appropriate internal $K(E_9)$ spin connection, which at the moment seems a formidable task given the non-reductiveness of $K(E_9)$ and the fact that it does not admit highest weight representations.

Our work can also be considered as a step towards understanding the E_{11} conjecture [58–60] as well as the E_{10} conjecture [61]. The advantage of the group E_9 considered here is that it admits an explicit realisation as a vertex operator algebra which allows to define explicitly the full non-linear theory. One can nevertheless expect that there exists an E_{11} exceptional field theory that would include all the others by considering specific partial solutions of its section constraint. The latter does not appear explicitly in the formulation of [60] but played a crucial role in a different linearised system extending E_{11} [62].

While we have focussed on the construction of the potential of E_9 exceptional field theory, our expressions and proof of invariance are equally valid for any affine Kac-Moody group based on a finite-dimensional simple Lie group G , in which case the rigid symmetry group of the potential is $\hat{G} \rtimes (\mathbb{R}_d^+ \times \mathbb{R}_{L_{-1}})$, with \hat{G} the centrally extended loop group

over G . The expressions for the generalised Lie derivative are entirely analogous as proved in [13]. Our result can then also be seen as the extension of the expressions for extended field theory potentials in [14] to the case of affine Kac-Moody groups with scalar fields in indecomposable representations. An especially interesting case is the affine group over $SO(8, n)$ governing two-dimensional half-maximal supergravity. Extended field theories for the duality groups of half-maximal supergravities in four and three dimensions have been recently formulated [63, 64] and capture ten-dimensional heterotic and six-dimensional $(2, 0)$ supergravities as solutions of the section constraint. The potential constructed in this paper corresponds to ten-dimensional $(1, 0)$ and six-dimensional $(2, 0)$ supergravities coupled to abelian supermultiplets. The introduction of gauge interactions for these half-maximal extended field theories requires a deformation of the generalised diffeomorphisms, of the potential and of the full dynamics [63, 65] which would also be interesting to pursue. Along these lines, a further interesting development would be the construction of an ‘ X -deformation’ of our potential (and later of the full E_9 exceptional field theory) that would also reproduce the dynamics of massive type IIA supergravity upon solving the section constraint, in analogy with the higher-dimensional cases [66].

Acknowledgments

We would like to thank Martin Cederwall, Blagoje Oblak and Jakob Palmkvist for useful discussions. This work was partially supported by a PHC PROCOPE, projet No 37667ZL, and by DAAD PPP grant 57316852 (XSUGRA). The work of GB was partially supported by the ANR grant Black-dS-String (ANR-16-CE31-0004). The work of GI was supported by STFC consolidated grant ST/P000754/1.

A Properties of the cocycle

A.1 E_9 group 1-cocycles in the co-adjoint

The group E_9 acts on its Lie algebra \mathfrak{e}_9 by conjugation and we aim to extend this group action on the extra \mathfrak{vir} generators L_m , $m \neq 0$. To this end let

$$X = X_K K + X_0 L_0 + \sum_{n \in \mathbb{Z}} X_A^n T_n^A = X_\alpha T^\alpha \tag{A.1}$$

be an element of \mathfrak{e}_9 in the $R(\Lambda_0)_0$ representation, where $\mathfrak{d} = L_0$. As in this section we are only concerned with the adjoint representation of \mathfrak{e}_9 and its extension by $\langle L_m \rangle$ for fixed $m \neq 0$, we are allowed to ignore the distinction between \mathfrak{d} and L_0 throughout our discussion. Notice also that compared to (4.52), there is no L_{-1} component here. The non-trivial commutator between the Virasoro generators and \mathfrak{e}_9 is given by

$$[X, L_m] = -m X_0 L_m + \sum_{n \in \mathbb{Z}} n X_A^n T_{n+m}^A = -m X_0 L_m + \sum_{n \in \mathbb{Z}} (n - m) X_A^{n-m} T_n^A. \tag{A.2}$$

We see that L_m transforms under \mathfrak{e}_9 by a rescaling proportional to the derivation component of X , plus extra elements in the loop algebra $\hat{\mathfrak{e}}_8$. The same happens for a finite

transformation $g \in E_9$, where we define

$$g^{-1}L_m g \equiv \rho(g)^{-2m}L_m - \omega_{-m}^\alpha(g)\eta_{\alpha\beta}T^\beta, \quad (\text{A.3})$$

where $\rho(g)^2$ is the component of g along the one-dimensional subgroup generated by the derivation:

$$g \equiv \rho(g)^{-2L_0}\hat{g}, \quad \hat{g} \in \hat{E}_8. \quad (\text{A.4})$$

By construction and for fixed m , the algebra $\mathfrak{e}_9 \oplus \langle L_m \rangle$ with commutation relations (A.2) defines a representation of E_9 under the adjoint action. For $X \in \mathfrak{e}_9$ and $e_m \in \mathbb{R}$, no sum over m , one gets

$$g^{-1}(X - e_m L_m)g = \left(g^{-1}Xg + e_m \eta_{\alpha\beta} \omega_{-m}^\alpha(g) T^\beta \right) - \rho(g)^{-2m} e_m L_m, \quad (\text{A.5})$$

such that there is a non-trivial effect of the L_m component on the \mathfrak{e}_9 component X , i.e., the representation matrices are block triangular and the representation is indecomposable. One can understand this representation to be built out of two E_9 representations, the adjoint representation \mathfrak{e}_9 and the one-dimensional representation $\rho(g)^{-2m}$ mentioned above (2.5), linked together by the non-trivial map from E_9 to \mathfrak{e}_9 defined by $g \mapsto \eta_{\alpha\beta} \omega_{-m}^\alpha(g) T^\beta$. The ω_{-m}^α are the components of a map from E_9 to the co-adjoint representation \mathfrak{e}_9^* that one calls a group 1-cocycle. The ω_{-m}^α determine in this sense the extension of the adjoint E_9 representation \mathfrak{e}_9 to the indecomposable E_9 representation $\mathfrak{e}_9 \oplus \langle L_m \rangle$. For the above formula (A.5) to define an action of E_9 , i.e., acting twice being compatible with the group multiplication, the map ω_{-m}^α must satisfy the 1-cocycle condition⁹

$$\omega_{-m}^\alpha(g_1 g_2) = R(g_2^{-1})^\alpha_\beta \omega_{-m}^\beta(g_1) + \rho(g_1)^{-2m} \omega_{-m}^\alpha(g_2), \quad (\text{A.6})$$

for any $g_1, g_2 \in E_9$ and where

$$g^{-1}T^\alpha g = R(g)^\alpha_\beta T^\beta \quad (\text{A.7})$$

defines representation matrices $R(g)^\alpha_\beta$ of the adjoint E_9 action: $R(g_1)^{\alpha\gamma} R(g_2)^\gamma_\beta = R(g_1 g_2)^\alpha_\beta$. Note also that the invariance of $\eta^{\alpha\beta}$ on \mathfrak{e}_9 implies $\eta_{\beta\gamma} R(g^{-1})^\gamma_\delta \eta^{\delta\alpha} = R(g)^\alpha_\beta$. We will discuss the proof of (A.6) momentarily. If the 1-cocycle was trivial, i.e. if there existed a co-adjoint vector v^α such that

$$\omega_{-m}^\alpha(g) \stackrel{?}{=} R^\alpha_\beta(g^{-1})v^\beta - \rho^{-2m}(g)v^\alpha, \quad (\text{A.8})$$

then the representation $\mathfrak{e}_9 \oplus \langle L_m \rangle$ would decompose into the direct sum of \mathfrak{e}_9 and the one-dimensional representation $\rho(g)^{-2m}$, but ω_{-m}^α is a non-trivial cocycle for all m and $\mathfrak{e}_9 \oplus \langle L_m \rangle$ is indecomposable.

The dual of the extended representation can be constructed in the usual way. Denoting the basis dual to $\eta_{\alpha\beta} T^\beta$ and L_m by Λ^α and Λ^m one finds for the action of E_9 that (no sum over m)

$$g^{-1}(j_\alpha \Lambda^\alpha + \chi_m \Lambda^m)g = j_\alpha R(g)^\alpha_\beta \Lambda^\beta + \left(\rho(g)^{2m} \chi_m - j_\alpha \rho(g)^{2m} R(g)^\alpha_\beta \omega_{-m}^\beta(g) \right) \Lambda^m, \quad (\text{A.9})$$

⁹In the standard mathematical definition it is $\rho^{2m}(g)\omega_{-m}^\alpha(g)$ that defines an E_9 1-cocycle in the co-adjoint representation.

from which one can read off the transformation of the coefficients j_α and χ_m . Using (A.6) we find

$$\omega_{-m}^\alpha(g) = -\rho(g)^{-2m} R(g^{-1})^\alpha \beta \omega_{-m}^\beta(g^{-1}) \quad (\text{A.10})$$

so that (A.9) can be rewritten succinctly as follows

$$(j_\alpha, \chi_m) \rightarrow \left(j_\beta R(g)^\beta \alpha, \rho(g)^{2m} \chi_m + \omega_{-m}^\alpha(g^{-1}) j_\alpha \right). \quad (\text{A.11})$$

Setting $m = -1$, this is the transformation we have for the currents and the field χ at $\tilde{\rho} = 0$ (4.13).

An important observation is that $\omega_{-m}^\alpha(g)$ can be defined in terms of the shift operators (2.18). First, we notice that the bilinear forms (2.17) transform under E_9 only by a rescaling

$$\eta_{m\alpha\beta} g^{-1} T^\alpha g \otimes g^{-1} T^\beta g = \rho(g)^{-2m} \eta_{m\alpha\beta} T^\alpha \otimes T^\beta. \quad (\text{A.12})$$

Then, using (2.19) for $n = 0$ and conjugating by g , we obtain for $m \neq 0$,

$$\omega_{-m}^\alpha(g) \mathbf{K} \equiv \rho(g)^{-2m} g \mathcal{S}_m(T^\alpha) g^{-1} - \mathcal{S}_m(g T^\alpha g^{-1}), \quad (\text{A.13})$$

which is therefore equivalent to (A.3) and allows to straightforwardly prove the cocycle condition (A.6). Crucially, (A.13) holds for any $T^\alpha \in \mathfrak{f}$ (with $g \in E_9$) and for this reason we will take it as our definition of $\omega_{-m}^\alpha(g)$. The cocycle condition (A.6) is still satisfied by this more general definition, where (A.7) also generalises to any $T^\alpha \in \mathfrak{f}$ and $\omega_{-m}^\alpha(g)$ is an E_9 group 1-cocycle in the conjugate representation \mathfrak{f}^* .

For the computation of section 4, it is useful to present the expansion of the loop components of the cocycle. Up to linear order in the components of $X \in \mathfrak{e}_9$ of $g = e^X$, we find using (A.13) that¹⁰

$$\omega_{-m}^A(g) = -(m+n) \eta^{AB} X_B^{-m-n} + \mathcal{O}(X^2). \quad (\text{A.14})$$

The expansion of the L_n components of the cocycle $\omega_{-m}^A(g) = \mathcal{O}(X^2)$ only starts at quadratic order in X while the component along \mathbf{K} simply vanishes. Another particularly important expression will be the cocycle associated with the Hermitian coset representative \mathcal{M} at $\tilde{\rho} = 0$. A convenient expression equivalent to (A.13) is

$$\omega_m^\alpha(\mathcal{M}) \mathbf{K} = \rho^{2m} \mathcal{M}^{-1} \mathcal{S}_m(T^{\alpha\dagger}) \mathcal{M} - \mathcal{S}_m(\mathcal{M}^{-1} T^{\alpha\dagger} \mathcal{M}) \quad \text{when } \tilde{\rho} = 0. \quad (\text{A.15})$$

Finally, it is instructive to write explicit expressions for $\omega_{-m}^\alpha(g)$ for $g = e^X$. Here we restrict to $m = \pm 1$ only. Going back to (A.2), it is useful to rewrite the second term as

$$-\sum_{n \in \mathbb{Z}} (n \pm 1) X_A^{n \pm 1} T_n^A = \langle [L_0, X], T^\alpha \rangle_{\pm 1} \eta_{\alpha\beta} T^\beta, \quad (\text{A.16})$$

¹⁰One computes that $[X, \mathcal{S}_m(T_n^A)] + m X_0 \mathcal{S}_m(T_n^A) - \mathcal{S}_m([X, T_n^A]) = -(m+n) \eta^{AB} X_B^{-m-n}$, while $[X, \mathcal{S}_m(L_n)] + m X_0 \mathcal{S}_m(L_n) - \mathcal{S}_m([X, L_n]) = 0$.

where the sum on α runs over all E_9 generators with indices raised and lowered with the standard $\eta_{\alpha\beta} = \eta_{0\alpha\beta}$ of (2.17) and the shifted form on the loop generators in \mathfrak{e}_9 is defined as

$$\langle T_m^A, T_n^B \rangle_{\pm 1} = \eta^{AB} \delta_{m, -n \pm 1} \quad (\text{A.17})$$

and agrees with the restriction to the loop part of the bilinear form $\eta_{\pm 1 \alpha\beta}$. To illustrate (A.16) further, we write out the right-hand side explicitly

$$\begin{aligned} -\langle [L_0, X], T^\alpha \rangle_{\pm 1} \eta_{\alpha\beta} T^\beta &= \langle [L_0, X], \mathbb{K} \rangle_{\pm 1} L_0 + \langle [L_0, X], L_0 \rangle_{\pm 1} \mathbb{K} - \sum_{n \in \mathbb{Z}} \langle [L_0, X], T_{-n}^A \rangle_{\pm 1} T_n^B \eta_{AB} \\ &= \sum_{m, n \in \mathbb{Z}} m X_C^m \langle T_m^C, T_{-n}^A \rangle_{\pm 1} T_n^B \eta_{AB} \\ &= \sum_{n \in \mathbb{Z}} (n \pm 1) X_A^{n \pm 1} T_n^A, \end{aligned} \quad (\text{A.18})$$

where we have used that $\eta_{\pm 1}$ does not pair \mathbb{K} and L_0 non-trivially with anything in \mathfrak{e}_9 .

Using (A.16), we can thus restate (A.2) as

$$\text{ad}_X L_{\pm 1} = \mp X_0 L_{\pm 1} - \langle [L_0, X], T^\alpha \rangle_{\mp 1} \eta_{\alpha\beta} T^\beta \quad (\text{A.19})$$

and we have also introduced the notation $\text{ad}_X L_{\pm 1} = [X, L_{\pm 1}]$ for the commutator between \mathfrak{e}_9 and $L_{\pm 1}$. This is the action we aim to exponentiate.

By induction one can show for any $k \geq 0$ that

$$\text{ad}_X^k L_{\pm 1} = (\mp X_0)^k L_{\pm 1} - \left\langle [L_0, X], \sum_{\ell=0}^{k-1} (\mp X_0)^\ell (-\text{ad}_X)^{k-1-\ell} T^\alpha \right\rangle_{\mp 1} \eta_{\alpha\beta} T^\beta, \quad (\text{A.20})$$

which can be exponentiated to $g = e^X$ easily since X_0 is central in the representation. This leads to

$$g^{-1} L_{\pm 1} g = e^{\pm X_0} L_{\pm 1} - \left\langle [L_0, X], \frac{e^{\text{ad}_X} - e^{\pm X_0}}{-\text{ad}_X \pm X_0} T^\alpha \right\rangle_{\mp 1} \eta_{\alpha\beta} T^\beta, \quad (\text{A.21})$$

which compared to (A.3) gives

$$\rho(g) = e^{-\frac{X_0}{2}}, \quad (\text{A.22a})$$

$$\omega_{\pm}^\alpha(g) = \left\langle [L_0, X], \frac{e^{\text{ad}_X} - e^{\mp X_0}}{-\text{ad}_X \mp X_0} T^\alpha \right\rangle_{\pm 1}, \quad (\text{A.22b})$$

This explicit expression of the cocycle is only valid for $T^\alpha \in \mathfrak{e}_9$.

A.2 Generalisation to $\hat{E}_8 \times \text{SL}(2)$

In the previous discussion g was an element of E_9 . When the axion $\tilde{\rho} \neq 0$, \mathcal{M} is no longer an element of E_9 and instead belongs to the group $\hat{E}_8 \times \text{SL}(2)$. We thus require an extension of some of the formulas above to this case. We shall now give a generalisation of (A.15) that can be expressed as an infinite power series in $\tilde{\rho}$ and reduces to the previous formula when $\tilde{\rho} = 0$.

As in (3.10) we shall decompose the Hermitian \mathcal{M} as

$$\mathcal{M} = m\hat{g}_{\mathcal{M}} = \hat{g}_{\mathcal{M}}^{\dagger}m \tag{A.23}$$

with Hermitian $m \in \text{SL}(2)$ and non-Hermitian $\hat{g}_{\mathcal{M}}$ in \hat{E}_8 . The generalisation of the cocycle that has the needed properties is

$$\Omega^{\alpha}(\mathcal{M})\mathbf{K} = \hat{g}_{\mathcal{M}}^{-1}\mathcal{S}_{+1}\left(m^{-1}T^{\alpha\dagger}m\right)\hat{g}_{\mathcal{M}} - \mathcal{S}_{+1}\left(\mathcal{M}^{-1}T^{\alpha\dagger}\mathcal{M}\right). \tag{A.24}$$

Compared with (A.15), we see that we have made a choice in the split between m and $\hat{g}_{\mathcal{M}}$ and that the character factor ρ does not appear explicitly anymore. This is natural since this part is contained in the action of $m \in \text{SL}(2)$. We have also defined this only for \mathcal{S}_{+1} . Notice that because of the presence of m inside the first shift operator, this expression does not satisfy (A.6) and is therefore not a group cocycle. However, when $\tilde{\rho} = 0$ we have $m \in \mathbb{R}_d^+$ and $\Omega^{\alpha}(\mathcal{M})$ reduces to $\omega_1^{\alpha}(\mathcal{M})$.

Since the T^{α} that belong to \hat{e}_8 can be represented by ϵ_8 elements that depend on a spectral parameter w via $T_m^A = w^m T^A$, leading to meromorphic functions of w , and $\text{SL}(2)$ acts on these generators by Möbius transformations of w , it is convenient to work out the conjugation by m in this picture. More explicitly, the form of the $\text{SL}(2)$ generators as differential operators in w is

$$L_{+1} = -w^2\partial_w, \quad L_0 = -w\partial_w, \quad L_{-1} = -\partial_w, \tag{A.25}$$

such that their exponentiated action on any function $f(w)$ is given by Möbius transformations leading to¹¹

$$m^{-1}f(w)m = f\left(\frac{\rho^{-2}(w - \tilde{\rho})}{1 + \tilde{\rho}\rho^{-2}(w - \tilde{\rho})}\right), \tag{A.26}$$

which when combined with the shift operator \mathcal{S}_{+1} that multiplies by w leads to

$$\begin{aligned} m\mathcal{S}_{+1}(m^{-1}f(w)m)m^{-1} &= \left(\rho^2\frac{w}{1 - \tilde{\rho}w} + \tilde{\rho}\right)f(w) = \rho^2\sum_{k \geq 0}\tilde{\rho}^k w^{k+1}f(w) + \tilde{\rho}f(w) \\ &= \rho^2\sum_{k \geq 0}\tilde{\rho}^k \mathcal{S}_{1+k}(f(w)) + \tilde{\rho}f(w). \end{aligned} \tag{A.27}$$

Inserting this into (A.24) leads to

$$\Omega^{\alpha}(\mathcal{M})\mathbf{K} = \mathcal{M}^{-1}\left(\rho^2\sum_{k \geq 0}\tilde{\rho}^k \mathcal{S}_{1+k}(T^{\alpha\dagger}) + \tilde{\rho}T^{\alpha\dagger}\right)\mathcal{M} - \mathcal{S}_{+1}\left(\mathcal{M}^{-1}T^{\alpha\dagger}\mathcal{M}\right) - \tilde{\rho}\delta_{\mathbf{K}}^{\alpha}\mathbf{K}, \tag{A.28}$$

where the last term is due to the fact that $\Omega^{\mathbf{K}}(\mathcal{M}) = 0$ in (A.24) but the expansion using the Möbius transformations above generates a spurious term.

¹¹The exponentiation of the individual transformations is

$$e^{\tilde{\rho}L_{+1}}f(w)e^{-\tilde{\rho}L_{+1}} = f\left(\frac{w}{1 + \tilde{\rho}w}\right), \quad \rho^{2L_0}f(w)\rho^{-2L_0} = f(\rho^{-2}w), \quad e^{\tilde{\rho}L_{-1}}f(w)e^{-\tilde{\rho}L_{-1}} = f(w - \tilde{\rho}).$$

A.3 Useful identities

Here we collect some useful identities for the generalisation of the cocycle $\Omega^\alpha(\mathcal{M})$ discussed above and the effect of $\text{SL}(2)$ conjugations on expressions appearing in the derivation of the potential.

Using the same argument as above based on Möbius transformations, one works out the conjugation under L_{-1} of the shift operator as

$$e^{\tilde{\rho}L_{-1}}\mathcal{S}_{-1}(e^{-\tilde{\rho}L_{-1}}f(w)e^{\tilde{\rho}L_{-1}})e^{-\tilde{\rho}L_{-1}} = \frac{1}{w-\tilde{\rho}}f(w) \quad (\text{A.29})$$

such that

$$e^{\tilde{\rho}L_{-1}}\mathcal{S}_{-1}(e^{-\tilde{\rho}L_{-1}}T^\alpha e^{\tilde{\rho}L_{-1}})e^{-\tilde{\rho}L_{-1}} = \sum_{k=0}^{\infty} \tilde{\rho}^k \mathcal{S}_{-1-k}(T^\alpha). \quad (\text{A.30})$$

It is sometimes convenient to rewrite the geometric series of shifted generators appearing in \mathcal{J}_α^- in this way.

The \hat{E}_8 invariant bilinear form $\eta_{n\alpha\beta}$ is not invariant under the action of $\text{SL}(2)$. To compute the effect of a conjugation with the Virasoro generator L_q for $q = -1, 0, 1$, it is useful to first show that at the Lie algebra level

$$\eta_{m\alpha\beta}[L_q, T^\alpha] \otimes T^\beta + \eta_{m\alpha\beta}T^\alpha \otimes [L_q, T^\beta] = (q-n)\eta_{m+q\alpha\beta}T^\alpha \otimes T^\beta. \quad (\text{A.31})$$

Using this formula, one computes that for $m \in \text{SL}(2)$ one has

$$\begin{aligned} \eta_{-1\alpha\beta}m^{-1}T^{\alpha\dagger}m \otimes T^\beta &= \left(\frac{1}{\rho^2}\eta_{1\alpha\beta} - 2\frac{\tilde{\rho}}{\rho^2}\eta_{\alpha\beta} + \frac{\tilde{\rho}^2}{\rho^2}\eta_{-1\alpha\beta} \right) T^\alpha \otimes m^{-1}T^{\beta\dagger}m \quad (\text{A.32}) \\ \eta_{\alpha\beta}m^{-1}T^{\alpha\dagger}m \otimes T^\beta &= \left(\frac{\tilde{\rho}}{\rho^2}\eta_{1\alpha\beta} + \left(1 - 2\frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{\alpha\beta} - \tilde{\rho}\left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{-1\alpha\beta} \right) T^\alpha \otimes m^{-1}T^{\beta\dagger}m \\ \eta_{1\alpha\beta}m^{-1}T^{\alpha\dagger}m \otimes T^\beta &= \left(\frac{\tilde{\rho}^2}{\rho^2}\eta_{1\alpha\beta} + 2\tilde{\rho}\left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{\alpha\beta} + \rho^2\left(1 - \frac{\tilde{\rho}^2}{\rho^2}\right)^2\eta_{-1\alpha\beta} \right) T^\alpha \otimes m^{-1}T^{\beta\dagger}m \\ \eta_{2\alpha\beta}m^{-1}T^{\alpha\dagger}m \otimes T^\beta &= \left(\frac{\tilde{\rho}^3}{\rho^2}\eta_{1\alpha\beta} + \tilde{\rho}^2\left(3 - 2\frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{\alpha\beta} + \tilde{\rho}\left(3\rho^2 - 3\tilde{\rho}^2 + \frac{\tilde{\rho}^4}{\rho^2}\right)\eta_{-1\alpha\beta} \right. \\ &\quad \left. + \rho^4\sum_{k=0}^{\infty}\tilde{\rho}^k\eta_{-2-k\alpha\beta} \right) T^\alpha \otimes m^{-1}T^{\beta\dagger}m. \end{aligned}$$

Note in particular that

$$(\eta_{\alpha\beta} - \tilde{\rho}\eta_{-1\alpha\beta})m^{-1}T^{\alpha\dagger}m \otimes T^\beta = (\eta_{\alpha\beta} - \tilde{\rho}\eta_{-1\alpha\beta})T^\alpha \otimes m^{-1}T^{\beta\dagger}m. \quad (\text{A.33})$$

These formulas can be used to obtain some properties of $\Omega^\alpha(\mathcal{M})$. One computes using the definition (A.24) that

$$\begin{aligned}
 \eta_{\alpha\beta}\Omega^\alpha(\mathcal{M})\mathbb{K}\otimes T^\beta &= \eta_{\alpha\beta}\left(\hat{g}_{\mathcal{M}}^{-1}\mathcal{S}_1\left(m^{-1}T^{\alpha\dagger}m\right)\hat{g}_{\mathcal{M}}-\mathcal{S}_1\left(\hat{g}_{\mathcal{M}}^{-1}m^{-1}T^{\alpha\dagger}m\hat{g}_{\mathcal{M}}\right)\right)\otimes T^\beta \quad (\text{A.34}) \\
 &= \left(\frac{\tilde{\rho}}{\rho^2}\eta_{1\alpha\beta}+\left(1-2\frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{\alpha\beta}-\tilde{\rho}\left(1-\frac{\tilde{\rho}^2}{\rho^2}\right)\eta_{-1\alpha\beta}\right) \\
 &\quad \times \left(\hat{g}_{\mathcal{M}}^{-1}\mathcal{S}_1(T^\alpha)\hat{g}_{\mathcal{M}}\otimes m^{-1}T^{\beta\dagger}m-\mathcal{S}_1(T^\alpha)\otimes\hat{g}_{\mathcal{M}}^{-1}m^{-1}T^{\beta\dagger}m\hat{g}_{\mathcal{M}}\right) \\
 &= \mathbb{K}\otimes\left(\tilde{\rho}(L_0-\hat{g}_{\mathcal{M}}^{-1}L_0\hat{g}_{\mathcal{M}})+\rho^2\sum_{k=0}^{\infty}\tilde{\rho}^k(L_{-1-k}-\hat{g}_{\mathcal{M}}^{-1}L_{-1-k}\hat{g}_{\mathcal{M}})\right),
 \end{aligned}$$

where we used (A.32) in the first step and (2.19) in the second. One therefore obtains the useful identity

$$\eta_{\alpha\beta}\Omega^\alpha(\mathcal{M})T^\beta = \tilde{\rho}(L_0 - \hat{g}_{\mathcal{M}}^{-1}L_0\hat{g}_{\mathcal{M}}) + \rho^2 \sum_{k=0}^{\infty} \tilde{\rho}^k (L_{-1-k} - \hat{g}_{\mathcal{M}}^{-1}L_{-1-k}\hat{g}_{\mathcal{M}}). \quad (\text{A.35})$$

We want also to compute $\eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma\Omega^\gamma(\mathcal{M})T^\beta$ with $H(\mathcal{M})^\alpha{}_\beta$ defined in (3.27). Using instead formula (A.28), one computes that

$$\begin{aligned}
 \eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma\Omega^\gamma(\mathcal{M})\mathbb{K}\otimes T^\beta &= \eta_{\alpha\beta}\left(\mathcal{M}^{-1}\rho^2\sum_{k=0}^{\infty}\tilde{\rho}^k\mathcal{S}_{-1-k}\left(\mathcal{M}^{-1}T^{\alpha\dagger}\mathcal{M}\right)^\dagger\mathcal{M}+\tilde{\rho}T^\alpha-\mathcal{S}_1(T^\alpha)\right)\otimes T^\beta \\
 &\quad + \tilde{\rho}\mathbb{K}\otimes\left(\frac{\tilde{\rho}}{\rho^2}\mathcal{M}^{-1}L_{-1}\mathcal{M}+\left(1-2\frac{\tilde{\rho}^2}{\rho^2}\right)\mathcal{M}^{-1}L_0\mathcal{M}-\tilde{\rho}\left(1-\frac{\tilde{\rho}^2}{\rho^2}\right)\mathcal{M}^{-1}L_1\mathcal{M}\right) \\
 &= \mathbb{K}\otimes\left(\hat{g}_{\mathcal{M}}^{-1}L_1\hat{g}_{\mathcal{M}}-L_1\right) \quad (\text{A.36})
 \end{aligned}$$

so one obtains the very simple result that

$$\eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma\Omega^\gamma(\mathcal{M})T^\beta = \hat{g}_{\mathcal{M}}^{-1}L_1\hat{g}_{\mathcal{M}} - L_1. \quad (\text{A.37})$$

Using this formula and reinserting the m matrix using (A.32) one obtains that

$$L_1 + \eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma\Omega^\gamma(\mathcal{M})T^\beta = \mathcal{M}^{-1}\left(\rho^2\left(1-\frac{\tilde{\rho}^2}{\rho^2}\right)^2L_1+2\tilde{\rho}\left(1-\frac{\tilde{\rho}^2}{\rho^2}\right)L_0+\frac{\tilde{\rho}^2}{\rho^2}L_{-1}\right)\mathcal{M}. \quad (\text{A.38})$$

Using again (A.32) one computes that

$$\eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma\mathbb{K}T^\beta = \mathcal{M}^{-1}\left(\tilde{\rho}\left(1-\frac{\tilde{\rho}^2}{\rho^2}\right)L_1-\left(1-2\frac{\tilde{\rho}^2}{\rho^2}\right)L_0-\frac{\tilde{\rho}}{\rho^2}L_{-1}\right)\mathcal{M}, \quad (\text{A.39})$$

such that

$$L_1 + \eta_{\alpha\beta}H(\mathcal{M})^\alpha{}_\gamma(\Omega^\gamma(\mathcal{M}) + \tilde{\rho}\delta_{\mathbb{K}}^\gamma)T^\beta = \mathcal{M}^{-1}\left((\rho^2 - \tilde{\rho}^2)L_1 + \tilde{\rho}L_0\right)\mathcal{M}. \quad (\text{A.40})$$

By a similar reasoning, one also obtains

$$\eta_{-1\alpha\beta}\Omega^\alpha(\mathcal{M})T^\beta = \tilde{\rho}L_{-1} + \rho^2\sum_{k=0}^{\infty}\tilde{\rho}^kL_{-2-k} - \hat{g}_{\mathcal{M}}^{-1}\left(\tilde{\rho}L_{-1} + \rho^2\sum_{k=0}^{\infty}\tilde{\rho}^kL_{-2-k}\right)\hat{g}_{\mathcal{M}}, \quad (\text{A.41})$$

and

$$\eta_{-1\alpha\beta}\Omega^\gamma(\mathcal{M})H(\mathcal{M})^\alpha{}_\gamma T^\beta = \hat{g}_{\mathcal{M}}^{-1}L_0\hat{g}_{\mathcal{M}} - L_0. \quad (\text{A.42})$$

B Gauge algebra closure on $\langle \chi |$

The closure of the algebra of generalised diffeomorphisms on the $\hat{E}_8 \rtimes (\mathbb{R}_d^+ \ltimes \mathbb{R}_{L-1})$ scalars \mathcal{M} follows from its closure on a vector field $|V\rangle$ (3.15), which was derived in [13]. The field $\langle \chi |$ does not transform under generalised diffeomorphisms simply as a generalised Lie derivative, and the mixing with the $\hat{\mathfrak{e}}_8 \oplus \mathfrak{sl}(2)$ current makes it non-obvious that the algebra closes on this field. In this appendix we show that this is indeed the case.

Because the algebra closes on \mathcal{M} and therefore on $\langle \mathcal{J}_\alpha |$, one can check the closure of the algebra on any linear combination of $\langle \chi |$ and $\langle \mathcal{J}_\alpha |$. Since the transformation of $\langle \chi |$ is not manifestly covariant, it is indeed convenient to check the closure of the algebra on the combination

$$\langle \xi | \equiv \langle \chi | + \sum_{k=0}^{\infty} \tilde{\rho}^k \langle \mathcal{J}_\alpha | \mathcal{S}_{-1-k}(T^\alpha) + \frac{\tilde{\rho}}{\rho^2} (\langle \mathcal{J}_\alpha | T^\alpha + \rho^{-1} \langle \partial \rho |), \quad (\text{B.1})$$

that transforms as

$$\begin{aligned} \delta \langle \xi | &= \langle \partial_\xi | \Lambda \rangle \langle \xi | + \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \xi | T^\beta + \frac{1}{\rho^2} \eta_{1\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle \langle \partial_\Lambda | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\ &+ \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \xi | T^\beta + \frac{1}{\rho^2} \eta_{\alpha\beta} \text{Tr}(T^\alpha \Sigma) \langle \partial_\Sigma | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\ &- \frac{1}{\rho^2} \langle \partial_\Sigma | \Sigma + \frac{1}{\rho^2} \text{Tr}(\Sigma) (\langle \mathcal{J}_\alpha | T^\alpha + \rho^{-1} \langle \partial \rho |). \end{aligned} \quad (\text{B.2})$$

To compute the closure it is convenient to use the BRST formalism, for which Λ and Σ are understood as anticommuting ghost fields, with their own variation defined according to (3.21) as

$$\begin{aligned} \delta |\Lambda\rangle &= \frac{1}{2} \left(\langle \partial_{\Lambda_1} | \Lambda_2 \rangle | \Lambda_1 \rangle - \eta_{\alpha\beta} \langle \partial_{\Lambda_2} | T^\alpha | \Lambda_2 \rangle T^\beta | \Lambda_1 \rangle - \langle \partial_{\Lambda_2} | \Lambda_2 \rangle | \Lambda_1 \rangle \right), \\ \delta \Sigma &= \langle \partial_\Sigma | \Lambda \rangle \Sigma - \eta_{\alpha\beta} \langle \partial_\Lambda | T^\alpha | \Lambda \rangle T^\beta \Sigma - \Sigma | \Lambda \rangle \langle \partial_\Lambda | \\ &+ \frac{1}{4} \eta_{1\alpha\beta} \left(\langle \partial_{\Lambda_2} | T^\alpha | \Lambda_2 \rangle T^\beta | \Lambda_1 \rangle + \langle \partial_{\Lambda_2} | T^\alpha | \Lambda_1 \rangle T^\beta | \Lambda_2 \rangle \right) \langle \partial_{\Lambda_2} | - \frac{1}{2} \eta_{-1\alpha\beta} \text{Tr}(T^\alpha \Sigma) T^\beta \Sigma, \end{aligned} \quad (\text{B.3})$$

for which the labels on $|\Lambda_1\rangle$ and $|\Lambda_2\rangle$ only indicate on which $|\Lambda\rangle$ the derivative acts, despite the fact that they are the same anticommuting ghost $|\Lambda\rangle$. For example, in index notation one has

$$\delta \Lambda^M = \frac{1}{2} \left(\Lambda^N \partial_N \Lambda^M - \eta_{\alpha\beta} (T^\alpha)^P{}_N \partial_P \Lambda^N (T^\beta)^M{}_Q \Lambda^Q - \partial_N \Lambda^N \Lambda^M \right). \quad (\text{B.4})$$

In this notation, the closure on the algebra on a vector field $|V\rangle$ is equivalent to the property that $\delta^2 |V\rangle = 0$. Note that $\delta^2 |\Lambda\rangle \neq 0$, but it gives a trivial generalised diffeomorphism, whereas the definition of a truly nilpotent operator requires the introduction of an infinite chain of ghosts for ghosts generating an L_∞ algebra structure [7, 67, 68].

In the BRST formulation it is easier to check that $\delta^2 \langle \xi |$ indeed vanishes. Here, we shall only give some of the steps for the terms quadratic in $|\Lambda\rangle$. The parts of the transformations

corresponding to the Lie derivative of an ordinary vector field work as in [13], whereas the other give the following contributions

$$\begin{aligned}
\delta^2 \langle \xi | &= \eta_{1\alpha\beta} \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \langle \partial_1 | \Lambda_2 \rangle \langle \partial_1 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad - \frac{1}{4} \eta_{\alpha\beta} \eta_{1\gamma\delta} \left(\langle \partial_2 | [T^\alpha, T^\gamma] | \Lambda_1 \rangle \langle \partial_2 | T^\delta | \Lambda_2 \rangle + \langle \partial_2 | [T^\alpha, T^\gamma] | \Lambda_2 \rangle \langle \partial_2 | T^\delta | \Lambda_1 \rangle \right) \langle \partial_1 + \partial_2 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad + \frac{1}{4} \eta_{1\alpha\beta} \left(\langle \partial_1 | T^\alpha | \Lambda_2 \rangle \langle \partial_2 | T^\beta | \Lambda_1 \rangle + \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \langle \partial_2 | T^\beta | \Lambda_2 \rangle \right) \langle \partial_2 | \\
&\quad + \eta_{1\alpha\beta} \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \left(\langle \partial_2 | \Lambda_2 \rangle \langle \partial_1 | + \langle \partial_1 | \Lambda_2 \rangle \langle \partial_2 | \right) \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad + \eta_{1\alpha\beta} \eta_{\gamma\delta} \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \langle \partial_2 | T^\gamma | \Lambda_2 \rangle \langle \partial_1 | \mathcal{M}^{-1} [T^\beta, T^\delta]^\dagger \mathcal{M} \\
&\quad + \eta_{1\alpha\beta} \eta_{\gamma\delta} \langle \partial_1 + \partial_2 | [T^\alpha, T^\gamma] | \Lambda_1 \rangle \langle \partial_2 | T^\gamma | \Lambda_2 \rangle \langle \partial_1 + \partial_2 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad + \frac{1}{2} \eta_{1\alpha\beta} \left(\langle \partial_2 | T^\alpha | \Lambda_1 \rangle \langle \partial_1 | \Lambda_2 \rangle - \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \langle \partial_2 | \Lambda_2 \rangle \right) \langle \partial_1 + \partial_2 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad + \frac{1}{2} \eta_{1\alpha\beta} \langle \partial_1 + \partial_2 | T^\alpha | \Lambda_1 \rangle \langle \partial_2 - \partial_1 | \Lambda_2 \rangle \langle \partial_1 + \partial_2 | \\
&= 0,
\end{aligned} \tag{B.5}$$

where $\langle \partial_1 | = \langle \partial_{\Lambda_1} |$ and $\langle \partial_2 | = \langle \partial_{\Lambda_2} |$ for short, and one uses that all the terms involving commutators simplify according to

$$\begin{aligned}
&\eta_{1\alpha\beta} \eta_{\gamma\delta} \left(- \langle \partial_2 | [T^\alpha, T^\gamma] | \Lambda_1 \rangle \langle \partial_2 | - \langle \partial_1 | [T^\alpha, T^\gamma] | \Lambda_1 \rangle \langle \partial_1 | + 2 \langle \partial_1 + \partial_2 | [T^\beta, T^\gamma] | \Lambda_1 \rangle \langle \partial_2 | \right) T^\delta | \Lambda_2 \rangle \\
&\quad \times \langle \partial_1 + \partial_2 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&\quad - 4 \eta_{1\alpha\beta} \eta_{\gamma\delta} \langle \partial_1 | [T^\alpha, T^\gamma] | \Lambda_1 \rangle \langle \partial_2 | T^\delta | \Lambda_2 \rangle \langle \partial_1 | \mathcal{M}^{-1} T^{\beta\dagger} \mathcal{M} \\
&= \eta_{1\alpha\beta} \langle \partial_1 | T^\alpha | \Lambda_1 \rangle \langle \partial_2 | T^\beta | \Lambda_2 \rangle \langle \partial_1 - \partial_2 |.
\end{aligned} \tag{B.6}$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] C.M. Hull, *Generalised Geometry for M-theory*, *JHEP* **07** (2007) 079 [[hep-th/0701203](https://arxiv.org/abs/hep-th/0701203)] [[INSPIRE](https://arxiv.org/abs/hep-th/0701203)].
- [2] P. Pires Pacheco and D. Waldram, *M-theory, exceptional generalised geometry and superpotentials*, *JHEP* **09** (2008) 123 [[arXiv:0804.1362](https://arxiv.org/abs/0804.1362)] [[INSPIRE](https://arxiv.org/abs/0804.1362)].
- [3] C. Hillmann, *$E_{7(7)}$ and $d = 11$ supergravity*, Ph.D. thesis, Humboldt University Berlin, Germany, (2008), [arXiv:0902.1509](https://arxiv.org/abs/0902.1509) [[INSPIRE](https://arxiv.org/abs/0902.1509)].
- [4] D.S. Berman and M.J. Perry, *Generalized Geometry and M-theory*, *JHEP* **06** (2011) 074 [[arXiv:1008.1763](https://arxiv.org/abs/1008.1763)] [[INSPIRE](https://arxiv.org/abs/1008.1763)].
- [5] D.S. Berman, H. Godazgar and M.J. Perry, *SO(5,5) duality in M-theory and generalized geometry*, *Phys. Lett. B* **700** (2011) 65 [[arXiv:1103.5733](https://arxiv.org/abs/1103.5733)] [[INSPIRE](https://arxiv.org/abs/1103.5733)].
- [6] A. Coimbra, C. Strickland-Constable and D. Waldram, *$E_{d(d)} \times \mathbb{R}^+$ generalised geometry, connections and M-theory*, *JHEP* **02** (2014) 054 [[arXiv:1112.3989](https://arxiv.org/abs/1112.3989)] [[INSPIRE](https://arxiv.org/abs/1112.3989)].

- [7] D.S. Berman, M. Cederwall, A. Kleinschmidt and D.C. Thompson, *The gauge structure of generalised diffeomorphisms*, *JHEP* **01** (2013) 064 [[arXiv:1208.5884](#)] [[INSPIRE](#)].
- [8] A. Coimbra, C. Strickland-Constable and D. Waldram, *Supergravity as Generalised Geometry II: $E_{d(d)} \times \mathbb{R}^+$ and M-theory*, *JHEP* **03** (2014) 019 [[arXiv:1212.1586](#)] [[INSPIRE](#)].
- [9] J.-H. Park and Y. Suh, *U-geometry: $SL(5)$* , *JHEP* **04** (2013) 147 [Erratum *ibid.* **11** (2013) 210] [[arXiv:1302.1652](#)] [[INSPIRE](#)].
- [10] M. Cederwall, J. Edlund and A. Karlsson, *Exceptional geometry and tensor fields*, *JHEP* **07** (2013) 028 [[arXiv:1302.6736](#)] [[INSPIRE](#)].
- [11] M. Cederwall, *Non-gravitational exceptional supermultiplets*, *JHEP* **07** (2013) 025 [[arXiv:1302.6737](#)] [[INSPIRE](#)].
- [12] G. Aldazabal, M. Graña, D. Marqués and J.A. Rosabal, *Extended geometry and gauged maximal supergravity*, *JHEP* **06** (2013) 046 [[arXiv:1302.5419](#)] [[INSPIRE](#)].
- [13] G. Bossard, M. Cederwall, A. Kleinschmidt, J. Palmkvist and H. Samtleben, *Generalized diffeomorphisms for E_9* , *Phys. Rev. D* **96** (2017) 106022 [[arXiv:1708.08936](#)] [[INSPIRE](#)].
- [14] M. Cederwall and J. Palmkvist, *Extended geometries*, *JHEP* **02** (2018) 071 [[arXiv:1711.07694](#)] [[INSPIRE](#)].
- [15] E. Cremmer and B. Julia, *The $N = 8$ Supergravity Theory. 1. The Lagrangian*, *Phys. Lett. B* **80** (1978) 48 [[INSPIRE](#)].
- [16] B. Julia, *Kac-Moody symmetry of gravitation and supergravity theories*, *Lectures Appl. Math.* **21** (1985) 335.
- [17] H. Nicolai, *The Integrability of $N = 16$ Supergravity*, *Phys. Lett. B* **194** (1987) 402 [[INSPIRE](#)].
- [18] E. Cremmer, B. Julia, H. Lü and C.N. Pope, *Dualization of dualities. 1.*, *Nucl. Phys. B* **523** (1998) 73 [[hep-th/9710119](#)] [[INSPIRE](#)].
- [19] E. Cremmer, B. Julia, H. Lü and C.N. Pope, *Dualization of dualities. 2. Twisted self-duality of doubled fields and superdualities*, *Nucl. Phys. B* **535** (1998) 242 [[hep-th/9806106](#)] [[INSPIRE](#)].
- [20] B. de Wit and H. Samtleben, *Gauged maximal supergravities and hierarchies of nonAbelian vector-tensor systems*, *Fortsch. Phys.* **53** (2005) 442 [[hep-th/0501243](#)] [[INSPIRE](#)].
- [21] B. de Wit, H. Nicolai and H. Samtleben, *Gauged Supergravities, Tensor Hierarchies and M-theory*, *JHEP* **02** (2008) 044 [[arXiv:0801.1294](#)] [[INSPIRE](#)].
- [22] O. Hohm and H. Samtleben, *Exceptional Field Theory I: $E_{6(6)}$ covariant Form of M-theory and Type IIB*, *Phys. Rev. D* **89** (2014) 066016 [[arXiv:1312.0614](#)] [[INSPIRE](#)].
- [23] O. Hohm and H. Samtleben, *Exceptional field theory. II. $E_{7(7)}$* , *Phys. Rev. D* **89** (2014) 066017 [[arXiv:1312.4542](#)] [[INSPIRE](#)].
- [24] O. Hohm and H. Samtleben, *Exceptional field theory. III. $E_{8(8)}$* , *Phys. Rev. D* **90** (2014) 066002 [[arXiv:1406.3348](#)] [[INSPIRE](#)].
- [25] D.S. Berman, E.T. Musaev, D.C. Thompson and D.C. Thompson, *Duality Invariant M-theory: Gauged supergravities and Scherk-Schwarz reductions*, *JHEP* **10** (2012) 174 [[arXiv:1208.0020](#)] [[INSPIRE](#)].
- [26] E.T. Musaev, *Gauged supergravities in 5 and 6 dimensions from generalised Scherk-Schwarz reductions*, *JHEP* **05** (2013) 161 [[arXiv:1301.0467](#)] [[INSPIRE](#)].

- [27] H. Godazgar, M. Godazgar and H. Nicolai, *Embedding tensor of Scherk-Schwarz flux compactifications from eleven dimensions*, *Phys. Rev. D* **89** (2014) 045009 [[arXiv:1312.1061](#)] [[INSPIRE](#)].
- [28] K. Lee, C. Strickland-Constable and D. Waldram, *Spheres, generalised parallelisability and consistent truncations*, *Fortsch. Phys.* **65** (2017) 1700048 [[arXiv:1401.3360](#)] [[INSPIRE](#)].
- [29] O. Hohm and H. Samtleben, *Consistent Kaluza-Klein Truncations via Exceptional Field Theory*, *JHEP* **01** (2015) 131 [[arXiv:1410.8145](#)] [[INSPIRE](#)].
- [30] B. Julia, *Infinite Lie algebras in physics*, proceedings of *Johns Hopkins Workshop on Current Problems in Particle Theory*, LPTENS-81-14 [[INSPIRE](#)].
- [31] H. Samtleben and M. Weidner, *Gauging hidden symmetries in two dimensions*, *JHEP* **08** (2007) 076 [[arXiv:0705.2606](#)] [[INSPIRE](#)].
- [32] J. de Boer and M. Shigemori, *Exotic branes and non-geometric backgrounds*, *Phys. Rev. Lett.* **104** (2010) 251603 [[arXiv:1004.2521](#)] [[INSPIRE](#)].
- [33] J. de Boer and M. Shigemori, *Exotic Branes in String Theory*, *Phys. Rept.* **532** (2013) 65 [[arXiv:1209.6056](#)] [[INSPIRE](#)].
- [34] I. Bakhmatov, D. Berman, A. Kleinschmidt, E. Musaev and R. Otsuki, *Exotic branes in Exceptional Field Theory: the $SL(5)$ duality group*, *JHEP* **08** (2018) 021 [[arXiv:1710.09740](#)] [[INSPIRE](#)].
- [35] J.J. Fernández-Melgarejo, T. Kimura and Y. Sakatani, *Weaving the Exotic Web*, *JHEP* **09** (2018) 072 [[arXiv:1805.12117](#)] [[INSPIRE](#)].
- [36] D.S. Berman, E.T. Musaev and R. Otsuki, *Exotic Branes in Exceptional Field Theory: $E_{7(7)}$ and Beyond*, *JHEP* **12** (2018) 053 [[arXiv:1806.00430](#)] [[INSPIRE](#)].
- [37] P. Goddard and D.I. Olive, *Kac-Moody and Virasoro Algebras in Relation to Quantum Physics*, *Int. J. Mod. Phys. A* **1** (1986) 303 [[INSPIRE](#)].
- [38] H. Sugawara, *A field theory of currents*, *Phys. Rev.* **170** (1968) 1659 [[INSPIRE](#)].
- [39] B. Julia and H. Nicolai, *Conformal internal symmetry of 2-D σ -models coupled to gravity and a dilaton*, *Nucl. Phys. B* **482** (1996) 431 [[hep-th/9608082](#)] [[INSPIRE](#)].
- [40] R.P. Geroch, *A method for generating solutions of Einstein's equations*, *J. Math. Phys.* **12** (1971) 918 [[INSPIRE](#)].
- [41] P. Breitenlohner and D. Maison, *On the Geroch group*, *Ann. Inst. H. Poincaré Phys. Theor.* **46** (1987) 215.
- [42] P. Breitenlohner, D. Maison and G.W. Gibbons, *Four-Dimensional Black Holes from Kaluza-Klein Theories*, *Commun. Math. Phys.* **120** (1988) 295 [[INSPIRE](#)].
- [43] H. Nicolai and H. Samtleben, *On $K(E_9)$* , *Q. J. Pure Appl. Math.* **1** (2005) 180 [[hep-th/0407055](#)] [[INSPIRE](#)].
- [44] E. Cremmer, H. Lü, C.N. Pope and K.S. Stelle, *Spectrum generating symmetries for BPS solitons*, *Nucl. Phys. B* **520** (1998) 132 [[hep-th/9707207](#)] [[INSPIRE](#)].
- [45] V.G. Kac and D.H. Peterson, *Defining relations of certain infinite dimensional groups*, *Astérisque Hors-Série*, (1984), pp. 165–208.
- [46] J. Tits, *Uniqueness and presentation of Kac-Moody groups over fields*, *J. Algebra* **105** (1987) 542.
- [47] L. Carbone and H. Garland, *Existence of lattices in Kac-Moody groups over finite fields*, *Commun. Contemp. Math.* **5** (2003) 813.

- [48] T. De Medts, R. Gramlich, M. Horn *Iwasawa decompositions of split Kac-Moody groups*, *J. Lie Theory* **19** (2009) 311.
- [49] O. Hohm and H. Samtleben, *Exceptional Form of $D = 11$ Supergravity*, *Phys. Rev. Lett.* **111** (2013) 231601 [[arXiv:1308.1673](#)] [[INSPIRE](#)].
- [50] C.D.A. Blair, E. Malek and J.-H. Park, *M-theory and Type IIB from a Duality Manifest Action*, *JHEP* **01** (2014) 172 [[arXiv:1311.5109](#)] [[INSPIRE](#)].
- [51] G. Bossard and A. Kleinschmidt, *Loops in exceptional field theory*, *JHEP* **01** (2016) 164 [[arXiv:1510.07859](#)] [[INSPIRE](#)].
- [52] H. Godazgar, M. Godazgar and M.J. Perry, *E_8 duality and dual gravity*, *JHEP* **06** (2013) 044 [[arXiv:1303.2035](#)] [[INSPIRE](#)].
- [53] V.A. Belinsky and V.E. Zakharov, *Integration of the Einstein Equations by the Inverse Scattering Problem Technique and the Calculation of the Exact Soliton Solutions*, *Sov. Phys. JETP* **48** (1978) 985 [*Zh. Eksp. Teor. Fiz.* **75** (1978) 1953] [[INSPIRE](#)].
- [54] D. Maison, *Are the stationary, axially symmetric Einstein equations completely integrable?*, *Phys. Rev. Lett.* **41** (1978) 521 [[INSPIRE](#)].
- [55] D. Bernard and B. Julia, *Twisted self-duality of dimensionally reduced gravity and vertex operators*, *Nucl. Phys. B* **547** (1999) 427 [[hep-th/9712254](#)] [[INSPIRE](#)].
- [56] M. Cederwall and J.A. Rosabal, *E_8 geometry*, *JHEP* **07** (2015) 007 [[arXiv:1504.04843](#)] [[INSPIRE](#)].
- [57] A. Baguet and H. Samtleben, *$E_{8(8)}$ Exceptional Field Theory: Geometry, Fermions and Supersymmetry*, *JHEP* **09** (2016) 168 [[arXiv:1607.03119](#)] [[INSPIRE](#)].
- [58] P.C. West, *E_{11} and M-theory*, *Class. Quant. Grav.* **18** (2001) 4443 [[hep-th/0104081](#)] [[INSPIRE](#)].
- [59] P. West, *Generalised Space-time and Gauge Transformations*, *JHEP* **08** (2014) 050 [[arXiv:1403.6395](#)] [[INSPIRE](#)].
- [60] A.G. Tumanov and P. West, *E_{11} in 11D*, *Phys. Lett. B* **758** (2016) 278 [[arXiv:1601.03974](#)] [[INSPIRE](#)].
- [61] T. Damour, M. Henneaux and H. Nicolai, *E_{10} and a ‘small tension expansion’ of M-theory*, *Phys. Rev. Lett.* **89** (2002) 221601 [[hep-th/0207267](#)] [[INSPIRE](#)].
- [62] G. Bossard, A. Kleinschmidt, J. Palmkvist, C.N. Pope and E. Sezgin, *Beyond E_{11}* , *JHEP* **05** (2017) 020 [[arXiv:1703.01305](#)] [[INSPIRE](#)].
- [63] F. Ciceri, G. Dibitetto, J.J. Fernández-Melgarejo, A. Guarino and G. Inverso, *Double Field Theory at $SL(2)$ angles*, *JHEP* **05** (2017) 028 [[arXiv:1612.05230](#)] [[INSPIRE](#)].
- [64] O. Hohm, E.T. Musaev and H. Samtleben, *$O(d+1, d+1)$ enhanced double field theory*, *JHEP* **10** (2017) 086 [[arXiv:1707.06693](#)] [[INSPIRE](#)].
- [65] O. Hohm and S.K. Kwak, *Double Field Theory Formulation of Heterotic Strings*, *JHEP* **06** (2011) 096 [[arXiv:1103.2136](#)] [[INSPIRE](#)].
- [66] F. Ciceri, A. Guarino and G. Inverso, *The exceptional story of massive IIA supergravity*, *JHEP* **08** (2016) 154 [[arXiv:1604.08602](#)] [[INSPIRE](#)].
- [67] O. Hohm and B. Zwiebach, *L_∞ Algebras and Field Theory*, *Fortsch. Phys.* **65** (2017) 1700014 [[arXiv:1701.08824](#)] [[INSPIRE](#)].
- [68] M. Cederwall and J. Palmkvist, *L_∞ algebras for extended geometry from Borcherds superalgebras*, [arXiv:1804.04377](#) [[INSPIRE](#)].