

# Supporting Information: Determining Free Energy Differences Through Variational Intermediates

Martin Reinhardt and Helmut Grubmüller\*

*Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen,  
Germany*

E-mail: hgrubmu@gwdg.de

Phone: +49 551 201 2300. Fax: +49 551 201 2302

Table 1: Mean squared errors (MSE) as a function of total simulation time for the electrostatic decoupling of solvated butanol. All MSEs are given in units of  $[(k_B T)^2]$ . The linear intermediates are compared to three variants of VI. Firstly, with an exact initial estimate of the free energy difference, secondly, with an estimate that is  $1 k_B T$  smaller than the exact reference value, and, thirdly, with an estimate of  $0 k_B T$ , i.e.,  $8.708 k_B T$  smaller than the reference value. For the latter two variants see continuation in Table 2.

sim. time [ps]	linear	VI, $C$ exact
20	$(6.46 \pm 0.04) \cdot 10^{-1}$	$(3.89 \pm 0.04) \cdot 10^{-1}$
40	$(3.50 \pm 0.03) \cdot 10^{-1}$	$(2.22 \pm 0.04) \cdot 10^{-1}$
100	$(1.49 \pm 0.02) \cdot 10^{-1}$	$(7.3 \pm 0.2) \cdot 10^{-2}$
200	$(7.7 \pm 0.2) \cdot 10^{-2}$	$(3.40 \pm 0.08) \cdot 10^{-2}$
400	$(3.8 \pm 0.1) \cdot 10^{-2}$	$(1.57 \pm 0.04) \cdot 10^{-2}$
1000	$(1.5 \pm 0.1) \cdot 10^{-2}$	$(6.6 \pm 0.3) \cdot 10^{-3}$
2000	$(7.5 \pm 0.5) \cdot 10^{-3}$	$(3.4 \pm 0.2) \cdot 10^{-3}$
4000	$(3.7 \pm 0.3) \cdot 10^{-3}$	$(1.8 \pm 0.2) \cdot 10^{-3}$
10000	$(1.4 \pm 0.1) \cdot 10^{-3}$	$(8 \pm 1) \cdot 10^{-4}$

Table 2: Continuation of Table 1

sim. time [ps]	VI, $\Delta C = 1k_B T$	VI, $C = 0$
20	$(4.2 \pm 0.1) \cdot 10^{-1}$	$14.61 \pm 0.06$
40	$(2.6 \pm 0.1) \cdot 10^{-1}$	$8.36 \pm 0.06$
100	$(8.2 \pm 0.4) \cdot 10^{-2}$	$4.05 \pm 0.05$
200	$(4.1 \pm 0.2) \cdot 10^{-2}$	$2.82 \pm 0.05$
400	$(2.0 \pm 0.1) \cdot 10^{-2}$	$2.43 \pm 0.06$
1000	$(1.0 \pm 0.1) \cdot 10^{-2}$	$2.50 \pm 0.09$
2000	$(5.9 \pm 0.9) \cdot 10^{-3}$	$2.75 \pm 0.14$
4000	$(3.5 \pm 0.7) \cdot 10^{-3}$	$2.82 \pm 0.08$
10000	$(1.9 \pm 0.6) \cdot 10^{-3}$	$3.21 \pm 0.21$