

The Weyl Double Copy for Gravitational Waves

Hadi Godazgar*

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Mühlenberg 1, D-14476 Potsdam, Germany.

Mahdi Godazgar†

School of Mathematical Sciences, Queen Mary University of London, Mile End Road, London E1 4NS, UK.

Ricardo Monteiro‡ and David Peinador Veiga§

*Centre for Research in String Theory, School of Physics and Astronomy,
Queen Mary University of London, E1 4NS, United Kingdom.*

C. N. Pope¶

*George P. & Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy,
Texas A&M University, College Station, TX 77843, USA.***

(Dated: October 7, 2020)

We establish the status of the Weyl double copy relation for radiative solutions of the vacuum Einstein equations. We show that all type N vacuum solutions, which describe the radiation region of isolated gravitational systems with appropriate fall-off for the matter fields, admit a degenerate Maxwell field that squares to give the Weyl tensor. The converse statement also holds, i.e. if there exists a degenerate Maxwell field on a curved background, then the background is type N. This relation defines a scalar that satisfies the wave equation on the background. We show that for non-twisting radiative solutions, the Maxwell field and the scalar also satisfy the Maxwell equation and the wave equation on Minkowski spacetime. Hence, non-twisting solutions have a straightforward double copy interpretation.

The discovery of gravitational waves [1] one hundred years after Einstein formulated his general theory of relativity has led to an exciting new area of gravitational physics with possible important prospects for observational astrophysics; a development that has been anticipated eagerly for half a century [2]. An important theoretical breakthrough in this direction will include an efficient and cost-effective method of generating gravitational wave templates; waveforms computed from the theory to be compared with observed waveforms [3, 4]; see [5–9] for a recent reviews. Amongst the myriad approaches proposed to facilitate the easier and less time-consuming generation of templates is one [10, 11] based on techniques adapted from string theory and supergravity scattering amplitude calculations, in particular the double copy method [12–15], which describes gravitational amplitudes as a kind of inner product of gauge theory amplitudes (hence “double copy”).

While initially found at the level of scattering amplitude relations, the double copy also exists at the level of classical solutions, including beyond perturbation theory for certain classes of spacetimes. One class of solutions for which a double copy relation exists is (multi) Kerr-Schild solutions, which can be thought of as exact perturbative (around Minkowski) gravitational solutions [16, 17]. The correspondence between the double copy relations for scattering amplitudes and for classical solu-

tions has been verified in various works [18–25]; see [26, 27] for earlier ideas in this direction. Of particular interest in the present paper is the Weyl double copy relation that exists for vacuum type D solutions and pp-waves [28–30]. This relation is best expressed in spinor language [31]. In the type D case, it can be shown that the Weyl spinor $\Psi_{ABCD} = (-2\Phi^2)^{-1/4}\Phi_{(AB}\Phi_{CD)}$ with Φ_{AB} a non-degenerate Maxwell spinor and $\Phi^2 \equiv \Phi^{AB}\Phi_{AB}$. Of particular significance is the fact that the Maxwell spinor also solves the Maxwell equation on Minkowski spacetime [30]. Furthermore, $\Phi^{1/2}$ solves the wave equation on Minkowski spacetime. What lies behind these relations is the existence of the well-known hidden symmetry for type D vacuum solutions as expressed by the existence of a Killing 2-spinor [28, 32]. See [33–46] for related works.

In this paper, we extend the curved Weyl double copy relation to all type N vacuum solutions, which describe the radiation region of isolated gravitational systems. In particular, we show that $\Psi_{ABCD} = S^{-1}\Phi_{(AB}\Phi_{CD)}$ with Φ_{AB} a degenerate Maxwell spinor and S some scalar that in particular satisfies the wave equation on the curved background. For non-twisting radiative spacetimes, the Maxwell field and the scalar field also solve the Maxwell equation and the wave equation, respectively, on Minkowski spacetime. This establishes the Weyl double copy in the sense of [30] for this large class of spacetimes. Notice that, while the double copy for scattering ampli-

tudes involves two copies of non-Abelian gauge theory, the first step in that procedure is to consider the double copy of the asymptotic states, which for linearised gauge theory are solutions to the Maxwell equation. The fact that certain exact gravity solutions can be interpreted as a double copy of a Maxwell field means that they should be interpreted as coherent states, an exact extension of the linearised asymptotic states in scattering amplitudes. For twisting spacetimes, the Maxwell field and the scalar depend generically on the metric functions. Hence, they are solutions only on the curved spacetime. However, the standard double copy interpretation applies at the linearised level. This may be indicative of the fact that twisting solutions have an intrinsic non-Abelian nature.

As a necessary step in applying the classical double copy tools to gravitational wave physics, we provide a systematic understanding of the status of the double copy for radiative solutions, beyond the most special example of pp-waves. This study reveals interesting differences with the Weyl double copy for type D solutions. In particular, the construction does not lead to a unique Maxwell field, since there is functional freedom associated to the scalar S . In the cases where the Maxwell field can be thought of as living in Minkowski spacetime, i.e. for non-twisting solutions, it would be interesting to use novel approaches, e.g. [47–49], to relate this new construction to the double copy for scattering amplitudes, as has been done for certain type D solutions.

SPINOR CALCULUS

The homomorphism between the Lorentz group and $SL(2, \mathbb{C})$ can be used to convert spacetime indices μ, ν, \dots into spinor $A, B, \dots = \{1, 2\}$ and conjugate spinor $\dot{A}, \dot{B}, \dots = \{\dot{1}, \dot{2}\}$ indices, using the Van der Waerden matrices $\sigma_{AA'}^\mu$, which are constructed from the identity and Pauli matrices. It is convenient to work in a spinor basis $\{o^A, \iota^A\}$ with $\epsilon_{AB} o^A \iota^B = 1$. In this basis, $\epsilon_{AB} = 2o_{[A} \iota_{B]}$ and can be used to lower indices $\psi_A = \psi^B \epsilon_{BA}$. Similarly, $\psi^A = \epsilon^{AB} \psi_B$. For a vacuum spacetime, the curvature components can be expressed in terms of the totally symmetric Weyl spinor Ψ_{ABCD} , which satisfies the Bianchi identity

$$\nabla^{A\dot{A}} \Psi_{ABCD} = 0. \quad (1)$$

Similarly, a solution of the Maxwell equation can be written in terms of a symmetric 2-spinor Φ_{AB} that solves

$$\nabla^{A\dot{A}} \Phi_{AB} = 0. \quad (2)$$

For type N solutions, choosing a spinor basis adapted to the principal null direction (PND) $\ell^\mu \sim o_A \bar{o}_{\dot{A}}$, the Newman-Penrose (NP) Weyl scalars all vanish except Ψ_4 , and the Weyl spinor takes the simple form

$$\Psi_{ABCD} = \Psi_4 o_A o_B o_C o_D. \quad (3)$$

WEYL DOUBLE COPY

In spinor language, the curved background Weyl double copy relation is

$$\Psi_{ABCD} = \frac{1}{S} \Phi_{(AB} \Phi_{CD)}, \quad (4)$$

for some scalar S and Maxwell spinor Φ_{AB} . Note that Φ_{AB} satisfies the Maxwell equation (2) in the fixed curved background metric, but it is viewed as a test field that does not back-react on the geometry. From (3), it follows that the NP Maxwell scalars all vanish except Φ_2 , and we have $\Phi_{AB} = \Phi_2 o_{(A} o_{B)}$. Thus the type N double copy relation is

$$\Psi_4 = \frac{1}{S} (\Phi_2)^2. \quad (5)$$

The Maxwell 2-spinor is degenerate, which means that the electromagnetic field is null, i.e. the electric and magnetic fields are perpendicular and of equal magnitude. An example of a null electromagnetic field is that of a plane electromagnetic wave in flat spacetime. Now we must consider whether such a relation (4) exists. Expanding out the Bianchi identity (1) by substituting (3) gives two equations:

$$o_A \nabla^{A\dot{A}} \log \Psi_4 + 4 o_A \iota^B \nabla^{A\dot{A}} o_B - \iota_A o^B \nabla^{A\dot{A}} o_B = 0 \quad (6)$$

and $o_A o^B \nabla^{A\dot{A}} o_B = 0$. The second equation is equivalent to the statement that the null congruence generated by the PND is geodesic and shear-free, i.e. $\kappa = \sigma = 0$, which follows from the Goldberg-Sachs theorem [50]. Expanding out the Maxwell equation in a similar fashion gives

$$o_A \nabla^{A\dot{A}} \log \Phi_2 + 2 o_A \iota^B \nabla^{A\dot{A}} o_B - \iota_A o^B \nabla^{A\dot{A}} o_B = 0, \quad (7)$$

as well as the same equation above that is equivalent to $\kappa = \sigma = 0$. Now, substituting $\Psi_4 = (\Phi_2)^2/S$ into (6) and simplifying this using (7) gives

$$o_A \nabla^{A\dot{A}} \log S - \iota_A o^B \nabla^{A\dot{A}} o_B = 0. \quad (8)$$

There is a clear structure in equations (6)–(8), where the coefficient of the middle term is the rank of the respective spinor. Equation (8) translates to

$$\ell \cdot \nabla \log S - \rho = 0, \quad m \cdot \nabla \log S - \tau = 0, \quad (9)$$

where (ℓ, n, m, \bar{m}) form an NP null frame. A simple calculation shows that the integrability condition on equations (9) are satisfied, which means that they are simple integral equations that can always be solved. Thus, we are guaranteed the existence of a scalar S satisfying these equations, which then gives a Maxwell field $\Phi_2 = \sqrt{\Psi_4 S}$. In tensor language, this Maxwell spinor translates to a field strength (called the ‘single copy’) of the form

$$F = \Phi_2 \ell^b \wedge m^b + \bar{\Phi}_2 \ell^b \wedge \bar{m}^b. \quad (10)$$

This establishes the curved Weyl double copy relation for type N vacuum solutions.

Furthermore, it is simple to show using (8) that S solves the wave equation

$$\square S = \nabla_{A\dot{A}} \nabla^{A\dot{A}} S = 2o_{A\dot{A}} \nabla^A \nabla^{\dot{A}} S = 0. \quad (11)$$

The real scalar field in the double copy construction (called the ‘zero-th copy’) is the real part of S .

These results mirror those that exist for type D solutions. In order to investigate whether the Maxwell field and the scalar field also satisfy the equations of motion on Minkowski spacetime, we investigate the different classes of type N solutions in turn.

TYPE N VACUUM SOLUTIONS

Type N vacuum solutions are classified in terms of the optical properties of the congruence generated by the PND. We have that $\kappa = \sigma = 0$; the properties that remain are parametrised by $\rho = -(\Theta + i\omega)$ with Θ denoting the expansion of the congruence and ω denoting its twist. The different cases lead to three distinct classes of solutions:

- Kundt solutions: $\Theta = 0$, which implies that $\omega = 0$ [51].
- Robinson-Trautman solutions: $\Theta \neq 0$, $\omega = 0$.
- Twisting solutions: $\Theta \neq 0$, $\omega \neq 0$.

Choosing a null frame for which ℓ is the PND, so that $\Psi_0 = \Psi_1 = \Psi_2 = \Psi_3 = 0$, we consider each case separately.

Kundt solutions

There are two kinds of type N Kundt solutions, both corresponding to plane-fronted wave solutions [52]. Plane-fronted waves with parallel propagation (pp-waves) are given by the metric

$$ds^2 = -2du(dv + Hdu) + 2dzd\bar{z}, \quad (12)$$

with $H(u, z, \bar{z}) = f(u, z) + \bar{f}(u, \bar{z})$ for general functions f . Choosing

$$\ell = \partial_v, \quad n = \partial_u - H\partial_v, \quad m = \partial_z, \quad (13)$$

one has $\rho = \tau = 0$ and so (9) implies $S = S(u, \bar{z})$, while the Weyl scalar $\Psi_4 = \partial_{\bar{z}}^2 \bar{f}$, so (5) implies that

$$\Phi_2 = \sqrt{\partial_{\bar{z}}^2 \bar{f} S(u, \bar{z})}. \quad (14)$$

The other class of plane-fronted waves is given by

$$ds^2 = -2du(dv + Wdz + \bar{W}d\bar{z} + Hdu) + 2dzd\bar{z}, \quad (15)$$

with $W(v, z, \bar{z}) = -2v(z + \bar{z})^{-1}$ and

$$H(u, v, z, \bar{z}) = [f(u, z) + \bar{f}(u, \bar{z})](z + \bar{z}) - \frac{v^2}{(z + \bar{z})^2},$$

again $f(u, z)$ is arbitrary. Choosing

$$\ell = \partial_v, \quad n = \partial_u - (H + W\bar{W})\partial_v + \bar{W}\partial_z + W\partial_{\bar{z}}, \quad m = \partial_z,$$

one has $\rho = 0$, $\tau = 2\beta = -(z + \bar{z})^{-1}$, so (9) gives $S = \zeta(u, \bar{z})/(z + \bar{z})$. The Weyl scalar $\Psi_4 = (z + \bar{z})\partial_{\bar{z}}^2 \bar{f}$, so (5) implies that

$$\Phi_2 = \sqrt{\partial_{\bar{z}}^2 \bar{f} \zeta(u, \bar{z})}. \quad (16)$$

Given that the only non-zero components of $F_{\mu\nu}$ are for $\mu\nu = [uz]$ and $[u\bar{z}]$, the simple form of the relevant components of $g^{\mu\nu}$ and the fact that $g = 1$ give

$$\begin{aligned} \nabla_\nu F^{\mu\nu} &= \frac{1}{\sqrt{|g|}} \partial_\nu \left(\sqrt{|g|} g^{\mu\rho} g^{\nu\sigma} F_{\rho\sigma} \right) \\ &= \partial_\nu (\eta^{\mu\rho} \eta^{\nu\sigma} F_{\rho\sigma}) = 0. \end{aligned} \quad (17)$$

On the other hand, S does not depend on $f(u, z)$ or $\bar{f}(u, \bar{z})$, meaning that it must solve the wave equation on any member of the family. In particular, it solves the wave equation on Minkowski spacetime. This implies that the Maxwell and the scalar fields also satisfy their equations on Minkowski spacetime, establishing the Weyl double copy for type N Kundt solutions.

Robinson-Trautman solutions

Type N Robinson-Trautman solutions take the form [53]

$$ds^2 = -H du^2 - 2du dr + \frac{2r^2}{P^2} dz d\bar{z}, \quad (18)$$

with $H(u, r, z, \bar{z}) = k - 2r \partial_u \log P$ (where $k = 0, \pm 1$) and $2P^2 \partial_z \partial_{\bar{z}} \log P(u, z, \bar{z}) = k$. Choosing

$$\ell = \partial_r, \quad n = \partial_u - \frac{1}{2} H \partial_r, \quad m = -\frac{P}{r} \partial_z, \quad (19)$$

one has $\rho = -r^{-1}$, $\tau = 0$, so (9) gives $S = -\zeta(u, \bar{z})/r$. Now $\Psi_4 = -\frac{P^2}{r} \partial_u \left(\frac{\partial_{\bar{z}}^2 P}{P} \right)$, so (5) determines that

$$\Phi_2 = \frac{P}{r} \sqrt{\zeta(u, \bar{z}) \partial_u (\partial_{\bar{z}}^2 P/P)}. \quad (20)$$

As an example, consider Robinson-Trautman solutions with $k = 0$ in (18). Writing $P = e^W$ we have $\partial_z \partial_{\bar{z}} W = 0$ and hence $W = w(u, z) + \bar{w}(u, \bar{z})$, implying that $\Psi_4 = -P^2/r \partial_u [\partial_{\bar{z}}^2 \bar{w}(u, \bar{z}) + (\partial_{\bar{z}} \bar{w}(u, \bar{z}))^2]$. We can obtain type N solutions of the Maxwell equation in the Robinson-Trautman background by taking

$$A = \gamma(u, z, \bar{z}) du, \quad (21)$$

where $\partial_z \partial_{\bar{z}} \gamma = 0$ and hence $\gamma = h(u, z) + \bar{h}(u, \bar{z})$. Thus from (10) we have $\Phi_2 = -P/r \partial_{\bar{z}} \bar{h}(u, \bar{z})$. Plugging into (5) we have

$$\partial_u [\partial_{\bar{z}}^2 \bar{w}(u, \bar{z}) + (\partial_{\bar{z}} \bar{w}(u, \bar{z}))^2] = -\frac{1}{rS} (\partial_{\bar{z}} \bar{h}(u, \bar{z}))^2, \quad (22)$$

and so indeed we have that $S = -\zeta/r$, where ζ is a function only of u and \bar{z} , as required in the general result stated above.

As with Kundt solutions, the only non-zero components of $F_{\mu\nu}$ are for $\mu\nu = [uz]$ and $[u\bar{z}]$. As before, using the fact that $\sqrt{|g|} = r^2/P^2$ and the relevant components of $g^{\mu\nu}$, it can be shown that (17) holds. Once again, S is independent of P and solves the wave equation on any member of the family (18), including Minkowski. Hence, both $F_{\mu\nu}$ and S satisfy their equations also on the flat background, establishing the Weyl double copy for Robinson-Trautman solutions.

Twisting solutions

Type N solutions with non-vanishing twist are more complicated, with only one explicit solution known [54]. The general metric is given by [55]

$$ds^2 = -2(du + Ldz + \bar{L}d\bar{z}) \left[dr + Wdz + \bar{W}d\bar{z} + H(du + Ldz + \bar{L}d\bar{z}) \right] + \frac{2}{P^2 |\rho|^2} dzd\bar{z}, \quad (23)$$

$$\begin{aligned} \rho^{-1} &= -(r + i\Sigma), \quad 2i\Sigma(u, z, \bar{z}) = P^2(\bar{\partial}L - \partial\bar{L}), \\ W(u, r, z, \bar{z}) &= \rho^{-1} \partial_u L + i\partial\Sigma, \quad \partial = \partial_z - L\partial_u, \\ H(u, r, z, \bar{z}) &= \frac{1}{2}K - r\partial_u \log P, \end{aligned}$$

with $K = 2P^2 \Re [\partial(\bar{\partial} \log P - \partial_u \bar{L})]$. There exists a residual gauge freedom to choose $P = 1$, but we shall not yet impose this choice. The solution is determined by the complex scalar L , which satisfies

$$\Sigma K + P^2 \Re [\partial\bar{\partial}\Sigma - 2\partial_u \bar{L}\partial\Sigma - \Sigma\partial_u \partial\bar{L}] = 0, \quad \partial I = 0,$$

and $\partial_u I \neq 0$, with $I = \bar{\partial}(\bar{\partial} \log P - \partial_u \bar{L}) + (\bar{\partial} \log P - \partial_u \bar{L})^2$. Choosing

$$\ell = \partial_r, \quad n = \partial_u - H\partial_r, \quad m = -P\bar{\rho}(\partial - W\partial_r), \quad (24)$$

ρ is as defined above, while $\tau = 0$. Equation (9) then implies that $S = \rho \chi(u, z, \bar{z})$, with χ satisfying

$$\partial\chi - \partial_u L \chi = 0. \quad (25)$$

Defining new coordinates $(v, w) = (I, z)$, the above equation can be solved using the method of characteristics ($I = \text{constant}$ correspond to the characteristics)

$$\chi(v, w) = \zeta(I) e^{\int \left[\left(\frac{\partial I(u, z)}{\partial u} \right)_{(v, w')} \times \frac{\partial L(v, w')}{\partial v} \right] dw'}, \quad (26)$$

with $\zeta(I)$ arbitrary. The Weyl scalar $\Psi_4 = \rho P^2 \partial_u I$, and so (5) implies

$$\Phi_2 = \rho P \sqrt{\partial_u I \chi(u, z, \bar{z})}. \quad (27)$$

Only one twisting type N solution, found by Hauser [54], is known explicitly. The metric functions are given by

$$P = (z + \bar{z})^{3/2} f(t), \quad t \equiv \frac{u}{(z + \bar{z})^2}, \quad L = 2i(z + \bar{z}),$$

where f satisfies $16(1 + t^2)f''(t) + 3f(t) = 0$, which is a hypergeometric equation, and I turns out to be given by

$$I = \frac{3}{2[(z + \bar{z})^2 - iu]}. \quad (28)$$

The solution to (9) is

$$S = \rho \zeta(I), \quad (29)$$

where $\zeta(I)$ is arbitrary. As expected, this is consistent with the general result (26). The Weyl scalar is $\Psi_4 = (2i/3) \rho P^2 I^2$, implying that

$$\Phi_2 = \rho P I \sqrt{\frac{2i\zeta(I)}{3}}. \quad (30)$$

As a further remark about the twisting type N solutions, we note that if the gauge freedom to set $P = 1$ is employed, the metric is specified purely in terms of the function $L(u, z, \bar{z})$, and the type N and Ricci flat conditions may be succinctly condensed down to just

$$\partial I = 0, \quad \Im(\bar{\partial}\bar{\partial}\partial L) = 0, \quad \text{where } I = -\partial_u \bar{\partial} \bar{L}. \quad (31)$$

The Weyl curvature is given by $\Psi_4 = \rho \partial_u I$.

In contrast to non-twisting solutions, the second equality in (17) does not hold for twisting solutions. Therefore, while there is a curved Weyl double copy relation, in this case it does not translate to a relation where the Maxwell field and the scalar can be thought of as Minkowski fields, unless we consider all the fields (gravity, Maxwell and scalar) at the linearised level.

NON-UNIQUENESS

In all the cases above, neither the Maxwell field nor the scalar field are uniquely determined. They are fixed only up to an arbitrary function of some of the coordinates, which we are free to choose. This contrasts with the Weyl double copy for vacuum type D solutions, for which, in a spinor basis adapted to the principal null directions, we have $S^3 \propto (\Phi_2)^{3/2} \propto \Psi_4$, where the proportionality is up to complex parameters [30]; hence the Maxwell and scalar fields are functionally fixed. This feature is related

to the fact that vacuum type D spacetimes are fully determined up to a few parameters, whereas vacuum type N spacetimes (of any class, as seen above) have functional freedom. By analogy, there is additional freedom in the Maxwell and scalar fields in the curved background.

In considering a special choice, we may ask whether it is possible to choose Φ_2 and S to be given by specific powers of Ψ_4 , as in the type D case, i.e. there exists some constant a such that $\Phi_2 \propto (\Psi_4)^a$ and $S \propto (\Psi_4)^{2a-1}$. The functional dependence of the results above implies that this possibility holds only for Kundt solutions. For pp-waves, the power is actually undetermined, i.e. the relation above holds for any a . A simple choice is $a = 1/2$, where S is constant, and in fact this choice implies that Maxwell plane waves double copy to gravitational plane waves (Φ_2 and Ψ_4 are functions of u only). For the other plane-fronted Kundt solutions, such a relation is possible for $a = 0$, in which case $S \propto (\Psi_4)^{-1}$. Analogously simple choices for the other type N classes are: $S \propto 1/r$ for Robinson-Trautman solutions and $S \propto \rho$ for twisting solutions.

Interestingly, pp-waves are the only type N solutions admitting a Killing 2-spinor [29], another feature that they share with type D solutions.

Acknowledgements HG would like to thank Queen Mary University of London for hospitality during the course of this work. HG is supported by the ERC Advanced Grant Exceptional Quantum Gravity (Grant No. 740209). MG and RM are supported by Royal Society University Research Fellowships. CNP is partially supported by DOE grant DE-FG02-13ER42020. DPV is supported by a Royal Society studentship grant.

* hadi.godazgar@aei.mpg.de

† m.godazgar@qmul.ac.uk

‡ ricardo.monteiro@qmul.ac.uk

§ d.peinadorveiga@qmul.ac.uk

¶ pope@physics.tamu.edu

** Also at Centre for Mathematical Sciences, Cambridge University, Wilberforce Road, Cambridge, CB3 0WA, UK.

- [1] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [2] W. H. Press and K. S. Thorne, *Astronomy, Gravitational-Wave* *Ann. Rev. Astron. Astrophys.* **10**, 335 (1972).
- [3] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, *Phys. Rev. D* **59**, 084006 (1999), arXiv:gr-qc/9811091.
- [4] F. Pretorius, Evolution of binary black hole spacetimes, *Phys. Rev. Lett.* **95**, 121101 (2005), arXiv:gr-qc/0507014.
- [5] M. Shibata and K. Taniguchi, Coalescence of Black Hole-Neutron Star Binaries, *Living Rev. Rel.* **14**, 6 (2011).
- [6] N. T. Bishop and L. Rezzolla, Extraction of Gravitational Waves in Numerical Relativity, *Living Rev. Rel.* **19**, 2 (2016), arXiv:1606.02532 [gr-qc].
- [7] L. Blanchet, Gravitational Radiation from Post-Newtonian Sources and Inspiralling Compact Binaries, *Living Rev. Rel.* **17**, 2 (2014), arXiv:1310.1528 [gr-qc].
- [8] R. A. Porto, The effective field theorist's approach to gravitational dynamics, *Phys. Rept.* **633**, 1 (2016), arXiv:1601.04914 [hep-th].
- [9] L. Barack and A. Pound, Self-force and radiation reaction in general relativity, *Rept. Prog. Phys.* **82**, 016904 (2019), arXiv:1805.10385 [gr-qc].
- [10] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order, *Phys. Rev. Lett.* **122**, 201603 (2019), arXiv:1901.04424 [hep-th].
- [11] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Black Hole Binary Dynamics from the Double Copy and Effective Theory, *JHEP* **10**, 206, arXiv:1908.01493 [hep-th].
- [12] H. Kawai, D. Lewellen, and S. Tye, A Relation Between Tree Amplitudes of Closed and Open Strings, *Nucl. Phys. B* **269**, 1 (1986).
- [13] Z. Bern, J. Carrasco, and H. Johansson, New Relations for Gauge-Theory Amplitudes, *Phys. Rev. D* **78**, 085011 (2008), arXiv:0805.3993 [hep-ph].
- [14] Z. Bern, J. J. M. Carrasco, and H. Johansson, Perturbative Quantum Gravity as a Double Copy of Gauge Theory, *Phys. Rev. Lett.* **105**, 061602 (2010), arXiv:1004.0476 [hep-th].
- [15] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson, and R. Roiban, The Duality Between Color and Kinematics and its Applications, (2019), arXiv:1909.01358 [hep-th].
- [16] R. Monteiro, D. O'Connell, and C. D. White, Black holes and the double copy, *JHEP* **12**, 056, arXiv:1410.0239 [hep-th].
- [17] A. Luna, R. Monteiro, D. O'Connell, and C. D. White, The classical double copy for Taub-NUT spacetime, *Phys. Lett. B* **750**, 272 (2015), arXiv:1507.01869 [hep-th].
- [18] A. Luna, R. Monteiro, I. Nicholson, D. O'Connell, and C. D. White, The double copy: Bremsstrahlung and accelerating black holes, *JHEP* **06**, 023, arXiv:1603.05737 [hep-th].
- [19] W. D. Goldberger and A. K. Ridgway, Radiation and the classical double copy for color charges, *Phys. Rev. D* **95**, 125010 (2017), arXiv:1611.03493 [hep-th].
- [20] A. Luna, R. Monteiro, I. Nicholson, A. Ochirov, D. O'Connell, N. Westerberg, and C. D. White, Perturbative spacetimes from Yang-Mills theory, *JHEP* **04**, 069, arXiv:1611.07508 [hep-th].
- [21] N. Arkani-Hamed, Y.-t. Huang, and D. O'Connell, Kerr black holes as elementary particles, *JHEP* **01**, 046, arXiv:1906.10100 [hep-th].
- [22] Y.-T. Huang, U. Kol, and D. O'Connell, Double copy of electric-magnetic duality, *Phys. Rev. D* **102**, 046005 (2020), arXiv:1911.06318 [hep-th].

- [23] K. Kim, K. Lee, R. Monteiro, I. Nicholson, and D. Peinador Veiga, The Classical Double Copy of a Point Charge, *JHEP* **02**, 046, arXiv:1912.02177 [hep-th].
- [24] A. Luna, S. Nagy, and C. White, The convolutional double copy: a case study with a point, (2020), arXiv:2004.11254 [hep-th].
- [25] A. Cristofoli, Gravitational shock waves and scattering amplitudes, (2020), arXiv:2006.08283 [hep-th].
- [26] R. Saotome and R. Akhoury, Relationship Between Gravity and Gauge Scattering in the High Energy Limit, *JHEP* **01**, 123, arXiv:1210.8111 [hep-th].
- [27] D. Neill and I. Z. Rothstein, Classical Space-Times from the S Matrix, *Nucl. Phys. B* **877**, 177 (2013), arXiv:1304.7263 [hep-th].
- [28] M. Walker and R. Penrose, On quadratic first integrals of the geodesic equations for type [22] spacetimes, *Commun. Math. Phys.* **18**, 265 (1970).
- [29] W. Dietz and R. Rüdiger, Space-times admitting Killing-Yano tensors. I, *Proc. Roy. Soc. Lond. A* **375**, 361 (1981).
- [30] A. Luna, R. Monteiro, I. Nicholson, and D. O’Connell, Type D Spacetimes and the Weyl Double Copy, *Class. Quant. Grav.* **36**, 065003 (2019), arXiv:1810.08183 [hep-th].
- [31] We summarise some important equations regarding spinor calculus in curved spacetime below. For comprehensive introductions, see [56, 57].
- [32] B. Carter, Global structure of the kerr family of gravitational fields, *Phys. Rev.* **174**, 1559 (1968).
- [33] A. Ilderton, Screw-symmetric gravitational waves: a double copy of the vortex, *Phys. Lett. B* **782**, 22 (2018), arXiv:1804.07290 [gr-qc].
- [34] K. Lee, Kerr-Schild Double Field Theory and Classical Double Copy, *JHEP* **10**, 027, arXiv:1807.08443 [hep-th].
- [35] D. S. Berman, E. Chacón, A. Luna, and C. D. White, The self-dual classical double copy, and the Eguchi-Hanson instanton, *JHEP* **01**, 107, arXiv:1809.04063 [hep-th].
- [36] K. Andrzejewski and S. Prencel, From polarized gravitational waves to analytically solvable electromagnetic beams, *Phys. Rev. D* **100**, 045006 (2019), arXiv:1901.05255 [hep-th].
- [37] S. Sabharwal and J. W. Dalhuisen, Anti-Self-Dual Spacetimes, Gravitational Instantons and Knotted Zeros of the Weyl Tensor, *JHEP* **07**, 004, arXiv:1904.06030 [hep-th].
- [38] H. Godazgar, M. Godazgar, and C. Pope, Taub-NUT from the Dirac monopole, *Phys. Lett. B* **798**, 134938 (2019), arXiv:1908.05962 [hep-th].
- [39] I. Bah, R. Dempsey, and P. Weck, Kerr-Schild Double Copy and Complex Worldlines, *JHEP* **02**, 180, arXiv:1910.04197 [hep-th].
- [40] R. Alawadhi, D. S. Berman, B. Spence, and D. Peinador Veiga, S-duality and the double copy, *JHEP* **03**, 059, arXiv:1911.06797 [hep-th].
- [41] C. Keeler, T. Manton, and N. Monga, From Navier-Stokes to Maxwell via Einstein, *JHEP* **08**, 147, arXiv:2005.04242 [hep-th].
- [42] N. Bahjat-Abbas, R. Stark-Muchão, and C. D. White, Monopoles, shockwaves and the classical double copy, *JHEP* **04**, 102, arXiv:2001.09918 [hep-th].
- [43] G. Elor, K. Farnsworth, M. L. Graesser, and G. Herzog, The Newman-Penrose Map and the Classical Double Copy, (2020), arXiv:2006.08630 [hep-th].
- [44] R. Alawadhi, D. S. Berman, and B. Spence, Weyl doubling, *JHEP* **09**, 127, arXiv:2007.03264 [hep-th].
- [45] L. Alfonsi, C. D. White, and S. Wikeley, Topology and Wilson lines: global aspects of the double copy, *JHEP* **07**, 091, arXiv:2004.07181 [hep-th].
- [46] T. Adamo and A. Ilderton, Classical and quantum double copy of back-reaction, *JHEP* **09**, 200, arXiv:2005.05807 [hep-th].
- [47] T. Adamo, E. Casali, L. Mason, and S. Nekovar, Scattering on plane waves and the double copy, *Class. Quant. Grav.* **35**, 015004 (2018), arXiv:1706.08925 [hep-th].
- [48] A. Anastasiou, L. Borsten, M. Duff, S. Nagy, and M. Zoccali, Gravity as Gauge Theory Squared: A Ghost Story, *Phys. Rev. Lett.* **121**, 211601 (2018), arXiv:1807.02486 [hep-th].
- [49] D. A. Kosower, B. Maybee, and D. O’Connell, Amplitudes, Observables, and Classical Scattering, *JHEP* **02**, 137, arXiv:1811.10950 [hep-th].
- [50] J. N. Goldberg and R. K. Sachs, A theorem on Petrov types, *Acta Phys. Pol.* **22 (supplement)** (1962).
- [51] Substituting $\Theta = \sigma = R_{\mu\nu} = 0$ into the Raychaudhuri equation $\ell \cdot \nabla \Theta - \omega^2 + \Theta^2 + \sigma\bar{\sigma} + \frac{1}{2}R_{\mu\nu}\ell^\mu\ell^\nu = 0$ gives $\omega = 0$.
- [52] See Theorem 31.2 of [58].
- [53] See Theorem 28.1 and Section 28.1.2 of [58].
- [54] I. Hauser, Type-N gravitational field with twist, *Phys. Rev. Lett.* **33**, 1112 (1974).
- [55] See Chapter 29 of Ref. [58].
- [56] R. Penrose and W. Rindler, *Spinors and space-time. Vol. 1: Two-spinor Calculus and Relativistic Fields* (Cambridge Univ. Press, 1984).
- [57] J. Stewart, *Advanced General Relativity*, Cambridge Monographs on Mathematical Physics (CUP, 1991).
- [58] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers, and E. Herlt, *Exact solutions of Einstein’s field equations*, Cambridge Monographs on Mathematical Physics (Cambridge Univ. Press, Cambridge, 2003).