

# On the Decidability of Termination for Polynomial Loops<sup>\*</sup>

Florian Frohn<sup>1</sup>, Marcel Hark<sup>2</sup>, and Jürgen Giesl<sup>2</sup>

<sup>1</sup> Max Planck Institute for Informatics and Saarland Informatics Campus, Saarbrücken, Germany

<sup>2</sup> LuFG Informatik 2, RWTH Aachen University, Germany

**Abstract.** We consider the termination problem for triangular weakly non-linear loops (*twn*-loops) over some ring  $\mathcal{S}$  like  $\mathbb{Z}$ ,  $\mathbb{Q}$ , or  $\mathbb{R}$ . Essentially, the guard of such a loop is an arbitrary Boolean formula over (possibly non-linear) polynomial inequations, and the body is a single assignment  $\begin{bmatrix} x_1 \\ \dots \\ x_d \end{bmatrix} \leftarrow \begin{bmatrix} c_1 \cdot x_1 + p_1 \\ \dots \\ c_d \cdot x_d + p_d \end{bmatrix}$  where each  $x_i$  is a variable,  $c_i \in \mathcal{S}$ , and each  $p_i$  is a (possibly non-linear) polynomial over  $\mathcal{S}$  and the variables  $x_{i+1}, \dots, x_d$ . We present a reduction from the question of termination to the existential fragment of the first-order theory of  $\mathcal{S}$  and  $\mathbb{R}$ . For loops over  $\mathbb{R}$ , our reduction entails decidability of termination. For loops over  $\mathbb{Z}$  and  $\mathbb{Q}$ , it proves semi-decidability of non-termination.

Furthermore, we present a transformation to convert certain non-*twn*-loops into *twn*-form. Then the original loop terminates iff the transformed loop terminates over a specific subset of  $\mathbb{R}$ , which can also be checked via our reduction. This transformation also allows us to prove *tight* complexity bounds for the termination problem for two important classes of loops which can *always* be transformed into *twn*-loops.

## 1 Introduction

Let  $\mathbb{R}_{\mathbb{A}}$  denote the real algebraic numbers. We consider loops of the form

$$\mathbf{while} \ \varphi \ \mathbf{do} \ \vec{x} \leftarrow \vec{a}. \quad (1)$$

Here,  $\vec{x}$  is a vector<sup>3</sup> of  $d \geq 1$  pairwise different variables that range over a ring  $\mathbb{Z} \leq \mathcal{S} \leq \mathbb{R}_{\mathbb{A}}$ , where  $\leq$  denotes the subring relation. Moreover,  $\vec{a} \in (\mathcal{S}[\vec{x}])^d$  where  $\mathcal{S}[\vec{x}]$  is the set of all polynomials over  $\vec{x}$  with coefficients from  $\mathcal{S}$ . The condition  $\varphi$  is an arbitrary propositional formula over the atoms  $\{p \triangleright 0 \mid p \in \mathcal{S}[\vec{x}], \triangleright \in \{\geq, >\}\}$ .<sup>4</sup>

<sup>\*</sup> funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 389792660 as part of [TRR 248](#), by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - 235950644 (Project GI 274/6-2), and by the DFG Research Training Group 2236 UnRAVeL

<sup>3</sup> We use row- and column-vectors interchangeably to improve readability.

<sup>4</sup> Note that negation is syntactic sugar in our setting, as, e.g.,  $\neg(p > 0)$  is equivalent to  $-p \geq 0$ . So w.l.o.g.  $\varphi$  is built from atoms,  $\wedge$ , and  $\vee$ .

We require  $\mathcal{S} \leq \mathbb{R}_\mathbb{A}$  instead of  $\mathcal{S} \leq \mathbb{R}$ , as it is unclear how to represent transcendental numbers on computers. However, in [Sect. 5](#) we will see that the loops considered in this paper terminate over  $\mathbb{R}$  iff they terminate over  $\mathbb{R}_\mathbb{A}$ . Thus, our results immediately carry over to loops where the variables range over  $\mathbb{R}$ . Hence, we sometimes also consider loops over  $\mathcal{S} = \mathbb{R}$ . However, even then we restrict ourselves to loops [\(1\)](#) where all constants in  $\varphi$  and  $\vec{a}$  are algebraic.

We often represent a loop [\(1\)](#) by the tuple  $(\varphi, \vec{a})$  of the *loop condition*  $\varphi$  and the *update*  $\vec{a} = (a_1, \dots, a_d)$ . Unless stated otherwise,  $(\varphi, \vec{a})$  is always a loop on  $\mathcal{S}^d$  using the variables  $\vec{x} = (x_1, \dots, x_d)$  where  $\mathbb{Z} \leq \mathcal{S} \leq \mathbb{R}_\mathbb{A}$ . A *linear-update loop* has the form  $(\varphi, A \cdot \vec{x} + \vec{b})$  and it has *real spectrum* if  $A$  has real eigenvalues only. A *linear loop* is a linear-update loop where  $\varphi$  is linear (i.e., its atoms are only linear<sup>5</sup> inequations). A *conjunctive loop* is a loop  $(\varphi, \vec{a})$  where  $\varphi$  does not contain disjunctions.

There exist several decidability results for the termination of linear loops [\[6, 8, 15, 23, 33, 36, 40, 52\]](#), but there are only very few results on the decidability of termination for certain forms of non-linear loops [\[34, 35, 37, 54\]](#). Moreover, all of these previous works only consider conjunctive loops besides [\[37\]](#) which only allows for loop conditions defining compact sets. In this paper, we regard (linear and non-linear) loops with arbitrary conditions, i.e., they may also contain disjunctions and define non-compact sets. Furthermore, we study the decidability of termination for non-linear loops over  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}_\mathbb{A}$ , and  $\mathbb{R}$ , whereas the existing decidability results for non-linear loops are restricted to loops over  $\mathbb{R}$ . So we identify new sub-classes of loops of the form [\(1\)](#) where (non-)termination is (semi-)decidable. Moreover, we also investigate the complexity of the termination problem.

*Contributions:* We study a sub-class of loops of the form [\(1\)](#) (so-called *twn-loops* ([Sect. 2](#))), and present an (incomplete) transformation  $Tr$  from non-*twn-loops* to *twn-loops* ([Sect. 3](#)). Then we show that termination of *twn-loops* over  $\mathbb{R}_\mathbb{A}$  and  $\mathbb{R}$  is decidable and that non-termination over  $\mathbb{Z}$  and  $\mathbb{Q}$  is semi-decidable ([Sect. 4](#) and [5](#)). For those classes of non-*twn-loops* where our transformation  $Tr$  is complete, we obtain analogous decidability results. For all other loops of the form [\(1\)](#), our (semi-)decision procedures still apply if  $Tr$  is applicable.

Finally, we prove **Co-NP-completeness** of termination of linear loops over  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}_\mathbb{A}$ , and  $\mathbb{R}$  with real spectrum and  **$\forall\mathbb{R}$ -completeness** of termination of linear-update loops with real spectrum over  $\mathbb{R}_\mathbb{A}$  and  $\mathbb{R}$  ([Sect. 6](#)).

The appendix contains all missing proofs..

## 2 Preliminaries

For any entity  $s$ ,  $s[x/t]$  is the entity that results from  $s$  by replacing all free occurrences of  $x$  by  $t$ . Similarly, if  $\vec{x} = (x_1, \dots, x_d)$  and  $\vec{t} = (t_1, \dots, t_d)$ , then

<sup>5</sup> In this paper “linear” refers to “linear polynomials” and thus includes *affine* functions.

$s[\vec{x}/\vec{t}]$  results from  $s$  by replacing all free occurrences of  $x_i$  by  $t_i$ , for each  $1 \leq i \leq d$ .

Any vector of polynomials  $\vec{a} \in (\mathcal{S}[\vec{x}])^d$  can also be regarded as a function  $\vec{a} : (\mathcal{S}[\vec{x}])^d \rightarrow (\mathcal{S}[\vec{x}])^d$ , where for any  $\vec{p} \in (\mathcal{S}[\vec{x}])^d$ ,  $\vec{a}(\vec{p}) = \vec{a}[\vec{x}/\vec{p}]$  results from *applying* the polynomials  $\vec{a}$  to the polynomials  $\vec{p}$ . In a similar way, we can also apply a formula to polynomials  $\vec{p} \in (\mathcal{S}[\vec{x}])^d$ . To this end, we define  $\psi(\vec{p}) = \psi[\vec{x}/\vec{p}]$  for first-order formulas  $\psi$  with free variables  $\vec{x}$ . As usual, function application associates to the left, i.e.,  $\vec{a}(\vec{b})(\vec{p})$  stands for  $(\vec{a}(\vec{b}))(\vec{p})$ . However, since applying polynomials only means that one instantiates variables, we obviously have  $(\vec{a}(\vec{b}))(\vec{p}) = \vec{a}(\vec{b}(\vec{p}))$ .

**Def. 1** formalizes the intuitive notion of termination for a loop  $(\varphi, \vec{a})$ .

**Definition 1 (Termination).** *The loop  $(\varphi, \vec{a})$  is non-terminating (over  $\mathcal{S}$ ) if*

$$\exists \vec{c} \in \mathcal{S}^d. \forall n \in \mathbb{N}. \varphi(\vec{a}^n(\vec{c})).$$

*Then  $\vec{c}$  is a witness for non-termination. Otherwise,  $(\varphi, \vec{a})$  terminates (over  $\mathcal{S}$ ).*

Here,  $\vec{a}^n$  denotes the  $n$ -fold application of  $\vec{a}$ , i.e.,  $\vec{a}^0(\vec{c}) = \vec{c}$  and  $\vec{a}^{n+1}(\vec{c}) = \vec{a}(\vec{a}^n(\vec{c}))$ . Termination (which is sometimes also called *universal* termination) is not to be confused with the *halting problem*, where one is interested in termination w.r.t. a *given* input. In contrast, **Def. 1** considers termination w.r.t. *all* inputs.

For any entity  $s$ , let  $\mathcal{V}(s)$  be the set of all free variables that occur in  $s$ . Given an assignment  $\vec{x} \leftarrow \vec{a}$ , the relation  $\succ_{\vec{a}} \in \mathcal{V}(\vec{a}) \times \mathcal{V}(\vec{a})$  is the transitive closure of  $\{(x_i, x_j) \mid i, j \in \{1, \dots, d\}, i \neq j, x_j \in \mathcal{V}(a_i)\}$ . We call  $(\varphi, \vec{a})$  *triangular* if  $\succ_{\vec{a}}$  is well founded. So the restriction to triangular loops prohibits “cyclic dependencies” of variables (e.g., where the new values of  $x_1$  and  $x_2$  both depend on the old values of  $x_1$  and  $x_2$ ). For example, a loop with the body  $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leftarrow \begin{bmatrix} x_1 + x_2^2 \\ x_2 - 1 \end{bmatrix}$  is triangular since  $\succ = \{(x_1, x_2)\}$  is well founded, whereas a loop with the body  $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leftarrow \begin{bmatrix} x_1 + x_2^2 \\ x_1 - 1 \end{bmatrix}$  is not triangular. Triangularity is used to compute a *closed form* for the  $n$ -fold application of the loop update  $\vec{a}$ , i.e., a vector  $\vec{q}$  of  $d$  expressions over the variables  $\vec{x}$  and  $n$  such that  $\vec{q} = \vec{a}^n$ . From a practical point of view, the restriction to triangular loops seems quite natural. For example, in [17], 1511 polynomial loops were extracted from the *Termination Problems Data Base* [53], the benchmark collection which is used at the annual *Termination and Complexity Competition* [20], and only 26 of them were non-triangular.

The loop  $(\varphi, \vec{a})$  is *weakly non-linear* if for no  $i$ ,  $x_i$  occurs in a non-linear monomial of  $a_i$ . So for example, a loop with the body  $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leftarrow \begin{bmatrix} x_1 + x_2^2 \\ x_2 - 1 \end{bmatrix}$  is weakly non-linear, whereas a loop with the body  $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \leftarrow \begin{bmatrix} x_1 \cdot x_2 \\ x_2 - 1 \end{bmatrix}$  is not. Together with triangularity, weak non-linearity ensures that we can always compute closed forms. In particular, weak non-linearity excludes loops like  $(\varphi, x \leftarrow x^2)$  that need exponential space, as the values of some variables grow doubly exponentially.

A *twn*-loop is triangular and weakly non-linear. So in other words, by permuting variables every *twn*-loop can be transformed to the form

$$\begin{bmatrix} x_1 \\ \vdots \\ x_d \end{bmatrix} \leftarrow \begin{bmatrix} c_1 \cdot x_1 + p_1 \\ \vdots \\ c_d \cdot x_d + p_d \end{bmatrix}$$

where  $c_i \in \mathcal{S}$  and  $p_i \in \mathcal{S}[x_{i+1}, \dots, x_d]$ . If  $(\varphi, \vec{a})$  is weakly non-linear and  $x_i$ 's coefficient in  $a_i$  is non-negative for all  $1 \leq i \leq d$ , then  $(\varphi, \vec{a})$  is *non-negative*. A *tnn*-loop is triangular and non-negative (and thus, also weakly non-linear).

Our *tnw*-loops are a special case of *solvable loops* [47].

**Definition 2 (Solvable Loops).** A loop  $(\varphi, \vec{a})$  is solvable if there is a partitioning  $\mathcal{J} = \{J_1, \dots, J_k\}$  of  $\{1, \dots, d\}$  such that for each  $1 \leq i \leq k$  we have

$$\vec{a}_{J_i} = A_i \cdot \vec{x}_{J_i} + \vec{p}_i,$$

where  $\vec{a}_{J_i}$  is the vector of all  $a_j$  with  $j \in J_i$  (and  $\vec{x}_{J_i}$  is defined analogously),  $d_i = |J_i|$ ,  $A_i \in \mathcal{S}^{d_i \times d_i}$ , and  $\vec{p}_i \in (\mathcal{S}[x_{J_{i+1}}, \dots, x_{J_k}])^{d_i}$ . The eigenvalues of a solvable loop are defined as the union of the eigenvalues of all  $A_i$ .

So solvable loops allow for blocks of variables with linear dependencies, and *tnw*-loops correspond to the case that each such block has size 1. While our approach could easily be generalized to solvable loops with real eigenvalues, in Sect. 3 we show that such a generalization does not increase its applicability.

For a ring  $\mathbb{Z} \leq \mathcal{S} \leq \mathbb{R}_{\mathbb{A}}$ , the *existential fragment of the first-order theory of  $\mathcal{S}$*  is the set  $\text{Th}_{\exists}(\mathcal{S})$  of all formulas  $\exists \vec{y} \in \mathcal{S}^k. \psi$ , where  $\psi$  is a propositional formula over the atoms  $\{p \triangleright 0 \mid p \in \mathbb{Q}[\vec{y}, \vec{z}], \triangleright \in \{\geq, >\}\}$  and  $k \in \mathbb{N}$  [43, 50]. Here,  $\vec{y}$  and  $\vec{z}$  are pairwise disjoint vectors of variables (i.e., the variables  $\vec{z}$  are *free*). Moreover,  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  is the set of all formulas  $\exists \vec{y}' \in \mathbb{R}_{\mathbb{A}}^{k'}, \vec{y} \in \mathcal{S}^k. \psi$ , with a propositional formula  $\psi$  over  $\{p \triangleright 0 \mid p \in \mathbb{Q}[\vec{y}', \vec{y}, \vec{z}], \triangleright \in \{\geq, >\}\}$  where  $k', k \in \mathbb{N}$  and the variables  $\vec{y}'$ ,  $\vec{y}$ , and  $\vec{z}$  are pairwise disjoint. As usual, a formula without free variables is *closed*. In the following, we also consider formulas over inequations  $p \triangleright 0$  where  $p$ 's coefficients are from  $\mathbb{R}_{\mathbb{A}}$  to be elements of  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$  (resp.  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ ). The reason is that real algebraic numbers are  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ -definable.

Finally, note that validity of formulas from  $\text{Th}_{\exists}(\mathcal{S})$  or  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  is decidable if  $\mathcal{S} \in \{\mathbb{R}_{\mathbb{A}}, \mathbb{R}\}$  and semi-decidable if  $\mathcal{S} \in \{\mathbb{Z}, \mathbb{Q}\}$  [11, 51]. By undecidability of Hilbert's Tenth Problem, validity is undecidable for  $\mathcal{S} = \mathbb{Z}$ . While validity of full first-order formulas (i.e., also containing universal quantifiers) over  $\mathcal{S} = \mathbb{Q}$  is undecidable [44], it is still open whether validity of formulas from  $\text{Th}_{\exists}(\mathbb{Q})$  or  $\text{Th}_{\exists}(\mathbb{Q}, \mathbb{R}_{\mathbb{A}})$  is decidable. However, validity of *linear* formulas from  $\text{Th}_{\exists}(\mathcal{S})$  or  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  is decidable for all  $\mathcal{S} \in \{\mathbb{Z}, \mathbb{Q}, \mathbb{R}_{\mathbb{A}}, \mathbb{R}\}$ .

### 3 Transformation to Triangular Weakly Non-Linear Form

We first show how to handle loops that are not yet *tnw*. To this end, we introduce a transformation of loops via *polynomial automorphisms* in Sect. 3.1 and show that our transformation preserves (non-)termination (Thm. 10). In Sect. 3.2, we use results from algebraic geometry to show that the question whether a loop can be transformed into *tnw*-form is reducible to validity of  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ -formulas (Thm. 20). Moreover, we show that it is decidable whether a *linear* automorphism can transform a loop into a special case of the *tnw*-form (Thm. 23).

### 3.1 Transforming Loops

Clearly, the *polynomials*  $x_1, \dots, x_d$  are *generators* of the  $\mathcal{S}$ -algebra  $\mathcal{S}[\vec{x}]$ , i.e., every polynomial from  $\mathcal{S}[\vec{x}]$  can be obtained from  $x_1, \dots, x_d$  and the operations of the algebra (i.e., addition and multiplication). So far, we have implicitly chosen a special “representation” of the loop based on the generators  $x_1, \dots, x_d$ .

We now change this representation, i.e., we use a different set of  $d$  polynomials which are also generators of  $\mathcal{S}[\vec{x}]$ . Then the loop has to be modified accordingly in order to adapt it to this new representation. This modification does not affect the loop’s termination behavior, but it may transform a non-*twn*-loop into *twn*-form.

The desired change of representation is described by  $\mathcal{S}$ -*automorphisms* of  $\mathcal{S}[\vec{x}]$ . As usual, an  $\mathcal{S}$ -*endomorphism* of  $\mathcal{S}[\vec{x}]$  is a mapping  $\eta : \mathcal{S}[\vec{x}] \rightarrow \mathcal{S}[\vec{x}]$  which is  $\mathcal{S}$ -linear and multiplicative.<sup>6</sup> We denote the ring of  $\mathcal{S}$ -endomorphisms of  $\mathcal{S}[\vec{x}]$  by  $\text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  (where the operations on this ring are pointwise addition and function composition  $\circ$ ). The group of  $\mathcal{S}$ -automorphisms of  $\mathcal{S}[\vec{x}]$  is  $\text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ ’s group of units, and we denote it by  $\text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ . So an  $\mathcal{S}$ -automorphism of  $\mathcal{S}[\vec{x}]$  is an  $\eta \in \text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  that is *invertible*. Thus, there exists an  $\eta^{-1} \in \text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  such that  $\eta \circ \eta^{-1} = \eta^{-1} \circ \eta = \text{id}_{\mathcal{S}[\vec{x}]}$ , where  $\text{id}_{\mathcal{S}[\vec{x}]}$  is the identity function on  $\mathcal{S}[\vec{x}]$ .

*Example 3 (Automorphism).* Let  $\eta \in \text{End}_{\mathcal{S}}(\mathcal{S}[x_1, x_2])$  with  $\eta(x_1) = x_2$ ,  $\eta(x_2) = x_1 - x_2^2$ . Then  $\eta \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[x_1, x_2])$ , where  $\eta^{-1}(x_1) = x_1^2 + x_2$  and  $\eta^{-1}(x_2) = x_1$ .

As  $\mathcal{S}[\vec{x}]$  is free on the generators  $\vec{x}$ , an endomorphism  $\eta \in \text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  is uniquely determined by the images of the variables, i.e., by  $\eta(x_1), \dots, \eta(x_d)$ . Hence, we have a one-to-one correspondence between elements of  $(\mathcal{S}[\vec{x}])^d$  and  $\text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ . In particular, every tuple  $\vec{a} = (a_1, \dots, a_d) \in (\mathcal{S}[\vec{x}])^d$  corresponds to the unique endomorphism  $\tilde{a} \in \text{End}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  with  $\tilde{a}(x_i) = a_i$  for all  $1 \leq i \leq d$ . So for any  $p \in \mathcal{S}[\vec{x}]$  we have  $\tilde{a}(p) = p(\vec{a})$ . Thus, the update of a loop induces an endomorphism which operates on polynomials.

*Example 4 (Updates as Endomorphisms).* Consider the loop

**while**  $x_2^3 + x_1 - x_2^2 > 0$  **do**  $(x_1, x_2) \leftarrow (a_1, a_2)$

where  $a_1 = ((-x_2^2 + x_1)^2 + x_2)^2 - 2 \cdot x_2^2 + 2 \cdot x_1$  and  $a_2 = (-x_2^2 + x_1)^2 + x_2$ , i.e.,  $\varphi = (x_2^3 + x_1 - x_2^2 > 0)$  and  $\vec{a} = (a_1, a_2)$ . Then  $\vec{a}$  induces the endomorphism  $\tilde{a}$  with  $\tilde{a}(x_1) = a_1$  and  $\tilde{a}(x_2) = a_2$ . So we have  $\tilde{a}(2 \cdot x_1 + x_2^3) = (2 \cdot x_1 + x_2^3)(\vec{a}) = 2 \cdot a_1 + a_2^3$ .

For tuples of numbers (e.g.,  $\vec{c} = (5, 2)$ ), the endomorphism  $\tilde{c}$  is  $\tilde{c}(x_1) = 5$  and  $\tilde{c}(x_2) = 2$ . Thus, we have  $\tilde{c}(x_2^3 + x_1 - x_2^2) = (x_2^3 + x_1 - x_2^2)(5, 2) = 2^3 + 5 - 2^2 = 9$ .

We extend the application of endomorphisms  $\eta : \mathcal{S}[\vec{x}] \rightarrow \mathcal{S}[\vec{x}]$  to vectors of polynomials  $\vec{a} = (a_1, \dots, a_d)$  by defining  $\eta(\vec{a}) = (\eta(a_1), \dots, \eta(a_d))$  and to formulas  $\varphi \in \text{Th}_{\exists}(\mathcal{S})$  by defining  $\eta(\varphi) = \varphi(\eta(\vec{x}))$ , i.e.,  $\eta(\varphi)$  results from  $\varphi$  by applying  $\eta$  to all polynomials that occur in  $\varphi$ . This allows us to transform  $(\varphi, \vec{a})$  into a new loop  $\text{Tr}_{\eta}(\varphi, \vec{a})$  using any automorphism  $\eta \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ .

<sup>6</sup> So we have  $\eta(c \cdot p + c' \cdot p') = c \cdot \eta(p) + c' \cdot \eta(p')$ ,  $\eta(1) = 1$ , and  $\eta(p \cdot p') = \eta(p) \cdot \eta(p')$  for all  $c, c' \in \mathcal{S}$  and all  $p, p' \in \mathcal{S}[\vec{x}]$ .

**Definition 5 (Tr).** Let  $\eta \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ . We define  $\text{Tr}_{\eta}(\varphi, \vec{a}) = (\varphi', \vec{a}')$  where

$$\varphi' = \eta^{-1}(\varphi) \quad \text{and} \quad \vec{a}' = (\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x}).^7$$

*Example 6 (Transforming Loops).* We transform the loop  $(\varphi, \vec{a})$  from [Ex. 4](#) with the automorphism  $\eta$  from [Ex. 3](#). We obtain  $\text{Tr}_{\eta}(\varphi, \vec{a}) = (\varphi', \vec{a}')$  where

$$\begin{aligned} \varphi' &= \eta^{-1}(\varphi) = ((\eta^{-1}(x_2))^3 + \eta^{-1}(x_1) - (\eta^{-1}(x_2))^2 > 0) \\ &= (x_1^3 + x_1^2 + x_2 - x_1^2 > 0) = (x_1^3 + x_2 > 0) \quad \text{and} \\ \vec{a}' &= ((\eta^{-1} \circ \tilde{a} \circ \eta)(x_1), (\eta^{-1} \circ \tilde{a} \circ \eta)(x_2)) = (\eta^{-1}(\tilde{a}(x_2)), \eta^{-1}(\tilde{a}(x_1 - x_2^2))) \\ &= (\eta^{-1}(a_2), \eta^{-1}(a_1 - a_2^2)) = (x_1 + x_2^2, 2 \cdot x_2). \end{aligned}$$

So the resulting transformed loop is  $(x_1^3 + x_2 > 0, (x_1 + x_2^2, 2 \cdot x_2))$ . Note that while the original loop  $(\varphi, \vec{a})$  is neither triangular nor weakly non-linear, the resulting transformed loop is *twn*. Also note that we used a *non-linear* automorphism with  $\eta(x_2) = x_1 - x_2^2$  for the transformation.

While the above example shows that our transformation can indeed transform non-*twn*-loops into *twn*-loops, it remains to prove that this transformation preserves (non-)termination. Then we can use our techniques for termination analysis of *twn*-loops for *twn-transformable*-loops as well, i.e., for all loops  $(\varphi, \vec{a})$  where  $\text{Tr}_{\eta}(\varphi, \vec{a})$  is *twn* for some automorphism  $\eta$ . (The question how to find such automorphisms will be addressed in [Sect. 3.2](#).)

As a first step, by [Lemma 7](#), our transformation is “compatible” with the operation  $\circ$  of the group  $\text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ , i.e., it is an *action*.

**Lemma 7.** *Tr is an action of  $\text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$  on loops, i.e., for  $\eta_1, \eta_2 \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$*

$$\text{Tr}_{\text{id}_{\mathcal{S}[\vec{x}]}}(\varphi, \vec{a}) = (\varphi, \vec{a}) \quad \text{and} \quad \text{Tr}_{\eta_1 \circ \eta_2}(\varphi, \vec{a}) = \text{Tr}_{\eta_2}(\text{Tr}_{\eta_1}(\varphi, \vec{a})).$$

The next lemma shows that a witness for non-termination of  $(\varphi, \vec{a})$  is transformed by  $\eta(\vec{x})$  into a witness for non-termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$ .

**Lemma 8.** *If  $\vec{c}$  witnesses non-termination of  $(\varphi, \vec{a})$ , then  $\hat{\eta}(\vec{c})$  witnesses non-termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$ . Here,  $\hat{\eta} : \mathcal{S}^d \rightarrow \mathcal{S}^d$  maps  $\vec{c}$  to  $\hat{\eta}(\vec{c}) = \tilde{c}(\eta(\vec{x})) = (\eta(\vec{x}))(\vec{c})$ .*

*Example 9 (Transforming Witnesses).* For the tuple  $\vec{c} = (5, 2)$  from [Ex. 4](#) and the automorphism  $\eta$  from [Ex. 3](#) with  $\eta(x_1) = x_2$  and  $\eta(x_2) = x_1 - x_2^2$ , we obtain

$$\hat{\eta}(\vec{c}) = (\eta(x_1), \eta(x_2))(\vec{c}) = (2, 5 - 2^2) = (2, 1).$$

As  $\vec{c} = (5, 2)$  witnesses non-termination of [Ex. 4](#),  $\hat{\eta}(\vec{c}) = (2, 1)$  witnesses non-termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$  due to [Lemma 8](#).

Finally, [Thm. 10](#) states that transforming a loop preserves (non-)termination.

**Theorem 10 (Tr Preserves Termination).** *If  $\eta \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ , then  $(\varphi, \vec{a})$  terminates iff  $\text{Tr}_{\eta}(\varphi, \vec{a})$  terminates. Furthermore,  $\hat{\eta}$  is a bijection between the respective sets of witnesses for non-termination.*

<sup>7</sup> In other words, we have  $\vec{a}' = (\eta(\vec{x}))(\vec{a})(\eta^{-1}(\vec{x}))$ , since  $(\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x}) = \eta^{-1}(\eta(\vec{x})[\vec{x}/\vec{a}]) = \eta(\vec{x})[\vec{x}/\vec{a}][\vec{x}/\eta^{-1}(\vec{x})] = (\eta(\vec{x}))(\vec{a})(\eta^{-1}(\vec{x}))$ .

Up to now, we only transformed a loop  $(\varphi, \vec{a})$  on  $\mathcal{S}^d$  using elements of  $\text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ . However, we can also transform it into the loop  $\text{Tr}_{\eta}(\varphi, \vec{a})$  on  $\mathbb{R}_{\mathbb{A}}^d$  using an automorphism  $\eta \in \text{Aut}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$ . Nevertheless, our goal remains to prove termination on  $\mathcal{S}^d$  instead of  $\mathbb{R}_{\mathbb{A}}^d$ , which is not equivalent in general. Thus, in Sect. 5 we will show how to analyze termination of loops on certain subsets  $F$  of  $\mathbb{R}_{\mathbb{A}}^d$ . This allows us to analyze termination of  $(\varphi, \vec{a})$  on  $\mathcal{S}^d$  by checking termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$  on the subset  $\widehat{\eta}(\mathcal{S}^d) \subseteq \mathbb{R}_{\mathbb{A}}^d$  instead.

By our definition of loops over a ring  $\mathcal{S}$ , we have  $\vec{a}(\vec{c}) \in \mathcal{S}^d$  for all  $\vec{c} \in \mathcal{S}^d$ , i.e.,  $\mathcal{S}^d$  is  $\vec{a}$ -invariant. This property is preserved by our transformation.

**Definition 11 ( $\vec{a}$ -Invariance).** *Let  $(\varphi, \vec{a})$  be a loop on  $\mathcal{S}^d$  and let  $F \subseteq \mathcal{S}^d$ . We call  $F$   $\vec{a}$ -invariant or update-invariant if for all  $\vec{c} \in F$  we have  $\vec{a}(\vec{c}) \in F$ .*

**Lemma 12.** *Let  $(\varphi, \vec{a})$  be a loop on  $\mathcal{S}^d$ , let  $F \subseteq \mathcal{S}^d$  be  $\vec{a}$ -invariant, and let  $\eta \in \text{Aut}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$ . Furthermore, let  $\text{Tr}_{\eta}(\varphi, \vec{a}) = (\varphi', \vec{a}')$ . Then  $\widehat{\eta}(F)$  is  $\vec{a}'$ -invariant.*

Recall that our goal is to reduce termination to a  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -formula. Clearly, termination on  $F$  cannot be encoded with such a formula if  $F$  cannot be defined via  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ . Thus, we require that  $F$  is  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable.

**Definition 13 ( $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -Definability).** *A set  $F \subseteq \mathbb{R}_{\mathbb{A}}^d$  is  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable if there is a  $\psi \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  with free variables  $\vec{x}$  such that for all  $\vec{c} \in \mathbb{R}_{\mathbb{A}}^d$*

$$\vec{c} \in F \quad \text{iff} \quad \psi(\vec{c}) \text{ is valid.}$$

An example for a  $\text{Th}_{\exists}(\mathbb{Z}, \mathbb{R}_{\mathbb{A}})$ -definable set is  $\{(a, 0, a) \mid a \in \mathbb{Z}\}$ , which is characterized by the formula  $\exists a \in \mathbb{Z}. x_1 = a \wedge x_2 = 0 \wedge x_3 = a$ .

To analyze termination of  $(\varphi, \vec{a})$  on  $\mathcal{S}^d$ , we can analyze termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$  on  $\widehat{\eta}(\mathcal{S}^d) \subseteq \mathbb{R}_{\mathbb{A}}^d$  instead. The reason is that  $\vec{c} \in \mathcal{S}^d$  witnesses non-termination of  $(\varphi, \vec{a})$  iff  $\widehat{\eta}(\vec{c})$  witnesses non-termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$  due to Thm. 10, i.e.,  $\mathcal{S}^d$  contains a witness for non-termination of  $(\varphi, \vec{a})$  iff  $\widehat{\eta}(\mathcal{S}^d)$  contains a witness for non-termination of  $\text{Tr}_{\eta}(\varphi, \vec{a})$ . While  $\mathcal{S}^d$  is clearly  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable, the following lemma shows that  $\widehat{\eta}(\mathcal{S}^d)$  is  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable, too. More precisely,  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definability is preserved by polynomial endomorphisms.

**Lemma 14.** *Let  $\mathbb{Z} \leq \mathcal{S} \leq \mathbb{R}_{\mathbb{A}}$  and let  $\eta \in \text{End}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$ . If  $F \subseteq \mathbb{R}_{\mathbb{A}}^d$  is  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable then so is  $\widehat{\eta}(F)$ .*

*Example 15.* The set  $\mathbb{Z}^2$  is  $\text{Th}_{\exists}(\mathbb{Z}, \mathbb{R}_{\mathbb{A}})$ -definable, as we have  $(x_1, x_2) \in \mathbb{Z}^2$  iff

$$\exists a, b \in \mathbb{Z}. x_1 = a \wedge x_2 = b.$$

Let  $\eta \in \text{End}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$  with  $\eta(x_1) = \frac{1}{2} \cdot x_1^2 + x_2^2$  and  $\eta(x_2) = x_2^2$ . Then  $\widehat{\eta}(\mathbb{Z}^2)$  is also  $\text{Th}_{\exists}(\mathbb{Z}, \mathbb{R}_{\mathbb{A}})$ -definable, because for  $x_1, x_2 \in \mathbb{R}_{\mathbb{A}}$ , we have  $(x_1, x_2) \in \eta(\mathbb{Z}^2)$  iff

$$\exists y_1, y_2 \in \mathbb{R}_{\mathbb{A}}, a, b \in \mathbb{Z}. y_1 = a \wedge y_2 = b \wedge x_1 = \frac{1}{2} \cdot y_1^2 + y_2^2 \wedge x_2 = y_2^2.$$

The following theorem shows that instead of regarding *solvable loops* [47], w.l.o.g. we can restrict ourselves to *twn-loops*. The reason is that every solvable loop with real eigenvalues can be transformed into a *twn-loop* by a *linear* automorphism  $\eta$ , i.e., the degree  $\deg(\eta)$  of  $\eta$  is 1, where  $\deg(\eta) = \max_{1 \leq i \leq d} \deg(\eta(x_i))$ .

**Theorem 16.** *Let  $(\varphi, \vec{a})$  be a solvable loop with real eigenvalues. Then one can compute a linear automorphism  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n.*

We recapitulate our most important results on  $\text{Tr}$  in the following corollary.

**Corollary 17 (Properties of  $\text{Tr}$ ).** *Let  $(\varphi, \vec{a})$  be a loop,  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$ ,  $\text{Tr}_\eta(\varphi, \vec{a}) = (\varphi', \vec{a}')$ , and  $F \subseteq \mathcal{S}^d$  be  $\vec{a}$ -invariant and  $\text{Th}_\exists(\mathcal{S}, \mathbb{R}_\mathbb{A})$ -definable. Then*

1.  $\hat{\eta}(F) \subseteq \mathbb{R}_\mathbb{A}^d$  is  $\vec{a}'$ -invariant and  $\text{Th}_\exists(\mathcal{S}, \mathbb{R}_\mathbb{A})$ -definable,
2.  $(\varphi, \vec{a})$  terminates on  $F$  iff  $(\varphi', \vec{a}')$  terminates on  $\hat{\eta}(F)$ , and
3.  $\vec{c} \in F$  witnesses non-termination of  $(\varphi, \vec{a})$  iff  $\hat{\eta}(\vec{c}) \in \hat{\eta}(F)$  witnesses non-termination of  $(\varphi', \vec{a}')$ .

### 3.2 Finding Automorphisms to Transform Loops into *tw*n-Form

The goal of the transformation from Sect. 3.1 is to transform  $(\varphi, \vec{a})$  into *tw*n-form, such that termination of the resulting loop  $\text{Tr}_\eta(\varphi, \vec{a})$  can be analyzed by the technique which will be presented in Sect. 4 and 5. Hence, the remaining challenge is to find a suitable automorphism  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n. In this section, we will present two techniques to check the existence of such automorphisms *constructively*, i.e., these techniques can also be used to compute such automorphisms.

Note that the search for suitable automorphisms is closely related to the question if a polynomial endomorphism can be conjugated into a “de Jonquières”-automorphism, a difficult question from algebraic geometry [14]. So future advances in this field may help to improve the results of the current section.

The first technique (Thm. 20) reduces the search for a suitable automorphism of bounded degree to  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$ . It is known that for any automorphism the degree of its inverse has an upper bound in terms of the length  $d$  of  $\vec{x}$ , see [14, Cor. 2.3.4].

**Theorem 18.** *Let  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$ . Then we have  $\deg(\eta^{-1}) \leq (\deg(\eta))^{d-1}$ .*

By Thm. 18, checking if an endomorphism is indeed an automorphism can be reduced to  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$ . To do so, one encodes the existence of suitable coefficients of the polynomials  $\eta^{-1}(x_1), \dots, \eta^{-1}(x_d)$ , which all have at most degree  $(\deg(\eta))^{d-1}$ .

**Lemma 19.** *Let  $\eta \in \text{End}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$ . Then the question whether  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  holds is reducible to  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$ .*

Based on Lemma 19, we now present our first technique to find an automorphism  $\eta$  that transforms a loop into *tw*n-form.

**Theorem 20 (Tr with Automorphisms of Bounded Degree).** *For any  $\delta \geq 0$ , the question whether there exists an  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  with  $\deg(\eta) \leq \delta$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n is reducible to  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$ .*



So if the degree of  $\eta$  is bounded a priori, then it is decidable whether there exists an  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n, since  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$  is decidable.

We call a loop *tw*n-transformable if there is an  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n. By [Thm. 20](#), *tw*n-transformability is *semi-decidable*, since one can increment  $\delta$  until a suitable automorphism is found. So in other words, any loop which is transformable to a *tw*n-loop can be transformed via [Thm. 20](#).

We call our transformation *Tr* *complete* for a class of loops if *every* loop from this class is *tw*n-transformable. For such classes of loops, a suitable automorphism is *computable* by [Thm. 20](#). Together with [Thm. 16](#), we get the following corollary.

**Corollary 21.** *Tr* is complete for solvable loops with real eigenvalues.

Note that for solvable loops  $(\varphi, \vec{a})$ , instead of computing  $\eta$  using [Thm. 20](#), the proof of [Thm. 16](#) yields a more efficient way to compute a linear automorphism  $\eta$  such that  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n. To this end, one computes the Jordan normal form of each  $A_i$  (see [Def. 2](#)), which is possible in polynomial time (see e.g., [[18](#), [45](#)]).

Our second technique to find suitable automorphisms for our transformation is restricted to *linear* automorphisms. In this case, it is decidable whether a loop can be transformed into a *tw*n-loop  $(\varphi', \vec{a}')$  where the monomial for  $x_i$  has the coefficient 1 in each  $a'_i$ . The decision procedure checks whether a certain Jacobian matrix is *strongly nilpotent*, i.e., it is not based on a reduction to  $\text{Th}_\exists(\mathbb{R}_\mathbb{A})$ .

**Definition 22 (Strong Nilpotence).** Let  $J \in (\mathbb{R}_\mathbb{A}[\vec{x}])^{d \times d}$  be a matrix of polynomials. For all  $1 \leq i \leq d$ , let  $\vec{y}^{(i)}$  be a vector of fresh variables.  $J$  is strongly nilpotent if  $\prod_{i=1}^d J[\vec{x}/\vec{y}^{(i)}] = 0^{d \times d}$ , where  $0^{d \times d} \in (\mathbb{R}_\mathbb{A}[\vec{x}])^{d \times d}$  is the zero matrix.

Our second technique is formulated in the following theorem which follows from an existing result in linear algebra [[13](#), [Thm. 1.6.](#)].

**Theorem 23 (Tr with Linear Automorphisms [[13](#), [Thm. 1.6.](#)]).** Let  $(\varphi, \vec{a})$  be a loop. The Jacobian matrix  $(\frac{\partial(a_i - x_i)}{\partial x_j})_{1 \leq i, j \leq d} \in (\mathbb{R}_\mathbb{A}[\vec{x}])^{d \times d}$  is strongly nilpotent iff there exists a linear automorphism  $\eta \in \text{Aut}_{\mathbb{R}_\mathbb{A}}(\mathbb{R}_\mathbb{A}[\vec{x}])$  with

$$\text{Tr}_\eta(\varphi, \vec{a}) = (\varphi', (x_1 + p_1, \dots, x_d + p_d)) \quad (2)$$

and  $p_i \in \mathbb{R}_\mathbb{A}[x_{i+1}, \dots, x_d]$ . Thus,  $\text{Tr}_\eta(\varphi, \vec{a})$  is *tw*n.

As strong nilpotence of the Jacobian matrix is clearly decidable, [Thm. 23](#) gives rise to a decision procedure for the existence of a linear automorphism that transforms  $(\varphi, \vec{a})$  to the form [\(2\)](#).

*Example 24 (Finding Automorphisms).* The following loop on  $\mathcal{S}^3$  shows how our results enlarge the class of loops where termination is reducible to  $\text{Th}_\exists(\mathcal{S}, \mathbb{R}_\mathbb{A})$ .

$$\mathbf{while} \ 4 \cdot x_2^2 + x_1 + x_2 + x_3 > 0 \ \mathbf{do} \ (x_1, x_2, x_3) \leftarrow (a_1, a_2, a_3) \quad (3)$$

$$\begin{aligned}
\text{with } a_1 &= x_1 + 8 \cdot x_1 \cdot x_2^2 + 16 \cdot x_2^3 + 16 \cdot x_2^2 \cdot x_3 \\
a_2 &= x_2 - x_1^2 - 4 \cdot x_1 \cdot x_2 - 4 \cdot x_1 \cdot x_3 - 4 \cdot x_2^2 - 8 \cdot x_2 \cdot x_3 - 4 \cdot x_3^2 \\
a_3 &= x_3 - 4 \cdot x_1 \cdot x_2^2 - 8 \cdot x_2^3 - 8 \cdot x_2^2 \cdot x_3 + x_1^2 + 4 \cdot x_1 \cdot x_2 + \\
&\quad 4 \cdot x_1 \cdot x_3 + 4 \cdot x_2^2 + 8 \cdot x_2 \cdot x_3 + 4 \cdot x_3^2
\end{aligned}$$

It is clearly *not* in *tw*n-form. To transform it, we use [Thm. 23](#). The Jacobian matrix  $J$  of  $(a_1 - x_1, a_2 - x_2, a_3 - x_3)$  is:

$$\begin{bmatrix}
8 \cdot x_2^2 & 16 \cdot x_1 \cdot x_2 + 48 \cdot x_2^2 + 32 \cdot x_2 \cdot x_3 & 16 \cdot x_2^2 \\
-2 \cdot x_1 - 4 \cdot x_2 - 4 \cdot x_3 & -4 \cdot x_1 - 8 \cdot x_2 - 8 \cdot x_3 & -4 \cdot x_1 - 8 \cdot x_2 - 8 \cdot x_3 \\
-4 \cdot x_2^2 + 2 \cdot x_1 + 4 \cdot x_2 + 4 \cdot x_3 & -8 \cdot x_1 \cdot x_2 - 24 \cdot x_2^2 - 16 \cdot x_2 \cdot x_3 + 4 \cdot x_1 + 8 \cdot x_2 + 8 \cdot x_3 & -8 \cdot x_2^2 + 4 \cdot x_1 + 8 \cdot x_2 + 8 \cdot x_3
\end{bmatrix}$$

One easily checks that  $J$  is strongly nilpotent. Thus, by [Thm. 23](#) the loop can be transformed into *tw*n-form by a linear automorphism. Indeed, consider the linear automorphism  $\eta \in \text{Aut}_{\mathbb{R}_\Delta}(\mathbb{R}_\Delta[\vec{x}])$  induced by the matrix  $M = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 2 & 0 \\ 1 & 2 & 2 \end{bmatrix}$ , i.e.,

$$x_1 \mapsto x_1 + x_2 + x_3, \quad x_2 \mapsto 2 \cdot x_2, \quad x_3 \mapsto x_1 + 2 \cdot x_2 + 2 \cdot x_3$$

with its inverse  $\eta^{-1}$

$$x_1 \mapsto 2 \cdot x_1 - x_3, \quad x_2 \mapsto \frac{1}{2} \cdot x_2, \quad x_3 \mapsto -x_1 - \frac{1}{2} \cdot x_2 + x_3.$$

If we transform our loop with  $\eta$ , we obtain the following *tw*n-loop:

$$\mathbf{while} \ x_1 + x_2^2 > 0 \ \mathbf{do} \ \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \leftarrow \begin{bmatrix} x_1 + x_2^2 \cdot x_3 \\ x_2 - 2 \cdot x_3^2 \\ x_3 \end{bmatrix} \quad (4)$$

If  $\mathcal{S} = \mathbb{R}_\Delta$ , then (4) terminates on  $\mathbb{R}_\Delta^3$  iff (3) terminates on  $\mathbb{R}_\Delta^3$  by [Thm. 10](#). Now assume  $\mathcal{S} = \mathbb{Z}$ , i.e., we are interested in termination of (3) on  $\mathbb{Z}^3$  instead of  $\mathbb{R}_\Delta^3$ . Note that  $\hat{\eta}$  maps  $\mathbb{Z}^3$  to the set of all  $\mathbb{Z}$ -linear combinations of columns of  $M$ , i.e.,

$$\hat{\eta}(\mathbb{Z}^3) = \{a \cdot (1, 0, 1) + b \cdot (1, 2, 2) + c \cdot (1, 0, 2) \mid a, b, c \in \mathbb{Z}\}.$$

By [Cor. 17](#), (4) terminates on  $\hat{\eta}(\mathbb{Z}^3)$  iff (3) terminates on  $\mathbb{Z}^3$ . Moreover,  $\hat{\eta}(\mathbb{Z}^3)$  is  $\text{Th}_{\exists}(\mathbb{Z}, \mathbb{R}_\Delta)$ -definable: We have  $(x_1, x_2, x_3) \in \hat{\eta}(\mathbb{Z}^3)$  iff

$$\exists a, b, c \in \mathbb{Z}. \ x_1 = a \cdot 1 + b \cdot 1 + c \cdot 1 \wedge x_2 = b \cdot 2 \wedge x_3 = a \cdot 1 + b \cdot 2 + c \cdot 2.$$

In the following sections, we will see how to analyze termination of loops like (4) on sets that can be characterized by such formulas.

To summarize, if a loop is *tw*n-transformable, then we can *always* find a suitable automorphism via [Thm. 20](#). So whenever [Thm. 23](#) is applicable, a suitable linear automorphism can also be obtained by using [Thm. 20](#) for some fixed degree  $\delta \geq 1$ . Hence, our first technique from [Thm. 20](#) subsumes our second one from [Thm. 23](#). However, while [Thm. 20](#) is *always* applicable, [Thm. 23](#) is *easier* to apply. The reason is that for [Thm. 20](#) one has to check validity of a possibly *non-linear* formula over the reals, where the degree of the occurring polynomials depends on  $\delta$  and the update  $\vec{a}$  of the loop. So even when searching for a linear automorphism, one may obtain a non-linear formula if the loop is non-linear. In contrast, [Thm. 23](#) only requires linear algebra. So it is preferable to first check whether the loop can be transformed into a *tw*n-loop

$(\varphi', (x_1 + p_1, \dots, x_d + p_d))$  with  $x_i \notin \mathcal{V}(p_i)$  via a linear automorphism. This check is *decidable* due to [Thm. 23](#).

Note that the proof of [Thm. 20](#) and the proof of [[13](#), Thm. 1.6.] which implies [Thm. 23](#) are constructive. Thus, we can not only check the existence of a suitable automorphism, but we can also compute it whenever its existence can be proven.

## 4 Computing Closed Forms

Now we show how to reduce the termination problem of a *tw*n-loop on a  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable set to validity of a formula from  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ . Our reduction exploits that for *tw*n-loops  $(\varphi, \vec{a})$ , there is a closed form for the  $n$ -fold application of  $\vec{a}$  which can be represented as a vector of *poly-exponential expressions*.

As in [[15](#)], we restrict ourselves to *tn*n-loops (instead of *tw*n-loops), because each *tw*n-loop can be transformed into a *tn*n-loop via *chaining*.

**Definition 25 (Chaining).** Chaining a loop  $(\varphi, \vec{a})$  yields  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$ .

Clearly,  $(\varphi, \vec{a})$  terminates iff  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$  terminates. Moreover, if  $(\varphi, \vec{a})$  is a *tw*n-loop then  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$  is a *tn*n-loop, i.e., the coefficient of each  $x_i$  in  $a_i(\vec{a})$  is non-negative. Thus, analogous to [[15](#)], we obtain the following theorem.

**Theorem 26.** *Termination of tw*n-loops is reducible to termination of *tn*n-loops.

It is well known that closed forms for *tn*n-loops are computable, see, e.g., [[27](#)]. The reason is that the bodies of *tn*n-loops correspond to a special case of *C*-finite recurrences, which are known to be solvable [[25](#)]. The resulting closed forms may contain polynomial arithmetic and exponentiation w.r.t.  $n$  (as, e.g.,  $x_1 \leftarrow 2 \cdot x_1$  has the closed form  $x_1 \cdot 2^n$ ) as well as certain piecewise defined functions. For example, the closed form of  $x_1 \leftarrow 1$  is  $x_1^{(n)} = x_1$  if  $n = 0$  and  $x_1^{(n)} = 1$ , otherwise.

We use *poly-exponential expressions* [[15](#)]<sup>8</sup> to represent closed forms where piecewise defined functions are simulated via *characteristic functions*. Given a formula  $\psi$  over  $n$ , its characteristic function  $\llbracket \psi \rrbracket : \mathbb{N} \rightarrow \{0, 1\}$  evaluates to 1 iff  $\psi$  is satisfied (i.e.,  $\llbracket \psi \rrbracket(c) = 1$  if  $\psi[n/c]$  holds and  $\llbracket \psi \rrbracket(c) = 0$ , otherwise). In this way, we avoid handling piecewise defined functions via disjunctions (as done in the closed form of [[27](#)]). Poly-exponential expressions are sums of arithmetic terms over the variables  $\vec{x}$  and the additional designated variable  $n$ , where it is always clear which addend determines the asymptotic growth of the whole expression when increasing  $n$ . This is crucial for our reducibility proof in [Sect. 5](#). In the following, for any set  $X \subseteq \mathbb{R}$ , any  $k \in X$ , and  $\triangleright \in \{\geq, >\}$ , let  $X_{\triangleright k} = \{x \in X \mid x \triangleright k\}$ .

**Definition 27 (Poly-Exponential Expressions).** Let  $\mathcal{C}$  be the set of all finite conjunctions over  $\{n = c, n \neq c \mid c \in \mathbb{N}\}$  where  $n$  is a designated variable. The set of all poly-exponential expressions with the variables  $\vec{x}$  is

$$\text{PE}[\vec{x}] = \left\{ \sum_{j=1}^{\ell} \llbracket \psi_j \rrbracket \cdot \alpha_j \cdot n^{a_j} \cdot b_j^n \mid \ell, a_j \in \mathbb{N}, \psi_j \in \mathcal{C}, \alpha_j \in \mathbb{R}_{\mathbb{A}}[\vec{x}], b_j \in (\mathbb{R}_{\mathbb{A}})_{>0} \right\}.$$

<sup>8</sup> Our definition of poly-exponential expressions slightly generalizes [[15](#), Def. 9] (e.g., we allow polynomials over the variables  $\vec{x}$  instead of just linear combinations).

So an example for a poly-exponential expression is

$$\llbracket n \neq 0 \wedge n \neq 1 \rrbracket \cdot \left(\frac{1}{2} \cdot x_1^2 + \frac{3}{4} \cdot x_2 - 1\right) \cdot n^3 \cdot 3^n + \llbracket n = 1 \rrbracket \cdot (x_1 - x_2).$$

Note that the restriction to triangular loops ensures that the closed form does not contain complex numbers. For example, for arbitrary matrices  $A \in \mathcal{S}^{d \times d}$ , the update  $\vec{x} \leftarrow A \cdot \vec{x}$  is known to admit a closed form as in [Def. 27](#) with complex  $b_j$ 's, whereas real numbers suffice for triangular matrices. Moreover, non-negativity is required to ensure  $b_j > 0$  (e.g., the non-*tnn* loop  $x_1 \leftarrow -x_1$  has the closed form  $x_1 \cdot (-1)^n$ ). So together with triangularity, weak non-linearity ensures that for every *tnn*-loop, one can compute a closed form  $\vec{q} \in (\mathbb{P}\mathbb{E}[\vec{x}])^d$  with  $\vec{q} = \vec{a}^n$ .

*Example 28 (Closed Forms).* Reconsider the loop (4) from [Ex. 24](#). This loop is *tnn* as  $\succ_{(4)} = \{(x_1, x_2), (x_1, x_3), (x_2, x_3)\}$  is well founded. Moreover, every variable  $x_i$  occurs with a non-negative coefficient in its corresponding update  $a_i$ . A closed form for the update after  $n \in \mathbb{N}$  loop iterations is:

$$\vec{q} = \begin{bmatrix} \frac{4}{3} \cdot x_3^5 \cdot n^3 + (-2 \cdot x_3^5 - 2 \cdot x_2 \cdot x_3^3) \cdot n^2 + (x_2^2 \cdot x_3 + \frac{2}{3} \cdot x_3^5 + 2 \cdot x_2 \cdot x_3^3) \cdot n + x_1 \\ -2 \cdot x_3^3 \cdot n + x_2 \\ x_3 \end{bmatrix}$$

## 5 Reducing Termination of *tnn*-Loops to $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$

It is known that the bodies of *tnn*-loops can be linearized [\[38\]](#), i.e., one can reduce termination of a *tnn*-loop  $(\varphi, \vec{a})$  to termination of a linear-update *tnn*-loop  $(\varphi', \vec{a}')$  where  $\varphi'$  may be *non*-linear. Moreover, [\[54\]](#) showed decidability of termination for certain classes of conjunctive linear-update loops over  $\mathbb{R}$ , which cover conjunctive linear-update *tnn*-loops. So, by combining the results of [\[38\]](#) and [\[54\]](#), one can conclude that termination for *conjunctive tnn*-loops over  $\mathbb{R}$  is decidable.

However, we will now present a reduction of termination of *tnn*-loops to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  which applies to *tnn*-loops over *any* ring  $\mathbb{Z} \leq \mathcal{S} \leq \mathbb{R}$  and can handle also *disjunctions* in the loop condition. Moreover, our reduction yields tight complexity results on termination of linear loops over  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}_{\mathbb{A}}$ , and  $\mathbb{R}$ , and on termination of linear-update loops over  $\mathbb{R}_{\mathbb{A}}$  and  $\mathbb{R}$  ([Sect. 6](#)).

The idea of our reduction is similar to [\[15\]](#). However, in [\[15\]](#), we considered conjunctive linear loops over  $\mathbb{Z}$ . In contrast, we now analyze termination of  $(\varphi, \vec{a})$  on an  $\vec{a}$ -invariant  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable subset of  $\mathbb{R}_{\mathbb{A}}^d$  and allow arbitrary propositional formulas and *non*-linearity in the condition. So the correctness proofs differ substantially from [\[15\]](#). For reasons of space, we only show the major steps of our reduction and refer to [App. A](#) for more details.

In the following, let  $(\varphi, \vec{a})$  be *tnn*, let  $F \subseteq \mathbb{R}_{\mathbb{A}}^d$  be  $\vec{a}$ -invariant and  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable by the formula  $\psi_F$ , and let  $\vec{q} \in (\mathbb{P}\mathbb{E}[\vec{x}])^d$  be the closed form of  $\vec{a}^n$ .

We now show how to encode termination of  $(\varphi, \vec{a})$  on  $F$  into a  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -formula. More precisely, we show that there is a function with the following specification that is computable in polynomial time:

$$\begin{aligned}
\text{Input : } & (\varphi, \vec{a}), \vec{q}, \text{ and } \psi_F \text{ as above} \\
\text{Result : } & \text{ a closed formula } \chi \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}}) \text{ such that} \\
& \chi \text{ is valid iff } (\varphi, \vec{a}) \text{ does not terminate on } F
\end{aligned} \tag{5}$$

We use the concept of *eventual non-termination* [8, 52], where the loop condition may be violated finitely often, i.e.,  $\vec{c}$  witnesses eventual non-termination of  $(\varphi, \vec{a})$  if  $\vec{a}^{n_0}(\vec{c})$  witnesses non-termination for some  $n_0 \in \mathbb{N}$ . Clearly,  $(\varphi, \vec{a})$  is non-terminating iff it is eventually non-terminating [40]. The formula  $\chi$  in (5) will encode the existence of a witness for eventual non-termination.

By the definition of  $\vec{q}$ ,  $(\varphi, \vec{a})$  is eventually non-terminating on  $F$  iff

$$\exists \vec{x} \in F, n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \varphi(\vec{q}). \tag{6}$$

*Example 29.* Continuing Ex. 24 and 28, (4) is eventually non-terminating on

$$F = \widehat{\eta}(\mathbb{Z}^3) = \{a \cdot (1, 0, 1) + b \cdot (1, 2, 2) + c \cdot (1, 0, 2) \mid a, b, c \in \mathbb{Z}\}$$

iff there is a corresponding witness  $\vec{c} = (x_1, x_2, x_3)$ , i.e., iff

$$\begin{aligned}
& \exists x_1, x_2, x_3 \in F, n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p > 0, \quad \text{where} \\
& p = \left(\frac{4}{3} \cdot x_3^5\right) \cdot n^3 + (-2 \cdot x_3^5 - 2 \cdot x_2 \cdot x_3^3 + 4 \cdot x_3^4) \cdot n^2 + \\
& \quad (x_2^2 \cdot x_3 + \frac{2}{3} \cdot x_3^5 + 2 \cdot x_2 \cdot x_3^3 - 4 \cdot x_2 \cdot x_3^2) \cdot n + (x_1 + x_2^2).
\end{aligned} \tag{7}$$

Let  $\vec{q}_{norm}$  be like  $\vec{q}$ , but each factor  $\llbracket \psi \rrbracket$  is replaced by 0 if it contains an equation and by 1, otherwise. The reason is that for large enough  $n$ , equations in  $\psi$  become false and negated equations become true. Thus, (6) is equivalent to

$$\exists \vec{x} \in F, n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \varphi(\vec{q}_{norm}). \tag{8}$$

In this way, we obtain *normalized* poly-exponential expressions.

**Definition 30 (Normalized PEs).** We call  $p \in \text{PE}[\vec{x}]$  normalized if it is in

$$\text{NPE}[\vec{x}] = \left\{ \sum_{j=1}^{\ell} \alpha_j \cdot n^{a_j} \cdot b_j^n \mid \ell, a_j \in \mathbb{N}, \alpha_j \in \mathbb{R}_{\mathbb{A}}[\vec{x}], b_j \in (\mathbb{R}_{\mathbb{A}})_{>0} \right\}.$$

*W.l.o.g., we always assume  $(b_i, a_i) \neq (b_j, a_j)$  if  $i \neq j$ . We define  $\text{NPE} = \text{NPE}[\emptyset]$ .*

As  $\varphi$  is a propositional formula over  $\mathbb{R}_{\mathbb{A}}[\vec{x}]$ -inequations,  $\varphi(\vec{q}_{norm})$  is a propositional formula over  $\text{NPE}[\vec{x}]$ -inequations. By (8), we need to check if there is an  $\vec{x} \in F$  such that  $\varphi(\vec{q}_{norm})$  is valid for large enough  $n$ . To do so, we generalize [15, Lemma 24]. As usual,  $g : \mathbb{N} \rightarrow \mathbb{R}$  dominates  $f : \mathbb{N} \rightarrow \mathbb{R}$  asymptotically ( $f \in o(g)$ ) if for all  $m > 0$  there is an  $n_0 \in \mathbb{N}$  such that  $|f(n)| < m \cdot |g(n)|$  for all  $n \in \mathbb{N}_{>n_0}$ .

**Lemma 31.** *Let  $b_1, b_2 \in (\mathbb{R}_{\mathbb{A}})_{>0}$  and  $a_1, a_2 \in \mathbb{N}$ . If  $(b_2, a_2) >_{lex} (b_1, a_1)$ , then  $n^{a_1} \cdot b_1^n \in o(n^{a_2} \cdot b_2^n)$ , where  $(b_2, a_2) >_{lex} (b_1, a_1)$  iff  $b_2 > b_1$  or  $b_2 = b_1 \wedge a_2 > a_1$ .*

In the following, let  $p \geq 0$  or  $p > 0$  occur in  $\varphi(\vec{q}_{norm})$ . Then we can order the coefficients of  $p$  according to the asymptotic growth of their addends w.r.t.  $n$ .

**Definition 32 (Ordering Coefficients).** Marked coefficients are of the form  $\alpha^{(b,a)}$  where  $\alpha \in \mathbb{R}_{\mathbb{A}}[\vec{x}]$ ,  $b \in (\mathbb{R}_{\mathbb{A}})_{>0}$ , and  $a \in \mathbb{N}$ . We define  $\text{unmark}(\alpha^{(b,a)}) = \alpha$  and  $\alpha_2^{(b_2, a_2)} \succ_{coef} \alpha_1^{(b_1, a_1)}$  if  $(b_2, a_2) >_{lex} (b_1, a_1)$ . Let  $p = \sum_{j=1}^{\ell} \alpha_j \cdot n^{a_j} \cdot b_j^n \in \text{NPE}[\vec{x}]$ , where  $\alpha_j \neq 0$  for all  $1 \leq j \leq \ell$ . Then the marked coefficients of  $p$  are  $\text{coefs}(p) = \{0^{(1,0)}\}$  if  $\ell = 0$  and  $\text{coefs}(p) = \{\alpha_j^{(b_j, a_j)} \mid 0 \leq j \leq \ell\}$ , otherwise.

*Example 33.* Continuing [Ex. 29](#),  $\text{coefs}(p)$  is  $\{\alpha_1^{(1,3)}, \alpha_2^{(1,2)}, \alpha_3^{(1,1)}, \alpha_4^{(1,0)}\}$  where:

$$\begin{aligned} \alpha_1 &= \frac{4}{3} \cdot x_3^5 & \alpha_2 &= -2 \cdot x_3^5 - 2 \cdot x_2 \cdot x_3^3 + 4 \cdot x_3^4 \\ \alpha_3 &= x_2^2 \cdot x_3 + \frac{2}{3} \cdot x_3^5 + 2 \cdot x_2 \cdot x_3^3 - 4 \cdot x_2 \cdot x_3^2 & \alpha_4 &= x_2^2 + x_1 \end{aligned}$$

Note that  $p(\vec{c}) \in \text{NPE}$  for any  $\vec{c} \in \mathbb{R}_{\mathbb{A}}^d$ , i.e., the only variable in  $p(\vec{c})$  is  $n$ . Now the  $\succ_{\text{coef}}$ -maximal addend determines the asymptotic growth of  $p(\vec{c})$ :

$$o(p(\vec{c})) = o(k \cdot n^a \cdot b^n) \quad \text{where } k^{(b,a)} = \max_{\succ_{\text{coef}}}(\text{coefs}(p(\vec{c}))). \quad (9)$$

Note that (9) would be incorrect for the case  $k = 0$  if we replaced  $o(p(\vec{c})) = o(k \cdot n^a \cdot b^n)$  with  $o(p(\vec{c})) = o(n^a \cdot b^n)$  as  $o(0) = \emptyset \neq o(1)$ . Obviously, (9) implies

$$\exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \text{sign}(p(\vec{c})) = \text{sign}(k) \quad (10)$$

where  $\text{sign}(0) = 0$ ,  $\text{sign}(k) = 1$  if  $k > 0$ , and  $\text{sign}(k) = -1$  if  $k < 0$ . This already allows us to reduce eventual non-termination to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  if  $\varphi$  is an atom.

**Lemma 34.** *Given  $p \in \text{NPE}[\vec{x}]$  and  $\triangleright \in \{\geq, >\}$ , one can reduce validity of*

$$\exists \vec{x} \in F, n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p \triangleright 0 \quad (11)$$

*to validity of a closed formula from  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  in polynomial time.<sup>9</sup>*

More precisely, (11) can be reduced to a formula  $\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(p \triangleright 0)$ , where  $\text{red}(p \triangleright 0)$  is constructed as follows. By (10), we have  $p(\vec{c}) > 0$  for large enough values of  $n$  iff the coefficient of the asymptotically fastest-growing addend  $\alpha(\vec{c}) \cdot n^a \cdot b^n$  of  $p$  that does not vanish (i.e., where  $\alpha(\vec{c}) \neq 0$ ) is *positive*. Similarly, we have  $p(\vec{c}) < 0$  for large enough  $n$  iff  $\alpha(\vec{c}) < 0$ . If *all* addends of  $p$  vanish when instantiating  $\vec{x}$  with  $\vec{c}$ , then  $p(\vec{c}) = 0$ . In other words, (11) holds iff there is a  $\vec{c} \in F$  such that  $\text{unmark}(\max_{\succ_{\text{coef}}}(\text{coefs}(p(\vec{c})))) \triangleright 0$ . To express this in  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ , let  $\alpha_1, \dots, \alpha_\ell$  be the coefficients of  $p$ , ordered according to the asymptotic growth of the respective addends where  $\alpha_1$  belongs to the fastest-growing addend. Then

$$\begin{aligned} \text{red}(p > 0) &\text{ is } \bigvee_{j=1}^{\ell} \left( \alpha_j > 0 \wedge \bigwedge_{i=1}^{j-1} \alpha_i = 0 \right) \\ \text{and } \text{red}(p \geq 0) &\text{ is } \text{red}(p > 0) \vee \bigwedge_{i=1}^{\ell} \alpha_i = 0. \end{aligned}$$

Hence, (11) is equivalent to  $\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(p \triangleright 0)$ .

*Example 35 (Reducing Eventual Non-Termination to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ ).* We finish [Ex. 33](#) resp. [24](#) for  $\mathcal{S} = \mathbb{Z}$ , where  $\text{unmark}(\max_{\succ_{\text{coef}}}(\text{coefs}(p))) = \frac{4}{3} \cdot x_3^5$  and  $\psi_F$  is

$$\exists a, b, c \in \mathbb{Z}. x_1 = a + b + c \wedge x_2 = b \cdot 2 \wedge x_3 = a + b \cdot 2 + c \cdot 2.$$

Thus, (7) is valid iff  $\exists x_1, x_2, x_3 \in \mathbb{R}_{\mathbb{A}}. \psi_F \wedge \text{red}(p > 0)$  is valid where

$$\begin{aligned} \text{red}(p > 0) &= \alpha_1 > 0 && \vee (\alpha_2 > 0 \wedge \alpha_1 = 0) \\ &\vee (\alpha_3 > 0 \wedge \alpha_1 = \alpha_2 = 0) && \vee (\alpha_4 > 0 \wedge \alpha_1 = \alpha_2 = \alpha_3 = 0). \end{aligned}$$

<sup>9</sup> More precisely, the reduction of [Lemma 34](#) and of the following [Thm. 36](#) takes polynomially many steps in the size of the input of the function in (5).

Then  $[x_1/1, x_2/0, x_3/1]$  satisfies  $\psi_F \wedge \alpha_1 > 0$  as  $(1, 0, 1) \in F$  (see Ex. 29) and  $(\frac{4}{3} \cdot x_3^5) [x_1/1, x_2/0, x_3/1] > 0$ . Thus,  $(1, 0, 1)$  witnesses eventual non-termination of (4). So the original loop (3) is non-terminating on  $\mathbb{Z}^3$  by Cor. 17 resp. Thm. 10.

Now we lift our reduction to propositional formulas. Note that a version of the following Thm. 36 that only covers conjunctions is a direct corollary of Lemma 34. To handle disjunctions, the proof of Thm. 36 exploits the crucial additional insight that a *tnn*-loop  $(\varphi \vee \varphi', \vec{a})$  terminates iff  $(\varphi, \vec{a})$  and  $(\varphi', \vec{a})$  terminate, which is not true in general (as, e.g., witnessed by the loop  $(x > 0 \vee -x > 0, -x)$ ).

**Theorem 36.** *Given a propositional formula  $\xi$  over the atoms  $\{p \triangleright 0 \mid p \in \text{NPE}[\vec{x}], \triangleright \in \{\geq, >\}\}$ , one can reduce validity of*

$$\exists \vec{x} \in F, n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \xi \quad (12)$$

*to validity of a formula  $\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(\xi) \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  in polynomial time.*

Here,  $\text{red}(\xi)$  results from replacing each atom  $p \triangleright 0$  in  $\xi$  by  $\text{red}(p \triangleright 0)$ .

Thm. 36 shows that the function (5) is computable (in polynomial time). This leads to the main result of this section.

**Theorem 37 (Reducing Termination).** *Termination of *tnn*-loops (resp. *tw*-loops) on  $\vec{a}$ -invariant and  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable sets is reducible to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ .*

However, in general this reduction is not computable in polynomial time, not even in exponential time. The reason is that closed forms  $\vec{q}$  of  $\vec{a}^n$  cannot be computed in exponential time if the update  $\vec{a}$  contains *non-linear* terms. For example, consider the following *tnn*-loop:

$$\text{while true do } \vec{x} \leftarrow (d \cdot x_1, x_1^d, \dots, x_{d-2}^d, x_{d-1}^d) \quad (13)$$

The closed form for  $x_i^{(n)}$  is  $q_i = d^{(d^{i-1} \cdot (n-i+1))} \cdot x_1^{(d^{i-1})}$  for all  $n \geq d$ . Note that  $\log d^{(d^{d-1})}$  grows faster in  $d$  than any expression of the form  $c^d$ , where  $c \in \mathbb{N}$ . Thus, the closed form  $q_d \in \mathbb{PE}[\vec{x}]$  for  $x_d^{(n)}$  contains constants whose logarithm grows faster than any expression  $c^d$ . Hence,  $q_d$  cannot be computed in exponential time. As mentioned at the beginning of this section, the bodies of *tnn*-loops could also be linearized [38]. However, since the linearization of (13) contains these constants as well, it cannot be computed in exponential time, either.

Note that our reduction also works if  $\mathcal{S} = \mathbb{R}$ , i.e., termination over  $\mathbb{R}$  is reducible to  $\text{Th}_{\exists}(\mathbb{R}, \mathbb{R}_{\mathbb{A}})$ . As  $\mathbb{R}$  and  $\mathbb{R}_{\mathbb{A}}$  are *elementary equivalent*, i.e., a first-order formula is valid over  $\mathbb{R}$  iff it is valid over  $\mathbb{R}_{\mathbb{A}}$ , we get the following corollary.

**Corollary 38 ((Semi-)Decidability of (Non-)Termination).** *Let  $(\varphi, \vec{a})$  be a *tw*-loop and let  $F \subseteq \mathbb{R}_{\mathbb{A}}^d$  be  $\vec{a}$ -invariant and  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable.*

- (a) *The loop  $(\varphi, \vec{a})$  terminates over  $\mathbb{R}_{\mathbb{A}}$  iff it terminates over  $\mathbb{R}$ .*
- (b) *Termination of  $(\varphi, \vec{a})$  on  $F$  is decidable if  $\mathcal{S} = \mathbb{R}_{\mathbb{A}}$  or  $\mathcal{S} = \mathbb{R}$ .*
- (c) *Non-termination of  $(\varphi, \vec{a})$  on  $F$  is semi-decidable if  $\mathcal{S} = \mathbb{Z}$  or  $\mathcal{S} = \mathbb{Q}$ .*

Moreover, by [Thm. 20](#) it is semi-decidable if a loop is *twN-transformable*. For *conjunctive twN-loops*, [Cor. 38 \(b\)](#) also follows from combining [\[38\]](#) and [\[54\]](#).

Our technique does not yield witnesses for non-termination, but the formula constructed by [Thm. 36](#) describes *all* witnesses for *eventual* non-termination. So, the set of witnesses of *eventual* non-termination is  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ -definable whereas in general, the set of witnesses of non-termination is not (see [\[12\]](#)).

**Lemma 39.** *Let  $\xi = \varphi(\vec{q}_{norm})$ . Then  $\vec{c} \in \mathbb{R}_{\mathbb{A}}^d$  witnesses eventual non-termination of  $(\varphi, \vec{a})$  on  $F$  iff  $\psi_F(\vec{c}) \wedge (\text{red}(\xi))(\vec{c})$ .*

However, in [\[22\]](#) we showed how to compute witnesses for non-termination from witnesses for eventual non-termination of *twN-loops*. Thus, [Lemma 39](#) combined with our results from [\[22\]](#) yields a technique to enumerate all witnesses for non-termination.

If  $(\varphi, \vec{a})$  results from the original loop by first transforming it into *twN*-form ([Sect. 3](#)) and by subsequently chaining it in order to obtain a loop in *tnn*-form ([Sect. 4](#)), then our approach can also be used to obtain witnesses for eventual non-termination of the original loop. In other words, one can compute a witness for the original loop from the witness for the transformed loop as in [Cor. 17](#), since chaining clearly preserves witnesses for eventual non-termination. [Alg. 1](#) summarizes our technique to check termination of *twN-transformable-loops*.

**Algorithm 1: Checking Termination**

<p><b>Input:</b> a <i>twN-transformable-loop</i> <math>(\varphi, \vec{a})</math> and <math>\psi_F \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})</math>  <b>Result:</b> <math>\top</math> resp. <math>\perp</math> if (non-)termination of <math>(\varphi, \vec{a})</math> on <math>F</math> is proven, ? otherwise  <math>(\varphi, \vec{a}) \leftarrow \text{Tr}_{\eta}(\varphi, \vec{a})</math>, <math>\psi_F \leftarrow \psi_{\vec{\eta}(F)}</math>, such that <math>(\varphi, \vec{a})</math> becomes <i>twN</i>  <math>(\varphi, \vec{a}) \leftarrow (\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))</math>, such that <math>(\varphi, \vec{a})</math> becomes <i>tnn</i>  <math>\vec{q} \leftarrow</math> closed form of <math>\vec{a}^n</math>  if (un)satisfiability of <math>\psi_F \wedge \text{red}(\varphi(\vec{q}_{norm}))</math> cannot be proven <b>then return</b> ?  <b>if</b> <math>\psi_F \wedge \text{red}(\varphi(\vec{q}_{norm}))</math> is satisfiable <b>then return</b> <math>\perp</math> <b>else return</b> <math>\top</math></p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

## 6 Complexity Analysis

We now analyze the complexity of our technique. We first regard *linear-update* loops, i.e., where the update is of the form  $\vec{x} \leftarrow A \cdot \vec{x} + \vec{b}$  with  $A \in \mathcal{S}^{d \times d}$  and  $\vec{b} \in \mathcal{S}^d$ . More precisely, we show that termination of linear loops with real spectrum is Co-NP-complete if  $\mathcal{S} \in \{\mathbb{Z}, \mathbb{Q}, \mathbb{R}_{\mathbb{A}}\}$  and that termination of linear-update loops with real spectrum is  $\forall\mathbb{R}$ -complete if  $\mathcal{S} = \mathbb{R}_{\mathbb{A}}$ . Since our proof is based on a reduction to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ , and  $\mathbb{R}_{\mathbb{A}}$  and  $\mathbb{R}$  are elementary equivalent, our results also hold if the program variables range over  $\mathbb{R}$ .

For our complexity results, we assume the usual dense encoding of univariate polynomials, i.e., a polynomial of degree  $k$  is represented as a list of  $k + 1$  coefficients. As discussed in [\[46\]](#), many problems which are considered to be efficiently solvable become intractable if polynomials are encoded sparsely (e.g., as lists of monomials where each monomial is a pair of its non-zero coefficient and



its degree). With densely encoded polynomials, all common representations of algebraic numbers can be converted into each other in polynomial time [3].

When analyzing linear-update loops, w.l.o.g. we can assume  $\vec{b} = \vec{0}$ , since

$$\mathbf{while} \ \varphi \ \mathbf{do} \ \vec{x} \leftarrow A \cdot \vec{x} + \vec{b} \qquad \text{terminates iff} \qquad (14)$$

$$\mathbf{while} \ \varphi \wedge x_{\vec{b}} = 1 \ \mathbf{do} \ \begin{bmatrix} \vec{x} \\ x_{\vec{b}} \end{bmatrix} \leftarrow \begin{bmatrix} A & \vec{b} \\ \vec{0}^T & 1 \end{bmatrix} \cdot \begin{bmatrix} \vec{x} \\ x_{\vec{b}} \end{bmatrix} \qquad (15)$$

terminates, where  $x_{\vec{b}}$  is a fresh variable (see [23, 40]). Moreover,  $\vec{c}$  witnesses (eventual) non-termination for (14) iff  $\begin{bmatrix} \vec{c} \\ 1 \end{bmatrix}$  witnesses (eventual) non-termination for (15). Note that the only eigenvalue of  $\begin{bmatrix} A & \vec{b} \\ \vec{0}^T & 1 \end{bmatrix}$  whose multiplicity increases in comparison to  $A$  is 1. Thus, to decide termination of linear-update loops with real spectrum, it suffices to decide termination of loops of the form  $(\varphi, A \cdot \vec{x})$  where  $A$  has only real eigenvalues.

Such loops can *always* be transformed into *tw*n-form using our transformation *Tr* from Sect. 3. To compute the required automorphism  $\eta$ , we compute the Jordan normal form  $Q$  of  $A$  together with the corresponding transformation matrix  $T$ , i.e.,  $T$  is an invertible real matrix such that  $A = T^{-1} \cdot Q \cdot T$ . Then  $Q$  is a triangular real matrix whose diagonal consists of the eigenvalues  $\lambda \in \mathbb{R}_{\mathbb{A}}$  of  $A$ . Now we define  $\eta \in \text{End}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$  by  $\eta(\vec{x}) = T \cdot \vec{x}$ . Then  $\eta \in \text{Aut}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$  has the inverse  $\eta^{-1}(\vec{x}) = T^{-1} \cdot \vec{x}$ . Thus,  $\text{Tr}_{\eta}(\varphi, A \cdot \vec{x})$  is a *tw*n-loop with the update

$$(\eta(\vec{x})) (A \cdot \vec{x}) (\eta^{-1}(\vec{x})) = T \cdot A \cdot T^{-1} \cdot \vec{x} = Q \cdot \vec{x}.$$

The Jordan normal form  $Q$  as well as the matrices  $T$  and  $T^{-1}$  can be computed in polynomial time [18, 45]. Hence, we can decide whether all eigenvalues are real numbers in polynomial time by checking the diagonal entries of  $Q$ . Thus, we obtain the following lemma.

**Lemma 40.** *Let  $(\varphi, A \cdot \vec{x})$  be a linear-update loop.*

- (a) *It is decidable in polynomial time whether  $A$  has only real eigenvalues.*
- (b) *If  $A$  has only real eigenvalues, then we can compute a linear  $\eta \in \text{Aut}_{\mathbb{R}_{\mathbb{A}}}(\mathbb{R}_{\mathbb{A}}[\vec{x}])$  such that  $\text{Tr}_{\eta}(\varphi, A \cdot \vec{x})$  is a linear-update *tw*n-loop in polynomial time.*
- (c) *If  $(\varphi, A \cdot \vec{x})$  is a linear loop, then so is  $\text{Tr}_{\eta}(\varphi, A \cdot \vec{x})$ .*

Hence, the transformation from Sect. 3 is complete for linear(-update) loops with real spectrum, i.e., every such loop can be transformed into a linear(-update) *tw*n-loop. Note that the first part of Lemma 40 yields an efficient check whether a given linear(-update) loop has real spectrum.

As chaining (Def. 25) can clearly be done in polynomial time, w.l.o.g. we may assume that  $\text{Tr}_{\eta}(\varphi, A \cdot \vec{x}) = (\varphi', Q \cdot \vec{x})$  is *tn*n. Next, to analyze termination of a loop, our technique of Sect. 4 computes a closed form for the  $n$ -fold application of the update. For *tn*n-loops of the form  $(\varphi', Q \cdot \vec{x})$  where  $Q$  is a triangular matrix with non-negative diagonal entries, a suitable (i.e., poly-exponential) closed form can be computed in polynomial time analogously to [27, Prop. 5.2]. This closed form is linear in  $\vec{x}$ .

According to our approach in Sect. 5, we now proceed as in Alg. 1 and compute  $\text{red}(\varphi(\vec{q}_{norm})) \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ . The construction of this formula can be done in polynomial time due to Thm. 36. Hence, we get the following lemma.

**Lemma 41.** *Let  $(\varphi, A \cdot \vec{x})$  be a linear-update loop with real spectrum. Then we can compute a formula  $\psi \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$  in polynomial time, such that  $\psi$  is valid iff the loop is non-terminating. If  $\varphi$  is linear, then so is  $\psi$ .*

Note that  $\psi$  is existentially quantified. Hence, if  $\varphi$  (and thus, also  $\psi$ ) is linear and  $\mathcal{S} \in \{\mathbb{Z}, \mathbb{Q}, \mathbb{R}_{\mathbb{A}}, \mathbb{R}\}$ , then invalidity of  $\psi$  is in Co-NP as validity of such formulas is in NP [41]. Thus, we obtain the first main result of this section.

**Theorem 42 (Co-NP-Completeness).** *Termination of linear loops  $(\varphi, A \cdot \vec{x} + \vec{b})$  with real spectrum over  $\mathbb{Z}$ ,  $\mathbb{Q}$ ,  $\mathbb{R}_{\mathbb{A}}$ , and  $\mathbb{R}$  is Co-NP-complete.*

For Co-NP-hardness, let  $\xi$  be a propositional formula over the variables  $\vec{x}$ . Then  $(\xi[x_i/(x_i > 0) \mid 1 \leq i \leq d], \vec{x})$  terminates iff  $\xi$  is unsatisfiable. So Co-NP-hardness follows from Co-NP-hardness of unsatisfiability of propositional formulas.

We now consider linear-update loops with real spectrum (and possibly non-linear loop conditions) on  $\mathbb{R}_{\mathbb{A}}^d$  and  $\mathbb{R}^d$ . Here, non-termination is  $\exists\mathbb{R}$ -complete.

**Definition 43** ( $\exists\mathbb{R}$  [49, 50]). *Let  $\text{Th}_{\exists}(\mathbb{R})_{\top} = \{\psi \in \text{Th}_{\exists}(\mathbb{R}) \mid \psi \text{ is satisfiable}\}$ . The complexity class  $\exists\mathbb{R}$  is the closure of  $\text{Th}_{\exists}(\mathbb{R})_{\top}$  under poly-time-reductions.*

We have  $\text{NP} \subseteq \exists\mathbb{R} \subseteq \text{PSPACE}$ . By Lemma 41, non-termination of linear-update loops with real spectrum is in  $\exists\mathbb{R}$ . It is also  $\exists\mathbb{R}$ -hard since  $(\varphi, \vec{x})$  is non-terminating iff  $\varphi$  is satisfiable. So non-termination is  $\exists\mathbb{R}$ -complete, i.e., termination is Co- $\exists\mathbb{R}$ -complete (where  $\text{Co-}\exists\mathbb{R} = \forall\mathbb{R}$  [50]).

**Theorem 44 ( $\forall\mathbb{R}$ -Completeness).** *Termination of linear-update loops with real spectrum over  $\mathbb{R}_{\mathbb{A}}$  and  $\mathbb{R}$  is  $\forall\mathbb{R}$ -complete.*

Recall that the bodies of *tnn*-loops can be linearized [38]. The loop (13) showed that in general, this linearization is not computable in exponential time. However, if the degree of the polynomials in the update is bounded by a constant  $M$ , or if the number of variables  $d$  is bounded by a constant  $D$ , then the linearization is in EXPTIME (see the proof of Thm. 45). Combining this with Thm. 44 yields Thm. 45 (a). If the number of variables is bounded, then checking validity of an existential formula over the reals is in P (see [2]). So in this case, combining the fact that linearization is in EXPTIME with Lemma 41 yields Thm. 45 (b).

**Theorem 45.** *Let  $M, D \in \mathbb{N}$  be fixed. Termination of *twn*-loops over  $\mathbb{R}_{\mathbb{A}}$  and  $\mathbb{R}$*

- (a) *is in 2-EXPTIME if the maximal degree in the update is at most  $M$ .*
- (b) *is in EXPTIME if the number of variables is at most  $D$ .*

## 7 Related Work and Conclusion

We presented a reduction from termination of *tw*n-loops to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\Delta})$ . This implies decidability of termination over  $\mathcal{S} = \mathbb{R}_{\Delta}$  and  $\mathcal{S} = \mathbb{R}$  and semi-decidability of non-termination over  $\mathcal{S} = \mathbb{Z}$  and  $\mathcal{S} = \mathbb{Q}$ . Moreover, we showed how to transform certain non-*tw*n-loops into *tw*n-form, which generalizes our results to a wider class of loops, including *solvable loops with real eigenvalues*. We also showed that *tw*n-transformability is semi-decidable. Finally, we used our results to prove Co-NP-completeness (resp.  $\forall\mathbb{R}$ -completeness) of termination of linear (resp. linear-update) loops with real spectrum.

*Related Work:* In contrast to *automated* termination analysis (e.g., [1, 4, 5, 7, 9, 10, 16, 19, 20, 21, 29, 30, 31, 32, 42]), we investigate *decidability* of termination for certain classes of loops. Clearly, such decidability results can only be obtained for quite restricted classes of programs.

Nevertheless, many techniques used in automated tools for termination analysis (e.g., the application of ranking functions) focus on similar classes of loops, since such loops occur as sub-programs in (abstractions of) real programs. Tools based on these techniques have turned out to be very successful, also for larger classes of programs. Thus, these tools could benefit from integrating our (semi-)decision procedures and applying them instead of incomplete techniques for any sub-program that can be transformed into a *tw*n-loop.

Related work on decidability of termination also considers related (and often more restricted) classes of loops. For linear conjunctive loops (where the loop condition is a conjunction), termination over  $\mathbb{R}$  [33, 36, 52],  $\mathbb{Q}$  [8], and  $\mathbb{Z}$  [23] is decidable. Tiwari [52] uses the special case of our *tw*n-transformation from Sect. 6 where the loop and the automorphism are linear. In contrast to these techniques, our approach applies to *non-linear* loops with *arbitrary* loop conditions over *various rings*.

*Linearization* is an alternative attempt to handle non-linearity. While the *update* of solvable loops can be linearized [38, 48], the *guard* cannot. Otherwise, one could linearize any loop ( $p = 0, \vec{x}$ ), which terminates over  $\mathbb{Z}$  iff  $p$  has no integer root. With [23], this would imply decidability of Hilbert’s Tenth Problem.

Regarding complexity, [40] proves that termination of conjunctive linear loops over  $\mathbb{Z}$  with update  $\vec{x} \leftarrow A \cdot \vec{x} + \vec{b}$  is in EXPSPACE if  $A$  is diagonalizable resp. in PSPACE if  $|\vec{x}| \leq 4$ . Moreover, [40] states that the techniques from [8, 52] run in polynomial time. So termination of conjunctive linear loops over  $\mathbb{Q}$  and  $\mathbb{R}$  is in P.

Our Co-NP-completeness result is orthogonal to those results as we allow disjunctions in the loop condition. Moreover, Co-NP-completeness also holds for termination over  $\mathbb{Z}$ , whereas [8, 52] only consider termination over  $\mathbb{Q}$  resp.  $\mathbb{R}$ .

In the non-linear case, [34] proves decidability of termination for conjunctive loops on  $\mathbb{R}^d$  for the case that the loop condition defines a compact and connected subset of  $\mathbb{R}^d$ . In [54], decidability of termination of conjunctive linear-update loops on  $\mathbb{R}^d$  with the *non-zero minimum property* is shown, which covers conjunctive linear-update loops with real spectrum. In combination with [38],

this yields a decision procedure for termination of conjunctive *twn*-loops over  $\mathbb{R}$ . For general conjunctive linear-update loops on  $\mathbb{R}^d$  undecidability is conjectured. Furthermore, [37] proves that termination of (not necessarily conjunctive) linear-update loops is decidable if the loop condition describes a compact set. Finally, [55] gives sufficient criteria for (non-)termination of solvable loops and [35] introduces sufficient conditions under which termination of non-deterministic non-linear loops on  $\mathbb{R}^d$  can be reduced to satisfiability of a semi-algebraic system.

For linear-update loops with real spectrum over  $\mathbb{R}$ , we prove  $\forall\mathbb{R}$ -completeness of termination, whereas [54] does not give tight complexity results. The approach from [55] is incomplete, whereas we present a complete reduction from termination to the respective first-order theory. The work in [35] is orthogonal to ours as it only applies to loops that satisfy certain non-trivial conditions. Moreover, we consider loops with arbitrary loop conditions over various rings, whereas [34, 35, 54] only consider conjunctive loops over  $\mathbb{R}$  and [37] only considers loops over  $\mathbb{R}$  where the loop condition defines a compact set.

Finally, several other works exploit the existence of closed forms for solvable (or similar classes of) loops to, e.g., analyze termination for *fixed* inputs or deduce invariants (e.g., [22, 24, 26, 27, 28, 38, 39, 47, 48]). While our approach covers solvable loops with real eigenvalues (by Cor. 21), it also applies to loops which are not solvable, see Ex. 24. Note that our transformation from Sect. 3 may also be of interest for other techniques for solvable or other sub-classes of polynomial loops, as it may be used to extend the applicability of such approaches.

**Acknowledgments** We thank Alberto Fiori for help with the example loop (13) and Arno van den Essen for useful discussions.

## References

- [1] D. Babić, B. Cook, A. J. Hu, and Z. Rakamaric. “Proving Termination of Nonlinear Command Sequences”. In: *Formal Aspects of Computing* 25.3 (2013), pp. 389–403. DOI: [10.1007/s00165-012-0252-5](https://doi.org/10.1007/s00165-012-0252-5).
- [2] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in Real Algebraic Geometry*. Algorithms and Computation in Mathematics 10. Springer, 2006. DOI: [10.1007/3-540-33099-2](https://doi.org/10.1007/3-540-33099-2).
- [3] S. Basu and B. Mishra. “Computational and Quantitative Real Algebraic Geometry”. In: *Handbook of Discrete and Computational Geometry, Third Edition*. Ed. by J. E. Goodman, J. O’Rourke, and C. D. Tóth. Chapman and Hall/CRC, 2017, pp. 969–1002. DOI: [10.1201/9781315119601](https://doi.org/10.1201/9781315119601).
- [4] A. M. Ben-Amram and S. Genaim. “On Multiphase-Linear Ranking Functions”. In: *Proc. CAV ’17*. LNCS 10427. 2017, pp. 601–620. DOI: [10.1007/978-3-319-63390-9\\_32](https://doi.org/10.1007/978-3-319-63390-9_32).
- [5] A. M. Ben-Amram, J. J. Doménech, and S. Genaim. “Multiphase-Linear Ranking Functions and Their Relation to Recurrent Sets”. In: *Proc. SAS ’19*. LNCS 11822. 2019, pp. 459–480. DOI: [10.1007/978-3-030-32304-2\\_22](https://doi.org/10.1007/978-3-030-32304-2_22).
- [6] M. Bozga, R. Iosif, and F. Konečný. “Deciding Conditional Termination”. In: *Logical Methods in Computer Science* 10.3 (2014). DOI: [10.2168/LMCS-10\(3:8\)2014](https://doi.org/10.2168/LMCS-10(3:8)2014).

- [7] A. R. Bradley, Z. Manna, and H. B. Sipma. “Termination of Polynomial Programs”. In: *Proc. VMCAI '05*. LNCS 3385. 2005, pp. 113–129. DOI: [10.1007/978-3-540-30579-8\\_8](https://doi.org/10.1007/978-3-540-30579-8_8).
- [8] M. Braverman. “Termination of Integer Linear Programs”. In: *Proc. CAV '06*. LNCS 4144. 2006, pp. 372–385. DOI: [10.1007/11817963\\_34](https://doi.org/10.1007/11817963_34).
- [9] M. Brockschmidt, B. Cook, and C. Fuhs. “Better Termination Proving through Cooperation”. In: *Proc. CAV '13*. LNCS 8044. 2013, pp. 413–429. DOI: [10.1007/978-3-642-39799-8\\_28](https://doi.org/10.1007/978-3-642-39799-8_28).
- [10] H. Y. Chen, B. Cook, C. Fuhs, K. Nimkar, and P. W. O’Hearn. “Proving Nontermination via Safety”. In: *Proc. TACAS '14*. LNCS 8413. 2014, pp. 156–171. DOI: [10.1007/978-3-642-54862-8\\_11](https://doi.org/10.1007/978-3-642-54862-8_11).
- [11] P. J. Cohen. “Decision Procedures for Real and  $p$ -adic Fields”. In: *Communications on Pure and Applied Mathematics* 22.2 (1969), pp. 131–151. DOI: [10.1002/cpa.3160220202](https://doi.org/10.1002/cpa.3160220202).
- [12] L. Dai and B. Xia. “Non-Termination Sets of Simple Linear Loops”. In: *Proc. ICTAC 12*. LNCS 7521. 2012, pp. 61–73. DOI: [10.1007/978-3-642-32943-2\\_5](https://doi.org/10.1007/978-3-642-32943-2_5).
- [13] A. van den Essen and E. Hubbers. “Polynomial Maps with Strongly Nilpotent Jacobian Matrix and the Jacobian Conjecture”. In: *Linear Algebra and its Applications* 247 (1996), pp. 121–132. DOI: [10.1016/0024-3795\(95\)00095-X](https://doi.org/10.1016/0024-3795(95)00095-X).
- [14] A. van den Essen. *Polynomial Automorphisms and the Jacobian Conjecture*. Springer Basel, 2000. DOI: [10.1007/978-3-0348-8440-2](https://doi.org/10.1007/978-3-0348-8440-2).
- [15] F. Frohn and J. Giesl. “Termination of Triangular Integer Loops is Decidable”. In: *Proc. CAV '19*. LNCS 11562. 2019, pp. 269–286. DOI: [10.1007/978-3-030-25543-5\\_24](https://doi.org/10.1007/978-3-030-25543-5_24).
- [16] F. Frohn and J. Giesl. “Proving Non-Termination via Loop Acceleration”. In: *Proc. FMCAD '19*. 2019, pp. 221–230. DOI: [10.23919/FMCAD.2019.8894271](https://doi.org/10.23919/FMCAD.2019.8894271).
- [17] F. Frohn. “A Calculus for Modular Loop Acceleration”. In: *Proc. TACAS '20*. LNCS 12078. 2020, pp. 58–76. DOI: [10.1007/978-3-030-45190-5\\_4](https://doi.org/10.1007/978-3-030-45190-5_4).
- [18] M. Giesbrecht. “Nearly Optimal Algorithms for Canonical Matrix Forms”. In: *SIAM Journal on Computing* 24.5 (1995), pp. 948–969. DOI: [10.1137/S0097539793252687](https://doi.org/10.1137/S0097539793252687).
- [19] J. Giesl, C. Aschermann, M. Brockschmidt, F. Emmes, F. Frohn, C. Fuhs, J. Hensel, C. Otto, M. Plücker, P. Schneider-Kamp, T. Ströder, S. Swiderski, and R. Thiemann. “Analyzing Program Termination and Complexity Automatically with AProVE”. In: *Journal of Automated Reasoning* 58.1 (2017), pp. 3–31. DOI: [10.1007/s10817-016-9388-y](https://doi.org/10.1007/s10817-016-9388-y).
- [20] J. Giesl, A. Rubio, C. Sternagel, J. Waldmann, and A. Yamada. “The Termination and Complexity Competition”. In: *Proc. TACAS '19*. LNCS 11429. 2019, pp. 156–166. DOI: [10.1007/978-3-030-17502-3\\_10](https://doi.org/10.1007/978-3-030-17502-3_10).
- [21] A. Gupta, T. A. Henzinger, R. Majumdar, A. Rybalchenko, and R. Xu. “Proving Non-Termination”. In: *Proc. POPL '08*. 2008, pp. 147–158. DOI: [10.1145/1328438.1328459](https://doi.org/10.1145/1328438.1328459).
- [22] M. Hark, F. Frohn, and J. Giesl. “Polynomial Loops: Beyond Termination”. In: *Proc. LPAR 20*. EPiC 73. 2020, pp. 279–297. DOI: [10.29007/nxv1](https://doi.org/10.29007/nxv1).
- [23] M. Hosseini, J. Ouaknine, and J. Worrell. “Termination of Linear Loops over the Integers”. In: *Proc. ICALP '19*. LIPIcs 132. 2019, 118:1–118:13. DOI: [10.4230/LIPIcs.ICALP.2019.118](https://doi.org/10.4230/LIPIcs.ICALP.2019.118).

- [24] A. Humenberger, M. Jaroschek, and L. Kovács. “Invariant Generation for Multi-Path Loops with Polynomial Assignments”. In: *Proc. VMCAI '18*. LNCS 10747. 2018, pp. 226–246. DOI: [10.1007/978-3-319-73721-8\\_11](https://doi.org/10.1007/978-3-319-73721-8_11).
- [25] M. Kauers and P. Paule. *The Concrete Tetrahedron – Symbolic Sums, Recurrence Equations, Generating Functions, Asymptotic Estimates*. Springer, 2011. DOI: [10.1007/978-3-7091-0445-3](https://doi.org/10.1007/978-3-7091-0445-3).
- [26] Z. Kincaid, J. Cyphert, J. Breck, and T. W. Reps. “Non-Linear Reasoning for Invariant Synthesis”. In: *Proceedings of the ACM on Programming Languages* 2.POPL (2018), 54:1–54:33. DOI: [10.1145/3158142](https://doi.org/10.1145/3158142).
- [27] Z. Kincaid, J. Breck, J. Cyphert, and T. W. Reps. “Closed Forms for Numerical Loops”. In: *Proceedings of the ACM on Programming Languages* 3.POPL (2019), 55:1–55:29. DOI: [10.1145/3290368](https://doi.org/10.1145/3290368).
- [28] L. Kovács. “Reasoning Algebraically About P-Solvable Loops”. In: *Proc. TACAS '08*. LNCS 4963. 2008, pp. 249–264. DOI: [10.1007/978-3-540-78800-3\\_18](https://doi.org/10.1007/978-3-540-78800-3_18).
- [29] D. Larraz, A. Oliveras, E. Rodríguez-Carbonell, and A. Rubio. “Proving Termination of Imperative Programs using Max-SMT”. In: *Proc. FMCAD '13*. 2013, pp. 218–225. DOI: [10.1109/FMCAD.2013.6679413](https://doi.org/10.1109/FMCAD.2013.6679413).
- [30] D. Larraz, K. Nimkar, A. Oliveras, E. Rodríguez-Carbonell, and A. Rubio. “Proving Non-Termination Using Max-SMT”. In: *Proc. CAV '14*. LNCS 8559. 2014, pp. 779–796. DOI: [10.1007/978-3-319-08867-9\\_52](https://doi.org/10.1007/978-3-319-08867-9_52).
- [31] J. Leike and M. Heizmann. “Ranking Templates for Linear Loops”. In: *Logical Methods in Computer Science* 11.1 (2015). DOI: [10.2168/LMCS-11\(1:16\)2015](https://doi.org/10.2168/LMCS-11(1:16)2015).
- [32] J. Leike and M. Heizmann. “Geometric Nontermination Arguments”. In: *Proc. TACAS '18*. LNCS 10806. 2018, pp. 266–283. DOI: [10.1007/978-3-319-89963-3\\_16](https://doi.org/10.1007/978-3-319-89963-3_16).
- [33] Y. Li. “A Recursive Decision Method for Termination of Linear Programs”. In: *Proc. SNC '14*. 2014, pp. 97–106. DOI: [10.1145/2631948.2631966](https://doi.org/10.1145/2631948.2631966).
- [34] Y. Li. “Termination of Single-Path Polynomial Loop Programs”. In: *Proc. ICTAC '16*. LNCS 9965. 2016, pp. 33–50. DOI: [10.1007/978-3-319-46750-4\\_3](https://doi.org/10.1007/978-3-319-46750-4_3).
- [35] Y. Li. “Termination of Semi-algebraic Loop Programs”. In: *Proc. SETTA '17*. LNCS 10606. 2017, pp. 131–146. DOI: [10.1007/978-3-319-69483-2\\_8](https://doi.org/10.1007/978-3-319-69483-2_8).
- [36] Y. Li. “Witness to Non-Termination of Linear Programs”. In: *Theoretical Computer Science* 681 (2017), pp. 75–100. DOI: [10.1016/j.tcs.2017.03.036](https://doi.org/10.1016/j.tcs.2017.03.036).
- [37] E. Neumann, J. Ouaknine, and J. Worrell. “On Ranking Function Synthesis and Termination for Polynomial Programs”. In: *Proc. CONCUR '20*. LIPIcs 171. 2020, 15:1–15:15. DOI: [10.4230/LIPIcs.CONCUR.2020.15](https://doi.org/10.4230/LIPIcs.CONCUR.2020.15).
- [38] S. de Oliveira, S. Bensalem, and V. Prevosto. “Polynomial Invariants by Linear Algebra”. In: *Proc. ATVA '16*. LNCS 9938. 2016, pp. 479–494. DOI: [10.1007/978-3-319-46520-3\\_30](https://doi.org/10.1007/978-3-319-46520-3_30).
- [39] S. de Oliveira, S. Bensalem, and V. Prevosto. “Synthesizing Invariants by Solving Solvable Loops”. In: *Proc. ATVA '17*. LNCS 10482. 2017, pp. 327–343. DOI: [10.1007/978-3-319-68167-2\\_22](https://doi.org/10.1007/978-3-319-68167-2_22).
- [40] J. Ouaknine, J. S. Pinto, and J. Worrell. “On Termination of Integer Linear Loops”. In: *Proc. SODA '15*. 2015, pp. 957–969. DOI: [10.1137/1.9781611973730.65](https://doi.org/10.1137/1.9781611973730.65).

- [41] A. D. Pia, S. S. Dey, and M. Molinaro. “Mixed-Integer Quadratic Programming is in NP”. In: *Mathematical Programming* 162.1-2 (2017), pp. 225–240. DOI: [10.1007/s10107-016-1036-0](https://doi.org/10.1007/s10107-016-1036-0).
- [42] A. Podelski and A. Rybalchenko. “A Complete Method for the Synthesis of Linear Ranking Functions”. In: *Proc. VMCAI '04*. LNCS 2937. 2004, pp. 239–251. DOI: [10.1007/978-3-540-24622-0\\_20](https://doi.org/10.1007/978-3-540-24622-0_20).
- [43] J. Renegar. “On the Computational Complexity and Geometry of the First-Order Theory of the Reals, Part I: Introduction. Preliminaries. The Geometry of Semi-Algebraic Sets. The Decision Problem for the Existential Theory of the Reals.” In: *Journal of Symbolic Computation* 13.3 (1992), pp. 255–300. DOI: [10.1016/S0747-7171\(10\)80003-3](https://doi.org/10.1016/S0747-7171(10)80003-3).
- [44] J. Robinson. “Definability and Decision Problems in Arithmetic”. In: *Journal of Symbolic Logic* 14.2 (1949), pp. 98–114. DOI: [10.2307/2266510](https://doi.org/10.2307/2266510).
- [45] J.-L. Roch and G. Villard. “Fast Parallel Computation of the Jordan Normal Form of Matrices”. In: *Parallel Processing Letters* 06.02 (1996), pp. 203–212. DOI: [10.1142/S0129626496000200](https://doi.org/10.1142/S0129626496000200).
- [46] D. S. Roche. “What Can (and Can’t) we Do with Sparse Polynomials?” In: *Proc. ISSAC '18*. 2018, pp. 25–30. DOI: [10.1145/3208976.3209027](https://doi.org/10.1145/3208976.3209027).
- [47] E. Rodríguez-Carbonell and D. Kapur. “Automatic Generation of Polynomial Loop Invariants: Algebraic Foundation”. In: *Proc. ISSAC '04*. 2004, pp. 266–273. DOI: [10.1145/1005285.1005324](https://doi.org/10.1145/1005285.1005324).
- [48] E. Rodríguez-Carbonell and D. Kapur. “Generating all Polynomial Invariants in Simple Loops”. In: *Journal of Symbolic Computation* 42.4 (2007), pp. 443–476. DOI: [10.1016/j.jsc.2007.01.002](https://doi.org/10.1016/j.jsc.2007.01.002).
- [49] M. Schaefer. “Complexity of Some Geometric and Topological Problems”. In: *Proc. GD '09*. LNCS 5849. 2009, pp. 334–344. DOI: [10.1007/978-3-642-11805-0\\_32](https://doi.org/10.1007/978-3-642-11805-0_32).
- [50] M. Schaefer and D. Štefankovič. “Fixed Points, Nash Equilibria, and the Existential Theory of the Reals”. In: *Theory of Computing Systems* 60.2 (2017), pp. 172–193. DOI: [10.1007/s00224-015-9662-0](https://doi.org/10.1007/s00224-015-9662-0).
- [51] A. Tarski. “A Decision Method for Elementary Algebra and Geometry”. In: *Quantifier Elimination and Cylindrical Algebraic Decomposition*. Ed. by B. F. Caviness and J. R. Johnson. Originally appeared in 1951, University of California Press, Berkeley and Los Angeles. Springer Vienna, 1998, pp. 24–84.
- [52] A. Tiwari. “Termination of Linear Programs”. In: *Proc. CAV '04*. LNCS 3114. 2004, pp. 70–82. DOI: [10.1007/978-3-540-27813-9\\_6](https://doi.org/10.1007/978-3-540-27813-9_6).
- [53] *TPDB (Termination Problems Data Base)*. URL: <http://termination-portal.org/wiki/TPDB>.
- [54] B. Xia and Z. Zhang. “Termination of Linear Programs with Nonlinear Constraints”. In: *Journal of Symbolic Computation* 45.11 (2010), pp. 1234–1249. DOI: [10.1016/j.jsc.2010.06.006](https://doi.org/10.1016/j.jsc.2010.06.006).
- [55] M. Xu and Z. Li. “Symbolic Termination Analysis of Solvable Loops”. In: *Journal of Symbolic Computation* 50 (2013), pp. 28–49. DOI: [10.1016/j.jsc.2012.05.005](https://doi.org/10.1016/j.jsc.2012.05.005).

# Appendix

## A Proofs

### A.1 Proof of Lemma 7

*Proof.* Let  $(\varphi, \vec{a})$  be a loop. Since  $id_{\mathcal{S}[\vec{x}]}^{-1} = id_{\mathcal{S}[\vec{x}]}$ , we obtain  $Tr_{id_{\mathcal{S}[\vec{x}]}}(\varphi, \vec{a}) = (\varphi', \vec{a}')$  with

$$\begin{aligned}\varphi' &= id_{\mathcal{S}[\vec{x}]}^{-1}(\varphi) = \varphi \\ \vec{a}' &= (id_{\mathcal{S}[\vec{x}]}^{-1} \circ \tilde{a} \circ id_{\mathcal{S}[\vec{x}]})(\vec{x}) = \tilde{a}(\vec{x}) = \vec{a}\end{aligned}$$

Now we take  $\eta_1, \eta_2 \in \text{Aut}_{\mathcal{S}}(\mathcal{S}[\vec{x}])$ . Note that  $(\eta_1 \circ \eta_2)^{-1} = \eta_2^{-1} \circ \eta_1^{-1}$ . Let  $Tr_{\eta_1 \circ \eta_2}(\varphi, \vec{a}) = (\varphi', \vec{a}')$ ,  $Tr_{\eta_1}(\varphi, \vec{a}) = (\varphi'', \vec{a}'')$ , and  $Tr_{\eta_2}(\varphi'', \vec{a}'') = (\varphi''', \vec{a}''')$ . We have

$$\begin{aligned}\varphi' &= (\eta_2^{-1} \circ \eta_1^{-1})(\varphi) \\ \varphi'' &= \eta_1^{-1}(\varphi) \\ \varphi''' &= \eta_2^{-1}(\varphi'') \\ &= \eta_2^{-1}(\eta_1^{-1}(\varphi)) \\ &= (\eta_2^{-1} \circ \eta_1^{-1})(\varphi) \\ &= \varphi'\end{aligned}$$

Moreover, we have

$$\begin{aligned}\vec{a}' &= (\eta_2^{-1} \circ \eta_1^{-1} \circ \tilde{a} \circ \eta_1 \circ \eta_2)(\vec{x}) \\ &= (\eta_2(\vec{x})) (\eta_1(\vec{x})) (\vec{a}) (\eta_1^{-1}(\vec{x})) (\eta_2^{-1}(\vec{x})) \\ \vec{a}'' &= (\eta_1^{-1} \circ \tilde{a} \circ \eta_1)(\vec{x}) \\ &= (\eta_1(\vec{x})) (\vec{a}) (\eta_1^{-1}(\vec{x})) \\ \vec{a}''' &= (\eta_2^{-1} \circ \tilde{a}'' \circ \eta_2)(\vec{x}) \\ &= (\eta_2(\vec{x})) (\eta_1(\vec{x})) (\vec{a}) (\eta_1^{-1}(\vec{x})) (\eta_2^{-1}(\vec{x})) \\ &= \vec{a}'\end{aligned}$$

□

### A.2 Proof of Lemma 8

*Proof.* Let  $\vec{c}$  be a witness for non-termination of  $(\varphi, \vec{a})$ , i.e.,  $\varphi(\vec{a}^n(\vec{c}))$  holds for all  $n \in \mathbb{N}$ . Let  $Tr_{\eta}(\varphi, \vec{a}) = (\varphi', \vec{a}')$ . To prove the lemma, we show that

$$\varphi'(\vec{a}'^n((\eta(\vec{x}))(\vec{c}))) = \varphi(\vec{a}^n(\vec{c}))$$

for all  $n \in \mathbb{N}$ . We have



$$\begin{aligned}
& \varphi'(\vec{a}'^n((\eta(\vec{x}))(\vec{c}))) \\
&= \eta^{-1}(\varphi)(\vec{a}'^n((\eta(\vec{x}))(\vec{c}))) \\
&= \varphi[\vec{x}/\eta^{-1}(\vec{x})] \underbrace{[\vec{x}/\vec{a}']}_{n \text{ times}} [\vec{x}/\eta(\vec{x})] [\vec{x}/\vec{c}] \\
&= \varphi[\vec{x}/\eta^{-1}(\vec{x})] [\vec{x}/\eta(\vec{x})] \underbrace{[\vec{x}/\vec{a}']}_{n \text{ times}} [\vec{x}/\eta^{-1}(\vec{x})] [\vec{x}/\eta(\vec{x})] [\vec{x}/\vec{c}] \\
&= \varphi \underbrace{[\vec{x}/\vec{a}']}_{n \text{ times}} [\vec{x}/\vec{c}] \\
&= \varphi(\vec{a}'^n(\vec{c}))
\end{aligned}$$

□

### A.3 Proof of Thm. 10

*Proof.* The second statement implies the first. In Lemma 8 we have seen that if  $\vec{c}$  is a witness for non-termination of  $(\varphi, \vec{a})$ , then  $\widehat{\eta}(\vec{c})$  witnesses non-termination of  $Tr_\eta(\varphi, \vec{a})$ . Now let  $\vec{u}$  be a witness for non-termination of  $Tr_\eta(\varphi, \vec{a})$ . Then by Lemma 8,  $\widehat{\eta^{-1}}(\vec{u})$  witnesses non-termination of  $Tr_{\eta^{-1}}(Tr_\eta(\varphi, \vec{a})) \stackrel{\text{Lemma 7}}{=} Tr_{\eta \circ \eta^{-1}}(\varphi, \vec{a}) = (\varphi, \vec{a})$ . Hence,  $\widehat{\eta}$  maps witnesses for non-termination of  $(\varphi, \vec{a})$  to witnesses for non-termination of  $Tr_\eta(\varphi, \vec{a})$  and  $\widehat{\eta^{-1}}$  maps witnesses for non-termination of  $Tr_\eta(\varphi, \vec{a})$  to witnesses for non-termination of  $(\varphi, \vec{a})$ . These two mappings are inverse to each other: For  $\vec{u} \in \mathcal{S}^d$  we have

$$\begin{aligned}
& \widehat{\eta}(\widehat{\eta^{-1}}(\vec{u})) \\
&= \widehat{\eta}((\eta^{-1}(\vec{x}))(\vec{u})) && \text{by definition of } \widehat{\eta^{-1}} \\
&= (\eta(\vec{x}))((\eta^{-1}(\vec{x}))(\vec{u})) && \text{by definition of } \widehat{\eta} \\
&= \eta(\vec{x})[\vec{x}/\eta^{-1}(\vec{x})][\vec{x}/\vec{u}] \\
&= \vec{u}
\end{aligned}$$

$$\begin{aligned}
& \widehat{\eta^{-1}}(\widehat{\eta}(\vec{u})) \\
&= \widehat{\eta^{-1}}((\eta(\vec{x}))(\vec{u})) && \text{by definition of } \widehat{\eta} \\
&= (\eta^{-1}(\vec{x}))((\eta(\vec{x}))(\vec{u})) && \text{by definition of } \widehat{\eta^{-1}} \\
&= \eta^{-1}(\vec{x})[\vec{x}/\eta(\vec{x})][\vec{x}/\vec{u}] \\
&= \vec{u}.
\end{aligned}$$

Hence,  $\widehat{\eta}$  is indeed a bijection with inverse mapping  $\widehat{\eta^{-1}}$ . □

### A.4 Proof of Lemma 12

*Proof.* Let  $\vec{c}' \in \widehat{\eta}(F)$ . Then  $\vec{c}' = \widehat{\eta}(\vec{c})$  for some  $\vec{c} \in F$ . As  $F$  is  $\vec{a}$ -invariant, we have  $\vec{a}(\vec{c}) \in F$ . We obtain

$$\begin{aligned}
\vec{a}'(\vec{c}') &= (\eta(\vec{x}))(\vec{a})(\eta^{-1}(x))(\vec{c}') \\
&= (\eta(\vec{x}))(\vec{a})(\eta^{-1}(x))(\eta(\vec{x}))(\vec{c}) \\
&= (\eta(\vec{x}))(\vec{a})(\vec{c}) \\
&= \widehat{\eta}(\vec{a}(\vec{c})) \in \widehat{\eta}(F).
\end{aligned}$$

□

### A.5 Proof of Lemma 14

*Proof.* Let  $F$  be characterized by  $\psi_F \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ . Consider the following formula  $\psi \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ :

$$\exists \vec{y} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F(\vec{y}) \wedge \vec{x} = (\eta(\vec{x}))(\vec{y})$$

Then  $\psi(\vec{c})$  holds for a point  $\vec{c} \in \mathbb{R}_{\mathbb{A}}^d$  iff  $\vec{c} = \widehat{\eta}(\vec{u})$  for some  $\vec{u} \in \mathbb{R}_{\mathbb{A}}^d$  where  $\psi_F(\vec{u})$  holds, i.e., where  $\vec{u} \in F$ . □

### A.6 Proof of Thm. 16

*Proof.* As  $(\varphi, \vec{a})$  is solvable, there is a partitioning  $\mathcal{J} = \{J_1, \dots, J_k\}$  as in Def. 2, i.e.,  $\{1, \dots, d\} = \bigsqcup_{i=1}^k J_i$  and for all  $1 \leq i \leq k$  we have  $\vec{a}_{J_i} = A_i \cdot \vec{x}_{J_i} + \vec{p}_i$ , where  $\vec{p}_i \in (\mathcal{S}[\vec{x}_{J_{i+1}}, \dots, \vec{x}_{J_k}])^{d_i}$ . W.l.o.g., assume that  $\vec{x}$  is ordered according to  $\mathcal{J}$ , i.e., if  $x_{i_1} \in J_{j_1}$  and  $x_{i_2} \in J_{j_2}$  for  $j_1 < j_2$ , then  $i_1 < i_2$ .

For each  $A_i$ , let  $Q_i = T_i \cdot A_i \cdot T_i^{-1}$  be its Jordan normal form, where  $T_i$  is the corresponding transformation matrix. Since  $A_i$  has only real eigenvectors, this means that the entries of  $Q_i$ ,  $T_i$ , and  $T_i^{-1}$  are real algebraic numbers. Let  $\eta$  be the endomorphism defined by  $\eta(\vec{x}_{J_i}) = T_i \cdot \vec{x}_{J_i}$ . This means that  $\eta$  is induced by the block diagonal matrix  $\text{Diag}(T_1, T_2, \dots, T_k)$ . Then  $\eta$  is an automorphism and its inverse satisfies  $\eta^{-1}(\vec{x}_{J_i}) = T_i^{-1} \cdot \vec{x}_{J_i}$ . Furthermore, the degree of  $\eta$  is obviously 1. Moreover,  $\eta$  and  $\eta^{-1}$  are compatible with the partition, i.e., the images of the variables in  $\vec{x}_{J_i}$  under  $\eta$  and  $\eta^{-1}$  are polynomials only using the variables  $\vec{x}_{J_i}$ . For each  $1 \leq i \leq k$  we have:

$$\begin{aligned}
&(\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x}_{J_i}) \\
&= \eta(\vec{x}_{J_i})[\vec{x}/\vec{a}][\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot \vec{x}_{J_i})[\vec{x}/\vec{a}][\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot \vec{a}_{J_i})[\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot (A_i \cdot \vec{x}_{J_i} + \vec{p}_i))[\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot A_i \cdot \vec{x}_{J_i} + T_i \cdot \vec{p}_i)[\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot A_i \cdot \vec{x}_{J_i})[\vec{x}/\eta^{-1}(\vec{x})] + (T_i \cdot \vec{p}_i)[\vec{x}/\eta^{-1}(\vec{x})] \\
&= (T_i \cdot A_i \cdot T_i^{-1} \cdot \vec{x}_{J_i}) + (T_i \cdot \vec{p}_i)[\vec{x}/\eta^{-1}(\vec{x})] \\
&= (Q_i \cdot \vec{x}_{J_i}) + (T_i \cdot \vec{p}_i)[\vec{x}/\eta^{-1}(\vec{x})]
\end{aligned}$$

We have  $T_i \cdot \vec{p}_i \in \mathcal{S}[\vec{x}_{J_{i+1}}, \dots, \vec{x}_{J_k}]^{d_i}$ . Therefore, we have  $(T_i \cdot \vec{p}_i) [\vec{x}/\eta^{-1}(\vec{x})] \in \mathcal{S}[\vec{x}_{J_{i+1}}, \dots, \vec{x}_{J_k}]^{d_i}$  as well, as  $\eta^{-1}$  is compatible with the partitioning. This implies that  $Tr_\eta(\varphi, \vec{a})$  is weakly non-linear. Since we assumed that  $\vec{x}$  is ordered w.r.t. the partitioning and each  $Q_i$  is triangular,  $Tr_\eta(\varphi, \vec{a})$  is triangular, too. Thus,  $Tr_\eta(\varphi, \vec{a})$  is in *tnw*-form.  $\square$

### A.7 Proof of Lemma 19

*Proof.* Let  $\delta = \deg(\eta)$ . Note that for any  $e \in \mathbb{N}$ , there is only a finite number of monomials over  $\vec{x}$  of degree  $e$ . (More precisely, the number of monomials of exactly degree  $e$  is  $\binom{d+e-1}{e}$ .) Hence, for any  $1 \leq i \leq d$  we can construct the following term that stands for  $\eta^{-1}(x_i)$ :

$$\sum_{m \text{ is a monomial of (at most) degree } \delta^{d-1}} a_{i,m} \cdot m$$

Here, the monomials  $m$  contain the variables  $\vec{x}$  and the  $a_{i,m}$  are variables that stand for the unknown coefficients of the polynomial  $\eta^{-1}(x_i)$ .

Hence, for any  $1 \leq i \leq d$  we now build a formula  $\varphi_{r,i}$  which stands for the requirement “ $\eta(\eta^{-1}(x_i)) = x_i$ ” (i.e., that  $\eta^{-1}$  is a right inverse of  $\eta$ ):

$$\varphi_{r,i} : \sum_{m \text{ is a monomial of (at most) degree } \delta^{d-1}} a_{i,m} \cdot \eta(m) = x_i$$

Similarly, for any  $1 \leq i \leq d$  we construct a formula  $\varphi_{l,i}$  which stands for the requirement “ $\eta^{-1}(\eta(x_i)) = x_i$ ” (i.e., that  $\eta^{-1}$  is a left inverse of  $\eta$ ):

$$\varphi_{l,i} : \eta(x_i) \left( \sum_{m \text{ is a monomial of (at most) degree } \delta^{d-1}} a_{i,m} \cdot m \right) = x_i$$

Thus, the formula

$$\forall \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \bigwedge_{i=1}^d \varphi_{r,i} \wedge \bigwedge_{i=1}^d \varphi_{l,i} \tag{16}$$

is valid iff  $\eta$  has an inverse of degree at most  $\delta^{d-1}$ . By [Thm. 18](#), this is equivalent to the question whether  $\eta$  has an inverse, i.e., whether  $\eta$  is an automorphism. Unfortunately, (16)  $\notin \text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ . However,  $\bigwedge_{i=1}^d \varphi_{r,i} \wedge \bigwedge_{i=1}^d \varphi_{l,i}$  has to hold for all  $\vec{x} \in \mathbb{R}_{\mathbb{A}}^d$ . So, we can reduce this formula to a system of equations: one simply has to check whether there is an instantiation of the unknown coefficients  $a_{i,m}$  such that all monomials in  $\varphi_{r,i}$  and  $\varphi_{l,i}$  except  $x_i$  get the coefficient 0 and the monomial  $x_i$  gets the coefficient 1. When building the conjunction of these equations and existentially quantifying the unknown coefficients  $a_{i,m}$ , one indeed obtains a formula from  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ .  $\square$

### A.8 Proof of Thm. 20

*Proof.* For every  $1 \leq i \leq d$ , let

$$\eta(x_i) = \sum_{m \text{ is a monomial of (at most) degree } \delta} b_{i,m} \cdot m,$$

where the  $b_{i,m}$  are variables that stand for unknown coefficients. By [Lemma 19](#) there is a  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ -formula that contains both  $b_{i,m}$  and the variables  $a_{i,m}$  (for the coefficients of  $\eta^{-1}$ ) which expresses that  $\eta$  is an automorphism.

Furthermore, using these coefficients we can construct a formula from  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$  which expresses that the update  $\vec{a}' = (a'_1, \dots, a'_d) = (\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x})$  is a *twn*-loop: Note that we have  $\deg(\vec{a}') = \deg((\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x})) \leq \deg(\eta^{-1}) \cdot \deg(\tilde{a}) \cdot \deg(\eta) \leq \delta^{d-1} \cdot \deg(\tilde{a}) \cdot \delta$ . So there is a bound on the degree of the polynomials occurring in the transformed loop  $\text{Tr}_{\eta}(\varphi, \vec{a})$ . Hence, for every  $1 \leq i \leq d$ , let

$$a'_i = \sum_{m \text{ is a monomial of (at most) degree } \delta^{d-1} \cdot \deg(\tilde{a}) \cdot \delta} c_{i,m} \cdot m,$$

where the variables  $c_{i,m}$  stand again for unknown coefficients. Now we can build a  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ -formula which is valid iff  $\vec{a}'$  is in *twn*-form by requiring that certain coefficients  $c_{i,m}$  are zero. Moreover, we can construct a  $\text{Th}_{\exists}(\mathbb{R}_{\mathbb{A}})$ -formula which is valid iff  $\vec{a}' = (\eta^{-1} \circ \tilde{a} \circ \eta)(\vec{x})$ . This proves the theorem.  $\square$

## A.9 Proof of [Thm. 26](#)

*Proof.* We first prove:

$$\text{If } (\varphi, \vec{a}) \text{ is } \textit{twn}, \text{ then } (\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a})) \text{ is } \textit{tnn}. \quad (17)$$

Due to weak non-linearity, we have  $a_i = m_i \cdot x_i + \alpha_i$  with  $x_i \notin \mathcal{V}(\alpha_i)$  for all  $1 \leq i \leq d$ . Then

$$a_i(\vec{a}) = m_i \cdot (m_i \cdot x_i + \alpha_i) + \alpha_i(\vec{a}) = m_i^2 \cdot x_i + m_i \cdot \alpha_i + \alpha_i(\vec{a}).$$

Assume that  $x_i \in \mathcal{V}(\alpha_i(\vec{a}))$ . Since  $x_i \notin \mathcal{V}(\alpha_i)$  by weak non-linearity, there must be an  $x_j \in \mathcal{V}(\alpha_i)$  with  $x_j \neq x_i$  and  $x_i \in \mathcal{V}(\alpha_j)$ . By definition,  $x_j \neq x_i$  and  $x_i \in \mathcal{V}(\alpha_j)$  implies  $x_j \succ_{\vec{a}} x_i$ . But  $x_j \in \mathcal{V}(\alpha_i)$  also implies  $x_i \succ_{\vec{a}} x_j$ , which would violate well-foundedness of  $\succ_{\vec{a}}$ , i.e., it would contradict the triangularity of  $(\varphi, \vec{a})$ .

Hence,  $m_i^2$  is the coefficient of  $x_i$  in  $a_i(\vec{a})$ . Since  $m_i^2 \geq 0$ , this proves that  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$  is non-negative.

Note that  $x_i \succ_{\vec{a}(\vec{a})} x_j$  implies  $x_j \in \mathcal{V}(\alpha_i)$  (in this case we also have  $x_i \succ_{\vec{a}} x_j$ ) or it implies that there is an  $x_k \in \mathcal{V}(\alpha_i)$  with  $x_j \in \mathcal{V}(\alpha_k)$  (in this case we have  $x_i \succ_{\vec{a}} x_k$  and  $x_k \succeq_{\vec{a}} x_j$ ). So in both cases,  $x_i \succ_{\vec{a}(\vec{a})} x_j$  implies  $x_i \succ_{\vec{a}} x_j$ . Thus, we obtain  $\succ_{\vec{a}(\vec{a})} \subseteq \succ_{\vec{a}}$ . As  $\succ_{\vec{a}}$  is well founded, this means that  $\succ_{\vec{a}(\vec{a})}$  is well founded, too. Hence,  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$  is triangular.

Now we prove that  $(\varphi, \vec{a})$  terminates iff  $(\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a}))$  terminates. Then the claim immediately follows due to [\(17\)](#), as chaining is clearly computable.

$$\begin{aligned} & (\varphi, \vec{a}) \text{ does not terminate} \\ \iff & \exists \vec{c} \in F. \forall n \in \mathbb{N}. \varphi(\vec{a}^n(\vec{c})) \quad \text{by } \text{Def. 1} \\ \iff & \exists \vec{c} \in F. \forall n \in \mathbb{N}. \varphi(\vec{a}^{2 \cdot n}(\vec{c})) \wedge \varphi(\vec{a}^{2 \cdot n + 1}(\vec{c})) \\ \iff & \exists \vec{c} \in F. \forall n \in \mathbb{N}. \varphi(\vec{a}^{2 \cdot n}(\vec{c})) \wedge \varphi(\vec{a})(\vec{a}^{2 \cdot n}(\vec{c})) \\ \iff & \exists \vec{c} \in F. \forall n \in \mathbb{N}. (\varphi \wedge \varphi(\vec{a})) (\vec{a}(\vec{a}))^n \\ \iff & (\varphi \wedge \varphi(\vec{a}), \vec{a}(\vec{a})) \text{ does not terminate} \end{aligned}$$

□

### A.10 Proof of Lemma 31

*Proof.* Recall that for  $f, g : \mathbb{N} \rightarrow \mathbb{R}$ ,  $f(n) \in o(g(n))$  means

$$\forall m > 0. \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. |f(n)| < m \cdot |g(n)|.$$

First consider the case  $b_2 > b_1$ . We have  $b_2^n = b_1^n \cdot \left(\frac{b_2}{b_1}\right)^n$  where  $\frac{b_2}{b_1} > 1$ . As we clearly have  $n^{a_1} \in o\left(\left(\frac{b_2}{b_1}\right)^n\right)$ , we obtain  $n^{a_1} \cdot b_1^n \in o\left(\left(\frac{b_2}{b_1}\right)^n \cdot b_1^n\right) = o(b_2^n) \subseteq o(n^{a_2} \cdot b_2^n)$ , i.e.,  $n^{a_1} \cdot b_1^n \in o(n^{a_2} \cdot b_2^n)$ .

Now consider the case  $b_2 = b_1$  and  $a_2 > a_1$ . Then  $n^{a_1} \cdot b_1^n \in o(n^{a_2} \cdot b_2^n)$  trivially holds. □

### A.11 Proof of Equation (9)

*Proof.* If  $p(\vec{c}) = 0$ , then  $k = 0$  by Def. 32 and hence  $o(p(\vec{c})) = o(k \cdot n^a \cdot b^n) = o(0)$ . Otherwise,  $p(\vec{c})$  has the form

$$k \cdot n^a \cdot b^n + \sum_{i=1}^{\ell} k_i \cdot n^{a_i} \cdot b_i^n$$

for  $k \neq 0$  and  $\ell \geq 0$ . We have  $k_i^{(b_i, a_i)} \in \text{coefs}(p(\vec{c}))$  and hence  $(b, a) >_{lex} (b_i, a_i)$  for all  $1 \leq i \leq \ell$ . Thus, Lemma 31 implies  $n^{a_i} \cdot b_i^n \in o(n^a \cdot b^n)$  and hence we get

$$o(p(\vec{c})) = o\left(k \cdot n^a \cdot b^n + \sum_{i=1}^{\ell} k_i \cdot n^{a_i} \cdot b_i^n\right) = o(n^a \cdot b^n) = o(k \cdot n^a \cdot b^n).$$

□

### A.12 Proof of Equation (10)

*Proof.* If  $k = 0$ , then the claim is trivial, so assume  $k \neq 0$ , i.e.,  $p(\vec{c}) = k \cdot b^n \cdot n^a + p'$  for some  $p' \in \mathbb{NPE}$ . By Lemma 31 we have

$$p' \in o(k \cdot b^n \cdot n^a)$$

$$\iff \forall m > 0. \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. |p'| < m \cdot |k \cdot b^n \cdot n^a|$$

$$\implies \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. |p'| < |k \cdot b^n \cdot n^a|.$$

Assume  $k > 0$ . Then

$$\exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. |p'| < |k \cdot b^n \cdot n^a|$$

$$\implies \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. -p' < |k \cdot b^n \cdot n^a|$$

$$\iff \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. -p' < k \cdot b^n \cdot n^a$$

$$\iff \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. 0 < k \cdot b^n \cdot n^a + p'$$

$$\iff \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. 0 < p(\vec{c})$$

$$\iff \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \text{sign}(p(\vec{c})) = \text{sign}(k).$$

If  $k < 0$ , then

$$\begin{aligned}
& \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. |p'| < |k \cdot b^n \cdot n^a| \\
\implies & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p' < |k \cdot b^n \cdot n^a| \\
\iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p' < -k \cdot b^n \cdot n^a \\
\iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. k \cdot b^n \cdot n^a + p' < 0 \\
\iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p(\vec{c}) < 0 \\
\iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \text{sign}(p(\vec{c})) = \text{sign}(k).
\end{aligned}$$

□

### A.13 Proof of Lemma 34

*Proof.* By the definition of  $\vec{q}_{norm}$ , we have  $p \in \text{NPE}[\vec{x}]$  and thus  $p(\vec{c}) \in \text{NPE}$  for any  $\vec{c} \in \mathbb{R}_{\mathbb{A}}^d$ . Hence,

$$\exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. p(\vec{c}) \triangleright 0 \quad \text{iff} \quad \text{unmark}(\max_{>coef}(\text{coefs}(p(\vec{c})))) \triangleright 0 \quad (\text{by (10)}).$$

Let  $\text{coefs}(p) = \{\alpha_1^{(b_1, a_1)}, \dots, \alpha_\ell^{(b_\ell, a_\ell)}\}$  where  $\alpha_i^{(b_i, a_i)} \succ_{coef} \alpha_j^{(b_j, a_j)}$  for all  $1 \leq i < j \leq \ell$ . If  $p(\vec{c}) = 0$ , then  $\alpha_1(\vec{c}) = \dots = \alpha_\ell(\vec{c}) = 0$  and thus  $\text{coefs}(p(\vec{c})) = \{0^{(1,0)}\}$  and  $\text{unmark}(\max_{>coef}(\text{coefs}(p(\vec{c})))) = 0$ . Otherwise, there is a  $1 \leq j \leq \ell$  with  $\text{unmark}(\max_{>coef}(\text{coefs}(p(\vec{c})))) = \alpha_j(\vec{c}) \neq 0$  and  $\alpha_i(\vec{c}) = 0$  for all  $1 \leq i \leq j-1$ . Thus,

$$\text{unmark}(\max_{>coef}(\text{coefs}(p(\vec{c})))) \triangleright 0 \quad \text{iff} \quad (\text{red}(p \triangleright 0))(\vec{c}) \text{ holds,}$$

where  $\text{red}$  is defined as in the explanation after Lemma 34. Hence, (11) is equivalent to

$$\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(p \triangleright 0). \quad (18)$$

As  $\vec{q}$  can clearly be transformed into  $\vec{q}_{norm}$  in polynomial time,  $p$  can be obtained from the inputs  $\varphi$  and  $\vec{q}$  in (5) in polynomial time. Thus, the size of  $p$  is polynomial in the size of the input and hence we can compute and sort  $\text{coefs}(p)$  in polynomial time. Furthermore,  $\text{red}(p \triangleright 0)$  is a disjunction of at most  $\ell + 1$  subformulas, where each subformula consists of at most  $\ell$  (in-)equations over  $\text{coefs}(p)$ . As  $\ell$  is bounded by the size of  $p$ ,  $\text{red}(p \triangleright 0)$  can be computed in polynomial time. Since  $\psi_F$  is part of the input of (5), it follows that (18) can be computed in polynomial time. □

### A.14 Proof of Thm. 36

*Proof.* We have to prove

$$(12) \iff \exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(\xi), \quad (19)$$

where  $\text{red}(\xi)$  results from replacing each atom  $p \triangleright 0$  in  $\xi$  by  $\text{red}(p \triangleright 0)$ . Since each  $\text{red}(p \triangleright 0)$  can be computed in polynomial time due to Lemma 34, the computation of the formula “ $\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(\xi)$ ” clearly works in polynomial time.

To prove (19), we introduce the notion of a *fundamental* set. Let  $p_1 \triangleright_1 0, \dots, p_k \triangleright_k 0$  denote the atoms in  $\xi$ . We call a subset  $I \subseteq \{1, \dots, k\}$  *fundamental* if  $\bigwedge_{i \in I} p_i \triangleright_i 0 \implies \xi$ . Recall that w.l.o.g., we can assume that  $\xi$  does not

contain any Boolean connectives except  $\wedge$  and  $\vee$ . Thus, whenever  $\xi \neq \text{false}$ , the formula  $\xi$  must have fundamental sets. Clearly, we have

$$\exists \vec{x} \in \mathbb{R}_{\mathbb{A}}^d. \psi_F \wedge \text{red}(\xi) \iff \exists \text{ fundamental set } I. \psi_F \wedge \bigwedge_{i \in I} \text{red}(p_i \triangleright_i 0) \text{ is valid.}$$

Thus, to prove (19), it suffices to show the following:

$$(12) \iff \exists \text{ fundamental set } I. \psi_F \wedge \bigwedge_{i \in I} \text{red}(p_i \triangleright_i 0) \text{ is valid.} \quad (20)$$

For the “ $\Leftarrow$ ”-direction of (20), assume that there is such a fundamental set, i.e.,

$$\psi_F \wedge \bigwedge_{i \in I} \text{red}(p_i \triangleright_i 0)$$

is valid. Then as in the proof of Lemma 34, we obtain that for each  $i \in I$ , there is an  $n_i \in \mathbb{N}$  such that

$$\exists \vec{x} \in F. \forall n \in \mathbb{N}_{>n_i}. p_i \triangleright_i 0.$$

As  $I$  is finite,  $n_{\max} = \max\{n_i \mid i \in I\}$  exists. Hence, we get

$$\exists \vec{x} \in F. \forall n \in \mathbb{N}_{>n_{\max}}. \bigwedge_{i \in I} p_i \triangleright_i 0.$$

Since  $I$  is fundamental, this implies (12).

For the “ $\Rightarrow$ ”-direction, assume (12). Then there is a  $\vec{c} \in F$  and an  $n_0 \in \mathbb{N}$  such that for each  $n \in \mathbb{N}_{>n_0}$ , there is a fundamental set  $I_n$  such that  $\bigwedge_{i \in I_n} p_i(\vec{c}) \triangleright_i 0$  holds. As there are only finitely many fundamental sets, there is some fundamental set  $I$  that occurs infinitely often in  $(I_n)_{n \in \mathbb{N}_{>n_0}}$ . Hence we get

$$\exists n_0 \in \mathbb{N}. \exists^\infty n \in \mathbb{N}_{>n_0}. \bigwedge_{i \in I} p_i(\vec{c}) \triangleright_i 0. \quad (21)$$

By definition of poly-exponential expressions, each  $p_i(\vec{c})$  is weakly monotonic in  $n$  for large enough  $n$ . Thus, (21) implies

$$\exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \bigwedge_{i \in I} p_i(\vec{c}) \triangleright_i 0.$$

As  $\vec{c} \in F$ , this implies that there is a fundamental set  $I$  such that  $\psi_F \wedge \bigwedge_{i \in I} \text{red}(p_i \triangleright_i 0)$  holds.  $\square$

### A.15 Proof of Thm. 37

*Proof.* By Thm. 26, termination of *twn*-loops is reducible to termination of *tnn*-loops. Given a *tnn*-loop  $(\varphi, \vec{a})$ , we obtain  $\vec{q}_{norm} \in (\text{NPIE}[\vec{x}])^d$  such that  $(\varphi, \vec{a})$  is (eventually) non-terminating iff (8) holds, where  $\varphi$  is a propositional formula over the atoms  $\{\alpha \geq 0, \alpha > 0 \mid \alpha \in \mathbb{R}_{\mathbb{A}}[\vec{x}]\}$ . Hence,  $\varphi(\vec{q}_{norm})$  is a propositional formula over the atoms  $\{p \triangleright 0 \mid p \in \text{NPIE}[\vec{x}], \triangleright \in \{\geq, >\}\}$ . Thus, by Thm. 36, validity of (8) resp. (12) is reducible to  $\text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ .  $\square$

### A.16 Proof of Cor. 38

*Proof.* By Thm. 26, termination of *tw*n-loops is reducible to termination of *tn*n-loops. By Thm. 37, termination of *tn*n-loops is reducible to invalidity of a formula  $\chi \in \text{Th}_{\exists}(\mathcal{S}, \mathbb{R}_{\mathbb{A}})$ . If  $\mathcal{S} = \mathbb{R}_{\mathbb{A}}$ , then validity of  $\chi$  is decidable, and if  $\mathcal{S} = \mathbb{Z}$  or  $\mathcal{S} = \mathbb{Q}$ , then validity of  $\chi$  semi-decidable [11, 51]. But  $\chi$  is valid iff the loop is non-terminating. Hence, non-termination is decidable for  $\mathcal{S} = \mathbb{R}_{\mathbb{A}}$  and semi-decidable if  $\mathcal{S} = \mathbb{Z}$  or  $\mathcal{S} = \mathbb{Q}$ . The claim (b) for  $\mathcal{S} = \mathbb{R}_{\mathbb{A}}$  follows since *deciding* non-termination is equivalent to deciding termination. Finally, (a) and the claim (b) for  $\mathcal{S} = \mathbb{R}$  follow due to elementary equivalence of  $\mathbb{R}_{\mathbb{A}}$  and  $\mathbb{R}$ .  $\square$

### A.17 Proof of Lemma 39

*Proof.* Let  $\xi$  contain the atoms  $p_i \triangleright_i 0$  for  $1 \leq i \leq k$ . We have:

$$\begin{aligned}
 & \vec{c} \text{ witnesses eventual non-termination of } (\varphi, \vec{a}) \\
 \iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. (\varphi(\vec{q}_{norm}))(\vec{c}) && \text{(by (8))} \\
 \iff & \exists n_0 \in \mathbb{N}. \forall n \in \mathbb{N}_{>n_0}. \xi(\vec{c}) \\
 \iff & \psi_F(\vec{c}) \wedge \text{red}(\xi)(\vec{c}) && \text{(as in the proof of Thm. 36)}
 \end{aligned}$$

$\square$

### A.18 Proof of Thm. 45

*Proof.* By Thm. 44, termination of linear-update loops over the reals is in  $\forall\mathbb{R}$ . We have  $\forall\mathbb{R} \subseteq \text{EXPTIME}$ . Below, we will prove that linearizing a loop with the procedure described in [38] is also in EXPTIME under the restrictions stated in the theorem. Note that linearizing a *tw*n-loop with the technique of [38] yields a linear-update *tw*n-loop. So in particular, it has real spectrum (since the eigenvalues of a triangular matrix are the diagonal entries). Combining this result with the above observation that termination of linear-update loops is in EXPTIME then yields a decision procedure for termination of *tw*n-loops over the reals which is in 2-EXPTIME if the number of variables is unbounded, i.e., this proves (a).

In case (b), the number of variables is bounded by the constant  $D$ . In this case, linearizing the loop is again in EXPTIME. By Lemma 41, from this linear-update loop one can generate a formula  $\psi$  in polynomial time which is valid iff the loop is non-terminating. By [2], due to the bound on the number of variables, one can also check in polynomial time whether this formula is valid. So in case (b), we obtain a decision procedure for termination of *tw*n-loops over the reals which is in EXPTIME.

**Linearization is in EXPTIME:** We consider the procedure from [38] to linearize a *tw*n-loop. So let  $(\varphi, \vec{a})$  be a *tw*n-loop and let  $mdeg$  be the maximal degree of the polynomials occurring in  $\vec{a}$ . W.l.o.g. we may assume that  $x_1 \succ_{\vec{a}} \dots \succ_{\vec{a}} x_d$ .



The basic idea of the algorithm in [38] is to add variables which represent monomials of the form  $\prod_{i=1}^d x_i^{z_i}$  in the variables  $x_1, \dots, x_d$ , where  $z_1, \dots, z_d \in \mathbb{N}$ .<sup>10</sup> To ease notation, we abbreviate  $\prod_{i=1}^d x_i^{z_i}$  to  $\vec{x}^{\vec{z}}$  where  $\vec{z} = (z_1, \dots, z_d) \in \mathbb{N}^d$ . We denote the variable representing such a monomial  $m$  in the linearized system by  $y_m$ . So in particular,  $y_{x_i}$  represents  $x_i$  in the linearized update. Moreover, if  $m = \vec{x}^{\vec{z}}$  and  $m' = \vec{x}^{\vec{z}'}$  are two monomials, we define the lexicographic ordering  $\succ$  on monomials, where  $m \succ m'$  holds if  $\vec{z} \succ_{lex} \vec{z}'$ . As usual, we write  $m \succeq m'$  to denote  $m \succ m'$  or  $m = m'$ .

The procedure from [38] now computes a *linear-update twin-loop*  $(\varphi \wedge \varphi_{inv}, \vec{a}')$  over the variables  $y_m$  for all monomials  $m$  where  $\deg(m) \leq mdeg$ . Here,

$$\varphi_{inv} = \bigwedge_{m=\vec{x}^{\vec{z}}, \deg(m) \leq mdeg} \left( y_m = \prod_{i=1}^d y_{x_i}^{z_i} \right).$$

Moreover, for each monomial  $m$  with  $\deg(m) \leq mdeg$ , the entry  $a'_m$  of  $\vec{a}'$  corresponding to the update of  $y_m$  only contains the variables  $y_{m'}$  with  $m \succeq m'$ .

The computation of the entries in  $\vec{a}'$  works according to the lexicographic ordering  $\succ$ , i.e., when computing  $a'_m$ , we can assume that we have already computed  $a'_{m'}$  for all  $m'$  with  $m \succ m'$ . More precisely, when computing the update  $a'_{m_1 \cdot m_2}$  for a monomial  $m_1 \cdot m_2$ , we can assume that we already know the updates  $a'_{m_1}$  and  $a'_{m_2}$  for the monomials  $m_1$  and  $m_2$ . Let  $a'_{m_i} = \sum_{m_i \succeq m'_i} r_{m'_i} \cdot y_{m'_i} + C_i$ , where  $i \in \{1, 2\}$  and  $r_{m'_i}, C_i \in \mathbb{R}_{\mathbb{A}}$ . Then we have

$$\begin{aligned} a'_{m_1} \cdot a'_{m_2} &= \left( \sum_{m_1 \succeq m'_1} r_{m'_1} \cdot y_{m'_1} + C_1 \right) \cdot \left( \sum_{m_2 \succeq m'_2} r_{m'_2} \cdot y_{m'_2} + C_2 \right) \\ &= \sum_{m_1 \succeq m'_1, m_2 \succeq m'_2} (r_{m'_1} \cdot r_{m'_2}) \cdot y_{m'_1} \cdot y_{m'_2} \\ &\quad + \sum_{m_2 \succeq m'_2} (C_1 \cdot r_{m'_2}) \cdot y_{m'_2} + \sum_{m_1 \succeq m'_1} (C_2 \cdot r_{m'_1}) \cdot y_{m'_1} + C_1 \cdot C_2. \end{aligned}$$

Since the update  $a'_{m_1 \cdot m_2}$  for the monomial  $m_1 \cdot m_2$  should have the same value as  $a'_{m_1} \cdot a'_{m_2}$ , the technique from [38] defines the following.

$$\begin{aligned} a'_{m_1 \cdot m_2} &= \sum_{m_1 \succeq m'_1, m_2 \succeq m'_2} (r_{m'_1} \cdot r_{m'_2}) \cdot y_{m'_1 \cdot m'_2} \\ &\quad + \sum_{m_2 \succeq m'_2} (C_1 \cdot r_{m'_2}) \cdot y_{m'_2} + \sum_{m_1 \succeq m'_1} (C_2 \cdot r_{m'_1}) \cdot y_{m'_1} + C_1 \cdot C_2. \end{aligned}$$

Here, instead of the non-linear monomial  $y_{m'_1} \cdot y_{m'_2}$ , one uses the variable  $y_{m'_1 \cdot m'_2}$ . Hence, the resulting linear update  $a'_{m_1 \cdot m_2}$  indeed only contains variables  $y_{m'}$  where  $m_1 \cdot m_2 \succeq m'$ , since  $m_1 \succeq m'_1$  and  $m_2 \succeq m'_2$  implies  $m_1 \cdot m_2 \succeq m'_1 \cdot m'_2$ . So for these variables  $y_{m'}$ , the linear update  $a'_{m'}$  has already been computed.

<sup>10</sup> We only consider monomials with positive degree, i.e, we leave out the constant monomial  $m = 1$ .

We observe the following:

**Corollary 46.** *Given the updates of  $y_{m_1}$  and  $y_{m_2}$ , the time required for computing the coefficients in the update of  $y_{m_1 \cdot m_2}$  is polynomial in the coefficients of the updates of  $y_{m_1}$  and  $y_{m_2}$ .*

Now, in the worst case, the algorithm from [38] needs to compute the updates for *all* monomials in the variables  $x_1, \dots, x_d$  with degree at most  $mdeg$  in a bottom-up fashion in the way above. As stated in [38], there are  $\binom{d+mdeg}{mdeg} = \binom{d+mdeg}{d} \in \mathcal{O}((mdeg+d)^{mdeg}) \cap \mathcal{O}((mdeg+d)^d)$  monomials of degree at most  $mdeg$ , i.e., the algorithm needs  $\mathcal{O}((mdeg+d)^{mdeg}) \cap \mathcal{O}((mdeg+d)^d)$  many iterations to compute the update of the linearized loop. By the precondition of the theorem, we have  $mdeg \leq M$  or  $d \leq D$ , i.e., in this case  $\binom{d+mdeg}{mdeg}$  can be bounded by a polynomial in  $d$  if  $mdeg \leq M$ , or by a polynomial in  $mdeg$  if  $d \leq D$ . Each step of the algorithm takes polynomial time in the input of *this step* as seen above. However, one step increases the coefficients polynomially. Since we have to do  $\mathcal{O}((mdeg+d)^{mdeg}) \cap \mathcal{O}((mdeg+d)^d)$  many steps, this means that the coefficients can grow exponentially in comparison to the coefficients in the original update  $\vec{a}$ . But even then, the time needed for a step of the linearization algorithm is at most *exponential* in the coefficients in  $\vec{a}$ . Since the algorithm only takes polynomially many steps under the restrictions given in the theorem, this then means that the linearization algorithm from [38] is in EXPTIME.  $\square$

*Remark 47.* In the proof of [Thm. 45](#) we have seen that linearizing a *tnn*-loop is in EXPTIME, whenever the maximal degree occurring in the polynomials in the updates or the number of variables is bounded in advance. This is due to the fact that we also take the computed coefficients for the polynomials in the updates of the resulting linear-update loop into account. When only considering the number of newly added variables, i.e., when the coefficients do *not* need to be taken into account, then (as stated in [38]), the algorithm only introduces polynomially many new variables whenever the number of variables or the degree of the polynomials in the updates is bounded in advance.

However, as seen in the loop (13), if neither the number of variables nor the maximal degree occurring in the update is bounded, then the problem of linearizing arbitrary *tnn*-loops is *not* in EXPTIME.