- 1 Rationale and design of the Healthy Brain Study: an accessible resource for understanding the
- 2 human brain and how it dynamically and individually operates in its bio-social context

- **HBS** consortium
- 5 Corresponding authors on behalf of the HBS consortium: lucy.overbeek@radboudumc.nl and
- 6 <u>Guillen.fernandez@donders.ru.nl</u>

- Abstract
- The endeavor to understand the human brain has seen more progress in the last few decades than in the previous two millennia. Still, our understanding of how the human brain relates to behavior in the real world and how this link is modulated by biological, social, and environmental factors is limited. To address this, we designed the Healthy Brain Study (HBS), an interdisciplinary, longitudinal, cohort study based on multidimensional, dynamic assessments in both the laboratory and the real world. Here, we describe the rationale and design of the currently ongoing HBS. The HBS is examining a population-based sample of 1,000 healthy participants (age 30-39) who are thoroughly studied across an entire year. Data are collected through cognitive, affective, behavioral, and physiological testing, neuroimaging, bio-sampling, questionnaires, ecological momentary assessment, and real-world assessments using wearable devices. These data will become an accessible resource for the scientific community enabling the next step in understanding the human brain and how it dynamically and individually operates in its bio-social context. An access procedure to the collected data and bio-samples is in place and published on https://www.healthybrainstudy.nl/en/data-and-methods.
- 22 https://www.trialregister.nl/trial/7955

Introduction

The human brain is seen as civilization's most precious resource ¹, both creating and interacting with our increasingly complex environment, it enables us to be conscious and social human beings. Brain functioning also plays a pivotal role in major societal challenges such as health, demographic change, and well-being. Due to developments in different scientific fields, the endeavor to understand the human brain has seen more progress in the last few decades than in the two millennia before. However, we think that current brain research suffers from at least five key limitations and we set up the Healthy Brain Study (HBS) to tackle these five limitations together and, thereby, to facilitate our understanding of how the human brain relates to behavior in the real world and how this link is modulated by biological, social, and environmental factors. In the following paragraphs, we explain the five main design choices of the HBS.

Firstly, a reductionist approach – in which researchers try to understand reality by focusing on a limited number of variables – has been understandably popular as it is vital to obtain detailed mechanistic insights. However, complex dynamical systems, like the human brain, cannot be properly understood by focusing on just one aspect at a time ²⁻⁴. Human brain functioning includes enabling consciousness and cognition, generating emotions, and producing adaptive behavior, and it performs all of these functions while embedded in its biological and social (bio-social) environment ⁵. To enable researchers to understand the complexity of human brain functioning in its bio-social context, the HBS provides a *broad range of variables* within a holistic approach.

Secondly, the brain's operations cannot be fully understood by single assessments obtained at a specific point in time, but require repeated measurements or continuous monitoring. Single-session assessments may be sufficient to uncover stable traits or processes. However, they do not capture changes in brain functioning that constitute a core feature of our plastic and adaptive brain ^{6,7}. Similarly, the body and the social environment are subject to change. For example, seasonality is

observed in affect ^{8,9}, behavior ^{9,10}, and biological ¹¹⁻¹⁴ and social ⁹ factors. Most of the studies mentioned were cross-sectional and explicitly stress the need for longitudinal studies that assess within-subject variation. Therefore, in the HBS, participants perform *repeated assessments* in three different seasons over one year starting at varying time points within a year. Thereby, we aim to reliably and validly capture changes in human brain operations that may be related not only to seasonality, but also to relevant life events and incidental or dynamic changes in biological factors (e.g., inflammation markers), social factors (e.g., household composition, work relations, friendships, politics, media exposure, lockdown), and environmental factors (e.g., daylight hours, exposure to chemicals).

Thirdly, group averages are critical in revealing general principles, but they gloss over differences that make us individual human beings. The human brain is arguably the most individual organ we have and is shaped by our experiences throughout life. Therefore, a large and rich sample is required before *single subject inferences* can be made about underlying principles of diversity in cognition, affect, and behavior ^{15,16}. Given this, the HBS aims to include a broad range of repeated assessments of 1,000 participants.

Fourthly, laboratory assessments enable well-controlled analyses, but they may show low ecological validity in generalizing cognition, affect, and behavior to real-world settings. To understand cognition, affect, and behavior more comprehensively, there is a need for assessments both in the laboratory as well as in the real world ^{17,18}. In the HBS, we perform a *real-world assessment* of physical activity, stress, and sleep with validated wearable devices. Furthermore, we apply ecological momentary assessments using a smartphone application. Taken together, these assessments enable us to understand cognition, affect, and behavior in the context where they naturally occur.

Finally, a healthy volunteer selection bias is a frequent problem in both cohort studies and neuroscience studies. For example, UK Biobank participants were more likely to be female, have a healthy lifestyle, and live in less socioeconomically deprived areas compared to the general population ¹⁹. Also, students, the usual participants in cognitive neuroscience studies, function well, are often relatively healthy and have a high socioeconomic status ²⁰. Also, most population-based cohorts and large-scale studies include either developing populations ²¹⁻²³ or advanced aging populations ²⁴⁻²⁶. Therefore, the HBS includes a broad *population-based sample* of individuals who are 30-39 years old that reflects the general population in terms of gender and educational attainment. The age range was chosen to represent adults beyond the age of developmental brain changes and before the onset of brain changes due to advanced aging or neurodegenerative disease. The lower limit of 30 years excludes any neurodevelopment effect as the brain has matured by this point ²⁷. Also, 30-39 is a socially challenging age range because it is generally characterized by a relatively high number of rather impactful life events (e.g., family planning, career-related changes, buying a house).

In conclusion, the unique feature of the HBS is that it combines in-depth phenotyping of a large range of cognitive, affective, behavioral, and social dimensions with a biological sampling of brain and body-related processes. This enables the extraction of a detailed bio-social fingerprint for the participants in the cohort. Such a detailed fingerprint is currently not available. The availability of HBS will contribute to a better understanding of risks and potentials in behavior in the real world at the individual level. This paper describes the rationale and design of the currently ongoing HBS, which originated from an interdisciplinary, team science ²⁸ based cross-faculty initiative from the Radboud campus in Nijmegen, the Netherlands, including Radboud University, Radboud University Medical Center, and the Max Planck Institute for Psycholinguistics.

Methods/design

Study design and setting

The HBS is a longitudinal cohort study in both laboratory and real-world settings. All laboratory assessments take place at a single-center on Radboud campus, Nijmegen, the Netherlands.

Participants

The HBS aims to include 1,000 participants (500 men and 500 women) from the Nijmegen region (≤ 15 km) of whom 220 have a low, 340 a middle, and 430 a high level of education. Nijmegen is a medium-sized city in the east of the Netherlands with 176,731 citizens on the 1st of January 2019 of whom 74% are native Dutch, which is comparable to the overall proportion of native Dutch citizens of the Netherlands (76%) ²⁹. In contrast, large cities (> 500,000 citizens) in the west of the Netherlands like Amsterdam, Rotterdam, and The Hague have respectively 46%, 48%, 45% native Dutch citizens ²⁹. Regarding educational attainment, 22% of Nijmegen citizens are primary and secondary educated (low level), 34% are primary, secondary, and vocationally educated (middle level), and 43% of the population have also a university degree (high level). Nijmegen has less citizens with low and middle level of education and more citizens with high level of education compared to the overall proportions of Dutch citizens (28%, 41%, 30% of citizens have respectively low, middle, and high level of education) ²⁹. In comparison, some large cities in the Netherlands have a higher proportion of citizens with a high level of education (e.g., Amsterdam 48%, Utrecht 52%), while other large cities have a higher proportion of citizens with a low level of education (e.g., Rotterdam 32%, The Hague 31%) ²⁹.

Inclusion criteria are age 30-39 years, living in the Nijmegen region (≤ 15km; because of feasibility), willingness, and ability to follow the study protocol. Exclusion criteria are: not speaking, reading, and/or understanding the Dutch language (minimum B1 level), a prior history of significant psychiatric or neurological illness (self-report), a current disease that affects the brain, a current medication that is therapeutically targeted at the brain (e.g., antidepressants, methylphenidate), pregnancy, contra-

indication for MRI (metal or devices in the upper body (cardiac pacemaker, cochlear implant, aneurism clip), previous brain surgery, moderate to severe claustrophobia), contra-indication for the submaximal Åstrand cycle test (current use of beta-blockers, a current disease that hinders physical exercise), contra-indication for the cold pressor test (Raynaud's phenomenon, chronic pain syndrome in shoulder or arm, open wounds on arm or hand, scleroderma, arteriovenous fistula or shunt, presence of (unstable) angina pectoris).

Recruitment

We apply different strategies to recruit participants. Firstly, municipalities, general practitioners, and employers of different sectors based in the Nijmegen region send the HBS invitation and research flyer to their citizens, clients, and employees, respectively. Employers are asked to sponsor the study by (partly) exempting their employees from three working days which allows them to participate in three lab visits. Employees remain entirely free to decide whether or not they want to participate. Also, campaigns to increase awareness of the HBS have been launched.

Potential participants fill out contact details in an online form on the website https://www.healthybrainstudy.nl and receive the study brochure. Participants can watch short videos on the website that explain the various tests and assessments or learn about the experiences of an HBS participant. Participants are contacted via phone and invited to a face-to-face information meeting on the Radboud campus. Participants provide written informed consent at this meeting before participation.

Ethics

The Institutional Review Board of Radboud University Medical Center approved the HBS (reference number: 2018-4894) in accordance with the latest revision of the Declaration of Helsinki ³⁰. Incidental findings could occur both while conducting the study (e.g., observed during assessments) and while

using the data and biosamples in the future to answer research questions. If a researcher or research assistant notices a potential finding incidentally, he/she will contact the principal investigator, who approaches an incidental findings committee. At the Radboud Campus, such committees exist for neuroimaging and genetics. For other findings, the principal investigator contacts a medical doctor with relevant expertise. If, according to the committee or medical expert, no clinically relevant finding is identified, the participant remains uninformed. In all other cases, the participant's general practitioner is sent a letter describing the findings. At the same time, the participant receives a request to contact their general practitioner. Participants must consent to this procedure and provide the contact details of their general practitioner, otherwise, they are not allowed to participate.

Participant panel, feedback of participants, incentives, and citizen science

A participant panel consisting of twelve people (age 30-39, 6 women and 6 men) was set up to aid in the design of the study. The panel advises on communication materials and incentives. For example, the panel gives feedback on the website, study information, posters, and flyers. Moreover, the first 243 participants filled out a questionnaire on their experience of the first lab visit, which provided us with feedback on the study procedures and on keeping participants involved. For example, we developed an online dashboard, because participants indicated that they would prefer more individual feedback on results. Participants receive gadgets after each assessment, and we organize (online) participant events. After completion of the study protocol, participants receive €150 with a maximal addition of €50 for assessment specific incentives.

Besides, a citizen science platform is used to involve participants as well as other citizens in generating research topics and questions that can be investigated with the HBS resource ³¹. We 'crowdsource' lists of research topics and/or research questions that participants and citizens think are useful for examining with the HBS resource. At the same time, they also rate the importance of the crowd-

generated suggestions by other participants and citizens resulting in an overview that reflects the relevance and prioritization of their overall input.

Quality management and safety

Research assistants and nurses received extensive training for the assessments undertaken as part of the study protocol. We adapted existing standardized operating procedures (SOPs) if available and developed a new SOP otherwise. An independent study monitor annually performs checks to ensure that the study protocol is followed.


Data management

We use Ldot, which is a web application that only stores personal and logistical data, for communication with our participants (https://www.health-ri.nl/services/ldot-workflow-tool-clinical-research). For data acquisition, we use Castor EDC (https://data.castoredc.com) to provide electronic case report forms and online questionnaires. In addition, a smartphone application for ecological momentary assessments was developed. After participants have performed the real-world assessments, our data managers extract the raw data that is stored locally on the device. Biosamples are stored at the Radboud Biobank with their sample tracking system, sample processing SOPs, and standardized sample storage conditions being employed 32. Furthermore, a Polymorphic Encryption and Pseudonymization (PEP) infrastructure was developed for the HBS to protect all data streams and the privacy of participants 33,34 (Figure 1). Ldot, Castor EDC, the smartphone application, and PEP meet the requirements of the European General Data Protection Regulation.

For each participant, the PEP-system generates unique pseudonyms for the different assessments to avoid the coupling of data to an individual participant during the data collection phase (step 1: collect). A backup of the data is stored locally (step 2: produce) and a copy is encrypted and transferred to the data repository (step 3: store). In the same step, the data are cryptographically pseudonymized. The

data can only be decrypted in the processing environment where scientific analyses are performed (step 4: process). The PEP method ensures that different datasets obtained from the repository cannot be linked easily by different research projects because pseudonyms identifying a single participant are personalized at the project level, and data transfer can be minimized based on researchers' requirements. Derived data, produced by researchers, can be stored in the data repository (step 5: contribute) for future use by other researchers using their researcher-specific pseudonyms.

The PEP-system was created to deal with the rigidity of the traditional encryption/decryption process by using polymorphic encryption. PEP ensures that there is no need to a priori fix the encryption key for the data. The PEP system enables different research teams to have access to the entire dataset or only a subset (of participants and variables) of the data source with a specific, personalized decryption key. Due to its additional security, the PEP system is an ideal approach to store, manage, and share sensitive personal data in a research data repository that reduces the risk of a participant's privacy being violated.



Who	Assessor	Producer	Repository	Researcher
Identity information	Identifiable	Pseudonymised Encrypted/ (by data source) Pseudonymised		Pseudonymised (researcher specific)
Data	Partly Clear text	Clear Text	Encrypted	Clear text
Linkability	Linkable to person	Unlinkable	Unlinkable	Linkable to researcher-specific participant-pseudonym

Figure 1. The Polymorphic Encryption and Pseudonymization (PEP) infrastructure.

Measures

The following paragraphs describe the measures briefly, while the supplementary information provides detailed descriptions. Each assessment starts with pre-visit online questionnaires, followed by a burst week of real-world assessments, followed by a whole day lab visit, which in turn is followed by post-visit online questionnaires and assessments (Figure 2). Only those constructs that may be sensitive to change during one year (states) are repeated during the second and third assessments. The stable (trait) measures are equally distributed over the three assessments. The majority of measures are validated in prior research.

Start assessment 1

4 months after assessment 2

Assessment 2

Assessment 3

Pre-visit online questionnaires

1 hour in 1 week

Burst week - real-world assessments

4 months after assessment 2

Assessment 3

1 hour in 1 week

45 minutes a day

45 hours

5 hours

Post-visit online questionnaires and assessments



Figure 2. Design of data collection in the Healthy Brain Study

Pre-visit online questionnaires

Participants fill out questionnaires before the start of the burst week to assess baseline characteristics.

The questionnaires cover general demographic questions and questions about lifestyle and well-being

(Table 1a).

Burst week with real-world assessments

The burst week consists of a real-world assessment of physical activity, stress, and sleep using validated wearable devices (Table 2) and ecological momentary assessments (EMA) using a smartphone application. The questionnaire for EMA covers mood, social company, online social interactions, context, control items, retrospection, anticipation, and substance use. In addition, participants perform the home collection of stool, urine, saliva, and diffusive sampling of chemicals using silicone

wristbands during the burst week (Table 3).

Lab visit Radboud campus

Each eight-hour lab visit includes bio-sampling (Table 3), neuroimaging (Table 4), physiological (Table 2), cognitive (Table 5), affective (Table 5), behavioral (Table 5), and sensory assessments (Table 6). To avoid systematic carry-over and fatigue effects, the order of assessments varies between and within participants except for fasting blood sampling and blood pressure at the start of the day.

Post-visit online questionnaires and assessments

Participants fill out an online questionnaire assessing (mental) health, life events, social/relationships, work, politics, personality, and literacy after each lab visit (Table 1b). Also, participants perform several online assessments about decision-making, narrative reading, and solidarity (Table 7). After their third and final lab visit, participants are invited to complete the 'Individual Differences in Language Skills' test battery (Table 8) assessing participants' linguistic knowledge, as well as linguistic processing and general cognitive skills.

COVID-19 questionnaire

From March until July 2020, when the assessment of participants was not allowed due to the COVID-19 measures, the included participants at that point (N=158) received a monthly questionnaire addressing behavior and worries regarding COVID-19, currently experienced anxiety ³⁵, stress ³⁶, and well-being ³⁷. Moreover, loneliness ³⁸, sedentary behavior ³⁹, and sleep quality ⁴⁰ were assessed. We used the same questionnaires as we use in the three repeated assessments (Table 1a and 1b).

Table 1a. Pre-visit online questionnaires

Domain	Name of the questionnaire	What does it measure?	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3	Ref
General information	Demographic and socio- economic background	Demographic data, the highest level of education, income, household composition	10	Х	Х	х	41
	Pregnancy	Number of pregnancies, time to pregnancy, pregnancy outcome, fertility treatment, hormones (anticonception), breastfeeding, current child wish	3	Х			
	Menstrual cycle	Menstrual cycle	1	X	Х	X	
Lifestyle	Smoking history	Past behavior, age of onset	1	Х			
	Smoking	Current behavior, frequency, and quantity	1	Х	Х	Х	
	Fagerstrom Test of Nicotine Dependence (FTND)	Nicotine dependence (for current or ever smokers)	2	х	х	х	42
	Alcohol	Frequency and quantity in the last month, age of onset of alcohol use, binge drinking	2	Х	Х	х	
	Alcohol Use Disorder Identification Test (AUDIT)	Heavy alcohol use and associated problems	3	х	х	х	43
	Substance matrix Mate-q	Amount and frequency of substance use	5	Х	Х	Х	44
	Food Frequency Questionnaire (FFQ)	Quantitative food intake	45	Х			45-48
	Sedentary Behavior Questionnaire (SBQ)	Sedentary behavior in various domains (e.g. home, work, transportation)	5	Х	Х	Х	39
	Pittsburgh Sleep Quality Index (PSQI)	Sleep quality	5	Х	Х	х	40
	Dream Recall Frequency Scale (DRFS)	Dream recall	1	Х	Х	Х	49
	The Internet Gaming Disorder Scale	Problematic gaming	2	х	х	х	50
	The Social Media Disorder Scale	Problematic social media use	2	Х	Х	X	51
	Short Media Multitasking Measure (S-MMM)	Use of different media simultaneously	1	Х	Х	х	52
Well-being	Satisfaction with life scale	Well-being	2	Х	Х	Х	53
_	Cantril ladder	Well-being	1	Х	Х	Х	37

Five Facet Mindfulness	Mindfulness	10	Х	Х	Х	54
Questionnaire – Short						
Form (FFMQ)						

Table 1b. Post-visit online questionnaires

Domain	Name of the questionnaire	What does it measure?	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3	Ref
Exposure	Exposure	Exposure from environment	5	Х	Х	Х	
Health	Over-the-counter medication	Use of nonprescription medication like pain relievers, cough suppressants, etc.	1	Х	Х	х	55
	Health complaints	Complaints like tiredness, nausea, back pain, headache, etc.	5	Х	Х	х	56
Mental Health	Adult ADHD Self-Report Scale (ASRS)	Symptom scale for ADHD	10	Х			57
	Autistic Trait Questionnaire (ATQ)	Autistic traits	5	х			58
	Self-Report Inventory of Depressive Symptomatology (IDS-SR) Anxiety Sensitivity Index Presence and severity of depressive symptoms Anxiety (trait)		5	х	х	х	59
	Anxiety Sensitivity Index (ASI)	Anxiety (trait)	5			Х	60
	State and Trait Anxiety Inventory (STAI-S)	Anxiety (state)	5	Х	Х	х	35
	Perceived Stress Scale (PSS)	Stress	5	х	х	Х	36
	Utrecht Burnout Scale (UBOS)	Burnout	3	Х	Х	х	61
	Reactive Proactive Aggression Questionnaire (RPQ)	Aggression	5	х	х	х	62
	Daily hassles	Daily hassles	5	Х	Х	Х	63
	Cognitive emotion regulation questionnaire (CERQ)	Cognitive regulation of emotion	5	Х	Х	х	64
Life events	Childhood Trauma Questionnaire (CTQ)	Adverse childhood experiences	5		Х		65
	Life events	Threatening life experiences	10	Х	Х	Х	66
Social/	UCLA loneliness scale	Loneliness	5	Х	Х	Х	38
Relationship	Need to belong scale	Belongingness	3	Х	Х	Х	67

	Multidimensional scale of Perceived Social Support (PSS)	Perceived social support	5	x	х	Х	68
Work	Exposure to work	Working hours, working schedules, type of employment	4	Х	Х	х	
	Survey Work-home Interaction – NijmeGen (SWING)	Work-life balance	4	х	х	Х	69
	Workplace commitment		5	Х	Х	Х	70
	Employability		5	Х	Х	Х	71,72
	Questionnaire on the Experience and Evaluation of Work (QEEW)	Job characteristics	7	X ¹			73
Politics	Populism index	Attitude toward populism	2	Х	Х	Х	
	Political efficacy Attitude towards national government and politics		2	х	Х	х	74
	Political participation	Political activities	1	Х	Х	Х	
	EU membership	Attitude towards EU membership	1	Х	Х	Х	75
Personality	BIG-5 NEO-FFI-3	Openness to experience, conscientiousness, neuroticism, extraversion, and agreeableness	10	Х			76
	Sensory Processing Sensitivity (SPS)	High sensitivity	5			х	77
	Barratt Impulsiveness Scale (BIS-11)	Impulsiveness	10		Х		78
	Self-control		10	Х			79
	New general self-efficacy scale	Self-efficacy	5		Х		80
	Dispositional greed	Greediness	3			Х	81
	Dark triad	Narcissism, Machiavellianism, psychopathy	5	Х			82
	Social investment attitudes	Attitudes toward corporate social responsibility	5		Х		83
Literacy	Numeracy test	Mathematical abilities	12		Х		84
,	Financial literacy	Financial attitudes, skills	20			Х	85
	Graph literacy	Ability to understand the meaning of graphs	10	Х			86
	Cultural intelligence	Ability to relate and work effectively across cultures	2			Х	87

¹ Participants fill out their job characteristics at the first assessment. In the second and third assessments, they fill out their job characteristics only in case of a new job.

Table 2. Physiological assessments

Domain	Measure	Location	Assessment 1	Assessment 2	Assessment 3	Ref
Physical	Fitness	Campus	Х	Х	Х	88
activity	Sedentary behavior	Home ¹	Х	Х	Х	89,90
Stress	Heart rate	Campus	Х	Х	Х	
		Home ¹	Х	Х	Х	91
	Heart rate variability	Home ¹	Х	Х	Х	91
	Skin conductance	Home ¹	Х	Х	Х	91
	Skin temperature	Home ¹	Х	Х	Х	91
	Startle eye-blink	Campus	Х	Х	Х	92
	Subjective stress levels	Campus	Х	Х	Х	93
		Home ²	Х	Х	Х	
Sleep	Sleep duration	Home ¹	Х	Х	Х	
	Sleep stages	Home ¹	Х	Х	Х	
Body	Weight	Campus	Х	Х	Х	
composition	Height	Campus	Х	Х	Х	
	Waist-hip circumference	Campus	Х	Х	Х	
	Body fat	Campus	Х	Х	Х	94
	Fat weight	Campus	Х	Х	Х	
	Total body water	Campus	Х	Х	Х	
	Skeletal muscle mass	Campus	Х	Х	Х	
	Body fat mass index	Campus	Х	Х	Х	
	Fat-free mass index	Campus	Х	Х	Х	
Pain	Subjective pain levels	Campus	Х	Х	Х	93,95
		Home ¹	Х	Х	Х	
	Electrical pain thresholds	Campus	Х	Х	Х	96,97
Cardiovascular	Blood pressure	Campus	Х	Х	Х	98
	Carotid artery reactivity	Campus	Х	Х	Х	99

¹ By wearable device, ² By ecological momentary assessment (EMA).

Table 3: Biosamples and silicone wristband

Biosample	Measure	Location	Assessment 1	Assessment 2	Assessment 3	Ref
Stool	Gut microbiome	Home	Х	Х	Х	12,32,100
Urine (first morning)	lons, such as calcium, potassium, sodium, magnesium	Home	х	х	Х	32,101
Saliva	Cortisol levels (short term; two baseline samples)	Home	Х	Х	Х	102
	Cortisol levels (short term; before, immediately after, and 20 minutes after acute challenge)	Campus	Х	Х	Х	
Blood - EDTA plasma	DNA	Campus	6 ml*			32
Blood - PAX gene	RNA	Campus	3x 2,5 ml*	3x 2,5 ml*	3x 2,5 ml*	
Blood - EDTA plasma	Future analyses	Campus	4x 10 ml* 1x 3 ml*	4x 10 ml* 1x 3 ml*	4x 10 ml* 1x 3 ml*	
Blood - serum	Future analyses (e.g., antibodies, proteomics)	Campus	10 ml*	10 ml*	10 ml*	
Blood - heparin	Future analyses (e.g., hormones, metabolomics)	Campus	2x 10 ml*	2x 10 ml*	2x 10 ml*	
plasma						
Hair	Cortisol levels (long term)	Campus	Х	Х	Х	103
Silicone wristband	Exposure to chemicals in the surrounding environment	Home	х	х	Х	104,105

*The indicated volumes refer to whole blood volumes

Table 4. Neuroimaging at the campus

Scan	Description	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3	Ref
Dummy scanner		10	Х			
T1w 3D MPRAGE	Anatomical scan	5	Х	Х	Х	
rfMRI	Resting-state functional scan followed by resting-state questionnaire	10	Х	Х	Х	106,107
nfMRI Movie functional scan		4,5	Х	Х	Х	
Scout, fieldmap, single-band reference EPIs	Auxiliary scans	2	Х	Х	Х	
Diffusion-weighted imaging scan	Structural connectivity characterizations and white matter tissue microstructural modelling	10	х			
High-resolution T1w 3D MP2RAGE anatomical scan	Quantitative T1 and cortical myelin mapping	10		Х		108
High-resolution T2*w scan	Quantitative T2* and magnetic susceptibility mapping for identification and quantification of iron deposition across the brain	10			х	109

Domain	Name of task	Measure	Description	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3	Ref
Cognition	Foraging task	The tendency to explore alternatives vs. to exploit a chosen alternative	Participants are presented with a tree and have to decide whether to harvest it for apples and incur a short harvest delay or move to a new tree and incur a longer travel delay	30	х	х	х	110
Cognition	Serial random- dot motion discrimination task	How predictions from the past are weighted with uncertain sensory information in the present	Participants judge the motion direction of moving dots (up vs. down) and receive auditory feedback about the correctness of their response	25	х	х	х	111
Cognition	Reward-driven reach-adaptation task	How willing people are to search for more rewarding outcomes in a motor task	Participants make shooting movements toward a target while holding a handle that records pulling and hand rotation movements	20	Х	х	х	112
Cognition	Paired associate memory task	Associative Memory	Participants memorize the associations between pictures of people and names in a study phase and the memory for these associations is tested in a test phase using a cuedrecall-test	7	х	х	х	113
Cognition	Tower of London	Executive function (planning)	Participants are presented with a startling array of different colored, same-sized balls and are requested to move the balls one-by-one, with as little moves as possible to a predefined goal array.	5	х	х	х	114
Affect	Contextual fear generalization task	Fear generalization	Participants are instructed to attend to the presented stimuli and learn to predict the shock in multiple contexts while assessing eye-blink startle electromyography, subjective	40	х	х	х	115

			report, and avoidance tendencies.					
Affect	Emotion regulation task	Emotion regulation	Participants are asked to actively regulate their emotions while either neutral or aversive pictures are presented on the computer screen	15	х	х	х	116
Affect	Self-referent encoding Task	Positive and negative memory bias	Participants endorse and memorize positive and negative words	8	Х	х	Х	117
Affect	Stimulus- response compatibility task	Automatic approach or avoidance tendency	Participants are presented with pictures (alcohol vs. soda) and are instructed to approach or avoid a certain condition	10	х	х	х	118
Behavior	Columbia card task	Risk preference	A card game that gives participants the repeated choice between risky options and safe options	22	Х	х	х	119
Behavior	Food auction task	Reliable index of people's preference for hedonic (short-term reward) vs. healthy food (long-term reward)	Participants bid on different food items (e.g., package of M&Ms, apple)	15	х	х	х	120

Table 6. Sensory assessments

Domain	Measure	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3
Vision	Contrast sensitivity	5	Х		
	Visual acuity	5		Х	
	Color vision	5			Х
Hearing	Hearing ability	1	Х	Х	Х

Table 7. Post-visit online assessments

Table 1.1 Ust-vi	on orinine assessifients						
Domain	Online task	What does it measure?	Duration (minutes)	Assessment 1	Assessment 2	Assessment 3	Ref
Decision	Llighor order riels	Diek ettitudes prudence and temperance in	15				121
Decision-	Higher-order risk	Risk attitudes, prudence, and temperance in	15	X	Х	Х	
making	preferences	financial decision-making					

	Equality equivalence test	Social preferences	10	Х	Х	Х	122
	Ambiguity	Ambiguous risk attitudes	10	Х	Х	Х	123
	Trust game	Trust and trustworthiness	10	Х	Х	Х	124
	Public good game	Altruism, conditional reciprocity	15	Х	Х	Х	125
	Time preferences	Temporal discounting	8	Х	Х	Х	126
Language	Narrative reading	Comprehension of and immersion into a narrative	15	Х	Х	Х	127
Solidarity	Vignettes	Culpability, in/out group	15	Х	Х	Х	128,129

Table 8. Individual Differences in Language Skills test battery


Domain	Online task	What does it measure?	Duration (minutes)	Ref
Cognition	Auditory simple and choice reaction time task	Processing speed	7	130
	Letter comparison	Processing speed	5	131,132
	Visual simple and choice reaction time task	Processing speed	7	130,133
	Digit span (forward & backward)	Auditory working memory	7	134
	Corsi block tapping (forward & backward)	Visual working memory	7	135,136
	Raven's advanced progressive matrices	Non-verbal intelligence	25	137
inguistic	Stairs4Words (2 Runs)	Linguistic experience: Vocabulary	7	
nowledge	Peabody Picture Vocabulary Test	Linguistic experience: Vocabulary	10	138,139
	Idiom recognition test	Linguistic experience: Knowledge of idiomatic expressions	3	
	Spelling test	Linguistic experience: Spelling	5	
	Author recognition test	Linguistic experience: Print exposure	5	140
	Prescriptive grammar	Linguistic experience: Prescriptive grammar knowledge	10	141
inguistic	Picture naming	Word production	7	130
rocessing	Rapid automatized naming	Word production	7	
	Verbal fluency	Word production	5	142
	Antonym production	Word production	5	143
	Maximal speech rate	Word production	3	
	Phrase generation	Sentence production	10	
	Sentence generation (active/passive sentence formulation)	Sentence production	12	
	Sentence generation (event apprehension)	Sentence production	10	
	Spontaneous speech	Sentence production	4	144
	Non-Word monitoring in non-word lists in noise	Word comprehension	10	
	Rhyme judgment	Word comprehension	5	
	Lexical decision	Word comprehension	7	130
	Semantic categorization	Word comprehension	5	
	Word monitoring in sentences in noise	Sentence comprehension	10	
	Grammatical gender cues	Sentence comprehension	10	145
	Verb-specific selective restrictions	Sentence comprehension	7	146,147
	Self-paced reading	Sentence comprehension	5	

Results - progress so far

Figure 3 presents the progress and milestones of the Healthy Brain Study. The first participant was included on the 9th of September, 2019. At the end of December, 2020, the HBS included 298 participants. Forty participants (13%) withdrew from the study so far, due to personal circumstances (e.g., pregnancy) or too much burden. Most participants withdrew after the first assessment. To date, participants performed 376 lab visits: 257 participants carried out the first assessment, 83 participants the second assessment, and 36 participants the third assessment completing the entire study protocol. The COVID-19 pandemic interrupted the data acquisition phase. At that point, the HBS included 158 participants. Due to the lockdown, we canceled all assessments involving physical interaction as of the 16th of March, 2020. The HBS resumed participant assessments on the 15th of July, 2020 in compliance with the directives in force in the Netherlands. As a result, some participants (48%) have more than four months between repeated assessments. Besides, some participants (10%) have a delay between the burst week with real-world assessments and the lab visit at the Radboud campus.

Healthy Brain Study Het dertigersbrein ontrafeld





304

Figure 3. Progress and milestones of the Healthy Brain Study

Discussion

This paper presents the design of the currently ongoing HBS, which will result in a unique and accessible resource for the scientific community and its public and private partners. Data are collected through cognitive, affective, behavioral, and physiological testing, neuroimaging, bio-sampling, questionnaires, ecological momentary assessment, and real-world assessment using wearable devices. We believe that the HBS complements other studies – small and large –, which together enable the scientific community to take the next step in understanding the human brain and how it dynamically and individually operates in its bio-social context. Here, we present examples of research opportunities including citizen science, reflect on the HBS design choices and study population, and discuss our data security system which enables future data sharing.

Examples of research opportunities

The HBS resource will be used to address expert and citizen-driven research questions that usually pertain to complex interactions between multiple factors. The first example of an expert-driven research question pertains to the association between income and positive affect. It was found among US residents that higher income was associated with more happiness and enjoyment, and less sadness and worry, but only up to a point (\$75.000 per year), above that, there was no relationship between income and emotional well-being ¹⁴⁸. The HBS resource can help explain the interplay between affect, social and biological data, and income. A second example of a complex interaction is that sedentary behavior is associated with poor health and higher mortality ^{149,150}. Merely standing up from time to time, e.g., to walk around a bit protects against part of this health risk ¹⁵¹. Existing research on this topic has mainly focused on the consequences of prolonged sitting and has overlooked the key question of why people choose to stand up (when they sit) or sit down (when they stand), in the first place. In other words, what psychological processes (e.g., related to effort, reward, affect, and fatigue) are associated with healthy and unhealthy sedentary behavior? Answering this question will pave the way for the development of novel, targeted interventions that will improve (occupational) health ¹⁵².

The HBS resource will also be used for citizen science. Different forms of citizen science exist. Projects can be led by experts, community-led, or co-created with different aims and levels of participation ¹⁵³. HBS participants and other citizens generate research questions to be answered with the HBS resource. In traditional designs, scientists test hypotheses that are often based on previous findings within their research domain or their intuitions. However, people living in or with specific conditions (i.e., being in their thirties and going through a key life event) may have additional insight on top of existing expert-knowledge. These insights are uncovered by a citizen science platform. The essence of the platform is to leverage collective intelligence from a large group of participants versus a smaller number of experts. This can reveal topics and research questions that have a significant influence on people's behavior in the real world and their health status, which experts may have left untouched ¹⁵⁴⁻¹⁵⁷. By giving citizens a voice in scientific research, it can contribute considerably to the valorization of research results.

Reflection on design choices and study population

Comprehension of complex interactions as illustrated above requires an interdisciplinary, team science approach ²⁸. The HBS design is the result of an orchestrated cross-campus process over 22 months in which 250 scientists from all classical faculties were involved and were challenged to look past the horizons of their disciplines in a few plenary meetings and several smaller working groups, all providing input to a multidisciplinary scientific board that made the final design decisions. Here, we reflect on our design choices and the study population selected.

First of all, to capture the complexity of the human brain and its environment, a large set of measures was provided. We sought a balance between comprehensiveness, local expertise, costs, and burden for the participants. This resulted in an extensive number, variety, duration of mostly validated assessments, albeit not perfectly comprehensive. For example, the neuroimaging protocol is largely

aligned with the Human Connectome Project ¹⁵⁸ and UK Biobank ¹⁵⁹ brain imaging, but includes a movie fMRI scan that is not included in the Human Connectome Project and UK Biobank, while the latter include scans that are not included in the HBS. Furthermore, instead of continuous monitoring over one year with validated wearable devices, the HBS covers three times a burst week of real-world assessments. Also, the collection of GPS data, financial transactions, and social media interactions were not allowed due to legal restrictions and privacy concerns.

Secondly, the HBS includes three repeated assessments for about one year. These repetitions aim to capture changes in human brain operations that may be related to relevant life events, seasonality, and/or incidental or dynamic changes in the biological and social environment. Regarding seasonality, the HBS participants start at varying time points within a year, so, although we sample only three instead of four times over one year, across participants we sample seasonal transitions in a fine-grained manner.

Thirdly, the HBS aims to include 1,000 participants. Due to differences in measurement-specific signal-to-noise properties, it is not possible to provide a straightforward power and sample size calculation because the data enables analyses of various cognitive, affective, and behavioral interactions in their bio-social context. To decide on the number of participants, we sought a balance between sensitivity and feasibility. The chosen number of participants is high compared to traditional neuroscientific experiments revealing general principles but is low compared to disease risk-oriented cohorts (which is not the aim of the HBS) like the UK Biobank ²⁶, the Rotterdam Study ²⁴, or the Rhineland Study ²⁵. However, the number is comparable to other studies designed to capture inter-individual differences like the Human Connectome Project, which included 1,200 young healthy adults ¹⁵⁸, or the Personalized Parkinson Project, which included 650 patients ¹⁶⁰. We consider the number large given the comprehensive range of repeated measures both in the laboratory as in the real world.

Fourthly, we developed a recruitment strategy targeted at a sample that represents the 30-39-year-old population of Nijmegen and its surroundings in terms of gender and educational attainment. However, a reasonable level of reading, speaking, and understanding Dutch (B1 level) is required to be able to complete the study protocol, e.g., to fill out questionnaires. This implies that the HBS participants do not fully represent the Nijmegen population at large, because in this example the illiterate, people with low literacy, or non-Dutch speaking individuals are excluded. However, the aim of including 220 participants with a low, 340 with a middle, and 430 with a high level of education enables the study of interacting social factors.

Digital security system and data sharing

The HBS resource will be accessible to the scientific community at large. The resource contains sensitive personal data that needs to be protected from unauthorized access and unintentional disclosure. The sharing of (big) data within the scientific community is necessary for progress and maximizes scientific benefits derived from valuable and costly data. The HBS data is protected by a digital security system, a Polymorphic Encryption and Pseudonymization (PEP) infrastructure ³³, which allows the sharing of data with researchers worldwide while safeguarding participants' privacy in line with the European General Data Protection Regulation. The digital security system is based on a multipoint, privacy-by-design strategy: (a) participants provide informed consent, also for the important element of data sharing; (b) signed contractual agreements with researchers are in place to ensure that no attempts towards de-pseudonymization, linking or commercialization of the raw data will be attempted; (c) governance policies limit access to the data to qualified researchers only; (d) an innovative pseudonymization and encryption process is applied ³⁴.

An access procedure is in place and published on https://www.healthybrainstudy.nl/en/data-and-methods. We stratify researchers into three tiers with different rights. Tier I consists of researchers from the Radboud campus that contributed to study design or data acquisition. Tier II consists of all

other researchers from the Radboud campus. Tier III consists of publicly financed researchers from other academic institutions. Companies can apply in all tiers, but they cannot apply independently. Application for data starts with the submission of a data request for a project that has been preregistered, e.g., in the Open Science Framework. Then, the HBS scientific board reviews the application. After approval, the researcher signs a data/material transfer agreement. Next, the researcher receives data and/or samples. The Radboud Biobank provides the samples ³². All processed data and samples with relevant documentation (including scripts and data and/or samples processing protocols) must be integrated back into the HBS resource so that it can be used by others. Finally, the researcher publishes the results by acknowledging the HBS consortium.

Conclusion

The HBS has been designed using a team science approach to integrate scientific disciplines and is characterized by a broad range of repeated assessments, a large number of participants, both laboratory and real-world assessments, and a population-based sample. Moreover, data is managed to allow data sharing with scientists worldwide while maintaining participants' privacy. With the HBS resource, the scientific community can take the next step in understanding the human brain and how it dynamically and individually operates in its bio-social context.

References

- 428 1 G-Science Academies Statement. Understanding, protecting, and developing global brain resources: https://royalsociety.org/about-us/international/international-work/g-science-academies-meetings/. (2016).
- Bruineberg, J., Kiverstein, J. & Rietveld, E. The anticipating brain is not a scientist: the freeenergy principle from an ecological-enactive perspective. *Synthese* **195**, 2417-2444 (2018).
- 433 3 Kiverstein, J. & Miller, M. The embodied brain: towards a radical embodied cognitive neuroscience. *Frontiers in Human Neuroscience* **9** (2015).
- 435 4 Clark, A. Being there: putting mind, body, and world together again. (1997).
- 436 5 Chiel, H. J. & Beer, R. D. The brain has a body: adaptive behavior emerges from interactions of nervous system, body and environment. *Trends Neurosci* **20**, 553-557 (1997).
- Hermans, E. J. *et al.* Stress-related noradrenergic activity prompts large-scale neural network reconfiguration. *Science* **334**, 1151-1153 (2011).
- Haeger, A., Costa, A. S., Schulz, J. B. & Reetz, K. Cerebral changes improved by physical activity during cognitive decline: A systematic review on MRI studies. *NeuroImage: Clinical* **23**, 101933 (2019).
- 443 8 Lyall, L. M. *et al.* Seasonality of depressive symptoms in women but not in men: A cross-444 sectional study in the UK Biobank cohort. *J Affect Disord* **229**, 296-305 (2018).
- Winthorst, W. H. *et al.* Self-attributed seasonality of mood and behavior: a report from the Netherlands study of depression and anxiety. *Depress Anxiety* **31**, 517-523 (2014).
- 447 10 Cepeda, M. *et al.* Seasonality of physical activity, sedentary behavior, and sleep in a middle-448 aged and elderly population: The Rotterdam study. *Maturitas* **110**, 41-50 (2018).
- 449 11 Miller, R. *et al.* The CIRCORT database: Reference ranges and seasonal changes in diurnal salivary cortisol derived from a meta-dataset comprised of 15 field studies.

 450 *Psychoneuroendocrinology* **73**, 16-23 (2016).
- Davenport, E. R. *et al.* Seasonal variation in human gut microbiome composition. *PLoS One* **9**, e90731 (2014).
- Ockene, I. S. *et al.* Seasonal variation in serum cholesterol levels: treatment implications and possible mechanisms. *Arch Intern Med* **164**, 863-870 (2004).
- 456 14 Marti-Soler, H. *et al.* Seasonality of cardiovascular risk factors: an analysis including over 230 000 participants in 15 countries. *Heart* **100**, 1517-1523 (2014).
- Llera, A., Wolfers, T., Mulders, P. & Beckmann, C. F. Inter-individual differences in human brain structure and morphology link to variation in demographics and behavior. *eLife* **8**, e44443 (2019).
- Smith, S. M. *et al.* A positive-negative mode of population covariation links brain connectivity, demographics and behavior. *Nat Neurosci* **18**, 1565-1567 (2015).
- 463 17 Myin-Germeys, I. *et al.* Experience sampling methodology in mental health research: new insights and technical developments. *World Psychiatry* **17**, 123-132 (2018).
- Hogenelst, K., Schoevers, R. A. & aan het Rot, M. Studying the neurobiology of human social interaction: Making the case for ecological validity. *Social Neuroscience* **10**, 219-229 (2015).
- Fry, A. *et al.* Comparison of sociodemographic and health-related characteristics of UK
 Biobank participants with those of the general population. *Am J Epidemiol* **186**, 1026-1034
 (2017).
- 470 20 Henrich, J., Heine, S. J. & Norenzayan, A. The weirdest people in the world? *The Behavioral and brain sciences* **33**, 61-83; discussion 83-135 (2010).
- Jaddoe, V. W. V. *et al.* The Generation R Study: Design and cohort profile. *European Journal of Epidemiology* **21**, 475 (2006).
- Volkow, N. D. *et al.* The conception of the ABCD study: From substance use to a broad NIH collaboration. *Developmental Cognitive Neuroscience* **32**, 4-7 (2018).
- Niermann, H. C. M. *et al.* The relation between infant freezing and the development of internalizing symptoms in adolescence: A prospective longitudinal study. *Dev Sci* **22**, e12763 (2019).
- Hofman, A. *et al.* The Rotterdam Study: objectives and design update. *Eur J Epidemiol* **22**, 819-829 (2007).
- Breteler, M. & Wolf, H. The Rhineland study: a novel platform for epidemiologic research into Alzeimer disease and related disorders. *Alzheimer's & Dementia* **10** (2014).

- Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* **12**, e1001779 (2015).
- Sowell, E. R. *et al.* Mapping cortical change across the human life span. *Nat Neurosci* **6**, 309-315 (2003).
- 487 28 Utzerath, C. & Fernández, G. Shaping science for increasing interdependence and specialization. *Trends in Neurosciences* **40**, 121-124 (2017).
- 489 29 Statistics Netherlands: https://www.cbs.nl
- 490 30 Association, W. M. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* **310**, 2191-2194 (2013).
- 492 31 https://stichtingcrowdience.nl/.
- 493 32 Manders, P., Lutomski, J., Smit, C., Swinkels, D. & Zielhuis, G. The Radboud Biobank: A central facility for disease-based biobanks to optimise use and distribution of biomaterial for scientific research in the Radboud University Medical Center, Nijmegen. *Open Journal of Bioresources* **5** (2018).
- Verheul, E. R., Jacobs, B., Meijer, C., Hildebrandt, M. & de Ruiter, J. Polymorphic encryption and pseudonymisation for personalised healthcare. *IACR Cryptology ePrint Archive*, 411 (2016).
- Jacobs, B. & Popma, J. Medical research, big data and the need for privacy by design. *Big Data* & Society **6**, 2053951718824352 (2019).
- 502 35 Spielberger, C. D., Gorsuch, R. L., Lushene, R., Vagg, P. R. & Jacobs, G. A. State-Trait Anxiety Inventory for adults -manual. (1983).
- 504 36 Cohen, S., Kamarck, T. & Mermelstein, R. A global measure of perceived stress. *J Health Soc Behav* **24**, 385-396 (1983).
- 506 37 Bjørnskov, C. How comparable are the Gallup world poll life satisfaction data? *Journal of Happiness Studies* **11**, 41-60 (2010).
- Russell, D., Peplau, L. A. & Cutrona, C. E. The revised UCLA Loneliness Scale: concurrent and discriminant validity evidence. *J Pers Soc Psychol* **39**, 472-480 (1980).
- Rosenberg, D. E. *et al.* Reliability and validity of the Sedentary Behavior Questionnaire (SBQ) for adults. *J Phys Act Health* **7**, 697-705 (2010).
- 512 40 Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R. & Kupfer, D. J. The Pittsburgh Sleep 513 Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* **28**, 193-514 213 (1989).
- 515 41 https://www.lissdata.nl/.
- Vink, J. M., Willemsen, G., Beem, A. L. & Boomsma, D. I. The Fagerstrom test for nicotine dependence in a Dutch sample of daily smokers and ex-smokers. *Addict Behav* **30**, 575-579 (2005).
- 519 43 Saunders, J. B., Aasland, O. G., Babor, T. F., de la Fuente, J. R. & Grant, M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption. *Addiction* **88**, 791-804 (1993).
- 522 44 Schippers, G. M. & Broekman, T. G. MATE-Q 2.1. Handleiding. Nijmegen: Bêta Boeken (2014).
- Feunekes, G. I., Van Staveren, W. A., De Vries, J. H., Burema, J. & Hautvast, J. G. Relative and biomarker-based validity of a food-frequency questionnaire estimating intake of fats and cholesterol. *Am J Clin Nutr* **58**, 489-496 (1993).
- Molag, M. L. *et al.* Selecting informative food items for compiling food-frequency questionnaires: comparison of procedures. *Br J Nutr* **104**, 446-456 (2010).
- 528 47 Siebelink, E., Geelen, A. & de Vries, J. H. Self-reported energy intake by FFQ compared with actual energy intake to maintain body weight in 516 adults. *Br J Nutr* **106**, 274-281 (2011).
- Verkleij-Hagoort, A. C. *et al.* Validation of the assessment of folate and vitamin B12 intake in women of reproductive age: the method of triads. *Eur J Clin Nutr* **61**, 610-615 (2007).
- 532 49 Schredl, M. Reliability and stability of a dream recall frequency scale. *Percept Mot Skills* **98**, 1422-1426 (2004).
- 534 50 Lemmens, J. S., Valkenburg, P. M. & Gentile, D. A. The internet gaming disorder scale. *Psychol Assess* **27**, 567-582 (2015).
- van den Eijnden, R. J. J. M., Lemmens, J. S. & Valkenburg, P. M. The social media disorder scale. *Comput Hum Behav* **61**, 478-487 (2016).
- 538 52 Baumgartner, S. E., Lemmens, J. S., Weeda, W. D. & Huizinga, M. Measuring media 539 multitasking: development of a short measure of media multitasking for adolescents. *ournal* 540 *of Media Psychology: Theories, Methods, and Applications* **92**, 92-101 (2017).

- 541 53 Diener, E., Emmons, R. A., Larsen, R. J. & Griffin, S. The satisfaction with life scale. *J Pers* 4ssess **49**, 71-75 (1985).
- 543 54 Bohlmeijer, E., ten Klooster, P. M., Fledderus, M., Veehof, M. & Baer, R. Psychometric 544 properties of the five facet mindfulness questionnaire in depressed adults and development 545 of a short form. *Assessment* **18**, 308-320 (2011).
- van Dijk, L., van der Maat, M., Salimans, R. & Bouvy, M. De balans tussen verkrijgbaarheid en veiligheid. Evaluatie van de indeling van zelfzorggeneesmiddelen en de rol van drogist en apotheek bij de verstrekking. *Nivel* (2010).
- 549 56 Yzermans, J., Baliatsas, C., van Dulmen, S. & Van Kamp, I. Assessing non-specific symptoms in 550 epidemiological studies: Development and validation of the Symptoms and Perceptions (SaP) 551 questionnaire. *International Journal of Hygiene and Environmental Health* **219**, 53-65 (2016).
- 552 57 Kessler, R. C. *et al.* The World Health Organization adult ADHD Self-Report Scale (ASRS): a short screening scale for use in the general population. *Psychol Med* **35**, 245-256 (2005).
- 554 58 Bralten, J. *et al.* Autism spectrum disorders and autistic traits share genetics and biology. *Mol Psychiatry* **23**, 1205-1212 (2018).
- Rush, A. J., Gullion, C. M., Basco, M. R., Jarrett, R. B. & Trivedi, M. H. The Inventory of Depressive Symptomatology (IDS): psychometric properties. *Psychol Med* **26**, 477-486 (1996).
- Rodriguez, B. F., Bruce, S. E., Pagano, M. E., Spencer, M. A. & Keller, M. B. Factor structure and stability of the Anxiety Sensitivity Index in a longitudinal study of anxiety disorder patients. *Behav Res Ther* **42**, 79-91 (2004).
- 561 61 Schaufeli, W. & Dierendonck, D. v. Utrechtse burn-out schaal: handleiding. *Amsterdam: Pearson Assessment and Information B.V.* (1981, 2001).
- 563 62 Cima, M., Raine, A., Meesters, C. & Popma, A. Validation of the Dutch Reactive Proactive Questionnaire (RPQ): differential correlates of reactive and proactive aggression from childhood to adulthood. *Aggress Behav* **39**, 99-113 (2013).
- Kanner, A. D., Coyne, J. C., Schaefer, C. & Lazarus, R. S. Comparison of two modes of stress measurement: daily hassles and uplifts versus major life events. *J Behav Med* **4**, 1-39 (1981).
- Garnefski, N., Kraaij, V. & Spinhoven, P. CERQ. Handleiding voor het gebruik van de Cognitive Emotion Regulation Questionnaire. *Leiderdorp: DATEC.* (2002).
- 570 65 Bernstein, D. P. *et al.* Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl* **27**, 169-190 (2003).
- 572 66 Brugha, T. S. & Cragg, D. The list of threatening experiences: the reliability and validity of a brief life events questionnaire. *Acta Psychiatr Scand* **82**, 77-81 (1990).
- 574 67 Leary, M. R. Need to belong scale. Measurement instrument database for the social science. 575 <u>www.midss.ie</u> (2013).
- Zimet, G. D., Powell, S. S., Farley, G. K., Werkman, S. & Berkoff, K. A. Psychometric
 characteristics of the Multidimensional Scale of Perceived Social Support. *J Pers Assess* 55, 610-617 (1990).
- Geurts, S. *et al.* Work-home interaction from a work psychological perspective: Development and validation of a new questionnaire, the SWING. *Work & Stress* **19** (2005).
- Klein, H. J., Cooper, J. T., Molloy, J. C. & Swanson, J. A. The assessment of commitment:
 Advantages of a unidimensional, target-free approach. *Journal of Applied Psychology* **99**, 222-238 (2014).
- van der Heijden, B. I. J. M. *et al.* Development and validation of the short-form employability five-factor instrument. *Journal of Vocational Behavior* **106**, 236 248 (2018).
- van der Heijden, B. I. Professional expertise of higher level employees; age stereotyping in self-assessments and supervisor ratings. *Tijdschr Gerontol Geriatr* **31**, 62-69 (2000).
- van Veldhoven, M. J. P. M., Prins, J., van der Laken, P. A. & Dijkstra, L. BBA2.0: Update van de standaard voor vragenlijstonderzoek naar werk, welbevinden en prestaties. *Amsterdam: SKB* (2014).
- 591 74 Craig, S. C. Efficacy, trust, and political behavior: an attempt to resolve a lingering conceptual dilemma. *American Politics Quarterly* **7**, 225-239 (1979).
- Kuhn, T., van Elsas, E., Hakhverdian, A. & van der Brug, W. An ever wider gap in an ever closer
 union: Rising inequalities and euroscepticism in 12 West European democracies, 1975–2009.
 Socio-Economic Review 14, 27-45 (2016).
- 596 76 Costa, P. & R. McCrae, R. The revised NEO personality inventory (NEO-PI-R). Vol. 2 (2008).
- 597 77 Aron, E. N., Aron, A. & Jagiellowicz, J. Sensory processing sensitivity: a review in the light of the evolution of biological responsivity. *Pers Soc Psychol Rev* **16**, 262-282 (2012).

- Patton, J. H., Stanford, M. S. & Barratt, E. S. Factor structure of the Barratt impulsiveness scale. *J Clin Psychol* **51**, 768-774 (1995).
- Tangney, J. P., Baumeister, R. F. & Boone, A. L. High self-control predicts good adjustment, less pathology, better grades, and interpersonal success. *J Pers* **72**, 271-324 (2004).
- 603 80 Chen, G., Gully, S. M. & Eden, D. Validation of a new general self-efficacy scale. *Organizational Research Methods* **4**, 62-83 (2001).
- Seuntjens, T. G., Zeelenberg, M., van de Ven, N. & Breugelmans, S. M. Dispositional greed. *J Pers Soc Psychol* **108**, 917-933 (2015).
- Jonason, P. K. & Webster, G. D. The dirty dozen: a concise measure of the dark triad. *Psychol Assess* **22**, 420-432 (2010).
- Williams, G. Some determinants of the socially responsible investment decision: A crosscountry study. *Journal of Behavioral Finance* **8**, 43-57 (2007).
- Weller, J. A. *et al.* Development and testing of an abbreviated numeracy scale: A rasch analysis approach. *Journal of Behavioral Decision Making* **26**, 198-212 (2013).
- Atkinson, A. & Messy, F.-A. Assessing financial literacy in 12 countries: an OECD/INFE international pilot exercise. *Journal of Pension Economics and Finance* **10**, 657-665 (2011).
- 615 86 Galesic, M. & Garcia-Retamero, R. Graph literacy: A cross-cultural comparison. *Medical decision making* **31**, 444-457 (2011).
- 617 87 Thomas, D. et al. Cultural intelligence: A theory-based, short form measure. (2015).
- Astrand, P. O. & Ryhming, I. A nomogram for calculation of aerobic capacity (physical fitness) from pulse rate during sub-maximal work. *J Appl Physiol* **7**, 218-221 (1954).
- Kozey-Keadle, S., Libertine, A., Lyden, K., Staudenmayer, J. & Freedson, P. S. Validation of wearable monitors for assessing sedentary behavior. *Med Sci Sports Exerc* **43**, 1561-1567 (2011).
- Ryan, C. G., Grant, P. M., Tigbe, W. W. & Granat, M. H. The validity and reliability of a novel activity monitor as a measure of walking. *British journal of sports medicine* **40**, 779-784 (2006).
- 626 91 Garbarino, M., Lai, M., Bender, D., Picard, R. & Tognetti, S. Empatica E3 A wearable wireless 627 multi-sensor device for real-time computerized biofeedback and data acquisition. 39-42 628 (2015).
- 629 92 Grillon, C., Ameli, R., Woods, S. W., Merikangas, K. & Davis, M. Fear-potentiated startle in 630 humans: effects of anticipatory anxiety on the acoustic blink reflex. *Psychophysiology* **28**, 588-631 595 (1991).
- Aitken, R. C. Measurement of feelings using visual analogue scales. *Proc R Soc Med* **62**, 989-993 (1969).
- Shanholtzer, B. A. & Patterson, S. M. Use of bioelectrical impedance in hydration status assessment: reliability of a new tool in psychophysiology research. *Int J Psychophysiol* **49**, 217-226 (2003).
- 637 95 Collins, S. L., Moore, R. A. & McQuay, H. J. The visual analogue pain intensity scale: what is moderate pain in millimetres? *Pain* **72**, 95-97 (1997).
- Timmerman, H., Wilder-Smith, O., van Weel, C., Wolff, A. & Vissers, K. Detecting the neuropathic pain component in the clinical setting: a study protocol for validation of screening instruments for the presence of a neuropathic pain component. *BMC Neurol* 14, 94 (2014).
- Timmerman, H., Wilder-Smith, O. H., Steegers, M. A., Vissers, K. C. & Wolff, A. P. The added value of bedside examination and screening QST to improve neuropathic pain identification in patients with chronic pain. *J Pain Res* **11**, 1307-1318 (2018).
- Lanier, G. M. *et al.* Validity and reliability of a novel slow cuff-deflation system for noninvasive
 blood pressure monitoring in patients with continuous-flow left ventricular assist device. *Circ Heart Fail* 6, 1005-1012 (2013).
- van Mil, A. C. *et al.* Correlation of carotid artery reactivity with cardiovascular risk factors and
 coronary artery vasodilator responses in asymptomatic, healthy volunteers. *J Hypertens* 35,
 1026-1034 (2017).
- Faith, J. J. *et al.* The long-term stability of the human gut microbiota. *Science* **341**, 1237439 (2013).
- 654 101 Cherbuin, N., Kumar, R., Sachdev, P. S. & Anstey, K. J. Dietary mineral intake and risk of mild cognitive impairment: the PATH through life project. *Front Aging Neurosci* **6**, 4 (2014).

- Schwabe, L., Haddad, L. & Schachinger, H. HPA axis activation by a socially evaluated coldpressor test. *Psychoneuroendocrinology* **33**, 890-895 (2008).
- Wright, K. D., Hickman, R. & Laudenslager, M. L. Hair cortisol analysis: A promising biomarker of HPA activation in older adults. *Gerontologist* **55 Suppl 1**, S140-145 (2015).
- Aerts, R. *et al.* Silicone wristband passive samplers yield highly individualized pesticide residue exposure profiles. *Environmental Science & Technology* **52**, 298-307 (2018).
- 662 105 O'Connell, S. G., Kincl, L. D. & Anderson, K. A. Silicone wristbands as personal passive samplers. *Environmental Science & Technology* **48**, 3327-3335 (2014).
- Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome Project.

 Neuroimage **80**, 105-124 (2013).
- Diaz, B. A. *et al.* The ARSQ 2.0 reveals age and personality effects on mind-wandering experiences. *Front Psychol* **5**, 271 (2014).
- Marques, J. P. *et al.* MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. *Neuroimage* **49**, 1271-1281 (2010).
- Langkammer, C. *et al.* Quantitative susceptibility mapping (QSM) as a means to measure brain iron? A post mortem validation study. *Neuroimage* **62**, 1593-1599 (2012).
- 672 110 Constantino, S. M. & Daw, N. D. Learning the opportunity cost of time in a patch-foraging task. *Cogn Affect Behav Neurosci* **15**, 837-853 (2015).
- Braun, A., Urai, A. E. & Donner, T. H. Adaptive history biases result from confidence-weighted accumulation of past choices. *J Neurosci* (2018).
- Therrien, A. S., Wolpert, D. M. & Bastian, A. J. Effective reinforcement learning following cerebellar damage requires a balance between exploration and motor noise. *Brain* **139**, 101-114 (2016).
- Persson, J., Kalpouzos, G., Nilsson, L. G., Ryberg, M. & Nyberg, L. Preserved hippocampus activation in normal aging as revealed by fMRI. *Hippocampus* **21**, 753-766 (2011).
- Oosterman, J. M., Wijers, M. & Kessels, R. P. Planning or something else? Examining neuropsychological predictors of Zoo Map performance. *Appl Neuropsychol Adult* **20**, 103-109 (2013).
- Andreatta, M. *et al.* Initial and sustained brain responses to contextual conditioned anxiety in humans. *Cortex* **63**, 352-363 (2015).
- Webb, T. L., Miles, E. & Sheeran, P. Dealing with feeling: a meta-analysis of the effectiveness of strategies derived from the process model of emotion regulation. *Psychol Bull* **138**, 775-808 (2012).
- Derry, P. A. & Kuiper, N. A. Schematic processing and self-reference in clinical depression. *J Abnorm Psychol* **90**, 286-297 (1981).
- 691 118 Groefsema, M., Engels, R., Kuntsche, E., Smit, K. & Luijten, M. Cognitive biases for social alcohol-related pictures and alcohol use in specific social settings: an event-level study.

 693 Alcohol Clin Exp Res 40, 2001-2010 (2016).
- Figner, B., Mackinlay, R. J., Wilkening, F. & Weber, E. U. Affective and deliberative processes
 in risky choice: age differences in risk taking in the Columbia Card Task. *J Exp Psychol Learn Mem Cogn* 35, 709-730 (2009).
- Veling, H. *et al.* Training impulsive choices for healthy and sustainable food. *J Exp Psychol Appl* **23**, 204-215 (2017).
- Noussair, C. N., Trautmann, S. T. & van de Kuilen, G. Higher order risk attitudes,
 demographics, and financial decisions. *The Review of Economic Studies* 81, 325-355 (2014).
- 701 122 Kerschbamer, R. The geometry of distributional preferences and a non-parametric
 702 identification approach: The Equality Equivalence Test. *European Economic Review* 76, 85-103
 703 (2015).
- 704 123 Dimmock, S. G., Kouwenberg, R. & Wakker, P. P. Ambiguity attitudes in a large representative sample. *Management Science* **62**, 1363-1380 (2016).
- Hergueux, J. & Jacquemet, N. Social preferences in the online laboratory: a randomized experiment. *Experimental Economics* **18**, 251-283 (2015).
- Fischbacher, U., Gächter, S. & Quercia, S. The behavioral validity of the strategy method in public good experiments. *Journal of Economic Psychology* **33**, 897-913 (2012).
- 710 126 Falk, A. *et al.* Global evidence on economic preferences. *The Quarterly Journal of Economics* 133, 1645-1692 (2018).

- 712 127 Kuijpers, M., Hakemulder, F., Tan, E. & Doicaru, M. Exploring absorbing reading experiences: 713 Developing and validating a self-report scale to measure story world absorption. *Scientific* 714 *Study of Literature* **4** (2014).
- Jasso, G. & Opp, K. D. Probing the character of norms: A factorial survey analysis of the norms of political action. *American Sociological Review* **62**, 947 (1997).
- 717 129 Rooks, G., Raub, W., Selten, R. & Tazelaar, F. How inter-firm co-operation depends on social embeddedness: A vignette study. *Acta Sociologica* **43**, 123-137 (2000).
- Hintz, F. et al. Shared lexical access processes in speaking and listening? An individual
 differences study. Journal of Experimental Psychology: Learning, Memory, and Cognition 46,
 1048–1063 (2020).
- 722 131 Salthouse, T. A. The processing-speed theory of adult age differences in cognition. *Psychol Rev* 103, 403-428 (1996).
- T24 132 Earles, J. L. & Salthouse, T. A. Interrelations of age, health, and speed. *J Gerontol B Psychol Sci Soc Sci* **50**, P33-p41 (1995).
- 726 133 Deary, I. J., Liewald, D. & Nissan, J. A free, easy-to-use, computer-based simple and four-727 choice reaction time programme: the Deary-Liewald reaction time task. *Behav Res Methods* 728 **43**, 258-268 (2011).
- 729 134 Wechsler, D. WAIS-III (3rd edition). Amsterdam: Harcourt Test Publishers (2004).
- 730 135 Chu, M., Meyer, A., Foulkes, L. & Kita, S. Individual differences in frequency and saliency of speech-accompanying gestures: the role of cognitive abilities and empathy. *J Exp Psychol Gen* 143, 694-709 (2014).
- 733 136 Berch, D. B., Krikorian, R. & Huha, E. M. The Corsi block-tapping task: methodological and theoretical considerations. *Brain Cogn* **38**, 317-338 (1998).
- 735 137 Raven, J., Raven, J. C. & Court, J. H. Raven manual section 4: advanced progressive matrices. 736 Oxford, UK: Oxford Psychologists Press (1998).
- 737 138 Dunn, L. M. & Dunn, D. Peabody Picture Vocabulary Test (3rd Edition). *Circle Pines: American Guidance Service* (1997).
- 739 139 Schlichting, L. Peabody Picture Vocabulary Test Dutch-III-NL. *Amsterdam, NL: Harcourt* 740 *Assessment BV* (2005).
- 741 140 Brysbaert, M., Sui, L., Dirix, N. & Hintz, F. Dutch Author Recognition Test. J Cogn 3, 6-6 (2020).
- Hubers, F., Snijders, T. & Hoop, H. How the brain processes violations of the grammatical norm: An fMRI study. *Brain and Language* **163**, 22-31 (2016).
- 744 142 Shao, Z., Janse, E., Visser, K. & Meyer, A. S. What do verbal fluency tasks measure? Predictors of verbal fluency performance in older adults. *Frontiers in psychology* **5**, 772-772 (2014).
- 746 143 Mainz, N., Shao, Z., Brysbaert, M. & Meyer, A. S. Vocabulary knowledge predicts lexical 747 processing: Evidence from a group of participants with diverse educational backgrounds. 748 *Front Psychol* **8**, 1164 (2017).
- Jongman, S., Khoe, Y. & Hintz, F. Vocabulary size influences spontaneous speech in native language users: validating the use of automatic speech recognition in individual differences research. *Language and Speech*, 002383092091107 (2020).
- Huettig, F. & Janse, E. Individual differences in working memory and processing speed predict anticipatory spoken language processing in the visual world. *Language, Cognition and Neuroscience* **31**, 80-93 (2016).
- 755 146 Altmann, G. T. M. & Kamide, Y. Incremental interpretation at verbs: restricting the domain of subsequent reference. *Cognition* **73**, 247-264 (1999).
- Hintz, F., Meyer, A. S. & Huettig, F. Predictors of verb-mediated anticipatory eye movements in the visual world. *J Exp Psychol Learn Mem Cogn* **43**, 1352-1374 (2017).
- Kahneman, D. & Deaton, A. High income improves evaluation of life but not emotional well-being. *Proc Natl Acad Sci U S A* **107**, 16489-16493 (2010).
- 761 149 Biswas, A. *et al.* Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med* **162**, 123-132 (2015).
- Ekelund, U. et al. Does physical activity attenuate, or even eliminate, the detrimental
 association of sitting time with mortality? A harmonised meta-analysis of data from more
 than 1 million men and women. The Lancet 388, 1302-1310 (2016).
- 767 151 Diaz, K. M. *et al.* Patterns of sedentary behavior and mortality in U.S. middle-aged and older adults: a national cohort study. *Ann Intern Med* **167**, 465-475 (2017).

- ten Broeke, P. *et al.* Temporal dynamics of sitting behavior at work. *Proceedings of the National Academy of Sciences* **117**, 14883-14889 (2020).
- den Broeder, L., Devilee, J., van Oers, H., Schuit, A. J. & Wagemakers, A. Citizen science for public health. *Health Promotion International* **33**, 505-514 (2016).
- 773 154 Bevelander, K. E. *et al.* Crowdsourcing novel childhood predictors of adult obesity. *PLOS ONE* **9**, e87756 (2014).
- 775 155 Bongard, J., Hines, P., Conger, D., Hurd, P. & Lu, Z. Crowdsourcing predictors of behavioral outcomes. *IEEE Transactions on Systems, Man, and Cybernetics Part A:Systems and Humans* 43 (2012).
- 778 156 Wang, C. *et al.* Crowdsourcing in health and medical research: a systematic review. *Infect Dis Poverty* **9**, 8 (2020).
- 780 157 Mount, M., Round, H. & Pitsis, T. S. Design thinking inspired crowdsourcing: Toward a 781 generative model of complex problem solving. *California Management Review* **62**, 103-120 782 (2020).
- 783 158 Van Essen, D. C. *et al.* The Human Connectome Project: a data acquisition perspective. *Neuroimage* **62**, 2222-2231 (2012).
- 785 159 Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective epidemiological study. *Nat Neurosci* **19**, 1523-1536 (2016).

787 160 Bloem, B. R. *et al.* The Personalized Parkinson Project: examining disease progression through broad biomarkers in early Parkinson's disease. *BMC Neurol* **19**, 160 (2019).

Acknowledgement

791 The HBS is funded by the Reinier Post Foundation and Radboud University, Nijmegen, the

792 Netherlands.

790

793 The HBS consortium consists of:

Last name	First name	Email address	Affiliation
Aarts	Esther	esther.aarts@donders.ru.nl	1
Akkerman	Agnes	a.akkerman@fm.ru.nl	2
Altgassen	Mareike	aaltgass@uni-mainz.de	3
Bartels	Ronald	ronald.bartels@radboudumc.nl	4
Beckers	Debby	d.beckers@psych.ru.nl	5
Bevelander	Kirsten	kris.bevelander@radboudumc.nl	4
Bijleveld	Erik	e.bijleveld@psych.ru.nl	5
Blaney Davidson	Esmeralda	esmeral da. blaney da vids on @radboudum c.nl	4
Boleij	Annemarie	annemarie.boleij@radboudumc.nl	4
Bralten	Janita	janita.bralten@radboudumc.nl	4
Cillessen	Toon	a.cillessen@psych.ru.nl	5
Claassen	Jurgen	jurgen.claassen@radboudumc.nl	4
Cools	Roshan	r.cools@donders.ru.nl	6
Cornelissen	Ineke	ineke.cornelissen@radboudumc.nl	4
Dresler	Martin	martin.dresler@donders.ru.nl	6
Eijsvogels	Thijs	thijs.eijsvogels@radboudumc.nl	4
Faber	Myrthe	m.faber@donders.ru.nl	6
Fernández	Guillén	g.fernandez@donders.ru.nl	6
Figner	Bernd	b.figner@psych.ru.nl	1-5
Fritsche	Matthias	m.fritsche@donders.ru.nl	1
Füllbrunn	Sascha	s.fullbrunn@fm.ru.nl	2
Gayet	Surya	s.gayet@donders.ru.nl	1
Gelder, van	Marleen M.H.J.	marleen.vangelder@radboudumc.nl	4
Gerven, van	Marcel	marcel.vangerven@donders.ru.nl	1
Geurts	Sabine	s.geurts@psych.ru.nl	5
Greven	Corina U.	c.greven@donders.ru.nl	6
Groefsema	Martine	m.groefsema@pwo.ru.nl	5
Haak	Koen	k.haak@donders.ru.nl	6
Hagoort	Peter	peter.hagoort@mpi.nl	1-7
Hartman	Yvonne	yvonne.hartman@radboudumc.nl	4

Heijden, van der	Beatrice	b.vanderheijden@fm.ru.nl	2
Hermans	Erno	erno.hermans@donders.ru.nl	6
Heuvelmans	Vivian	vivian.heuvelmans@radboudumc.nl	4
Hintz	Florian	florian.hintz@mpi.nl	7
Hollander, den	Janet	janet.denhollander@radboudumc.nl	4
Hulsman	Anneloes M.	a.hulsman@donders.ru.nl	1-5
Idesis	Sebastian	sebastian.idesis@upf.edu	8
Jaeger	Martin	martin.jaeger@radboudumc.nl	4
Janse	Esther	e.janse@let.ru.nl	9
Janzing	Joost	joost.janzing@radboudumc.nl	4
Kessels	Roy P.C.	r.kessels@donders.ru.nl	1-6
Kleijn, de	Willemien	w.dekleijn@psych.ru.nl	10
Klein	Marieke	marieke.klein@radboudumc.nl	4
Klumpers	Floris	f.klumpers@donders.rul.nl	1-5
Kohn	Nils	n.kohn@donders.ru.nl	6
Korzilius	Hubert	h.korzilius@fm.ru.nl	2
Krahmer	Bas	b1.krahmer@student.ru.nl	4
Lange, de	Floris	f.delange@donders.ru.nl	1
Leeuwen, van	Judith	j.vanleeuwen@donders.ru.nl	6
Liu	Huaiyu	l.huaiyu@psych.ru.nl	5
Luijten	Maartje	m.luijten@bsi.ru.nl	5
Manders	Peggy	peggy.manders@radboudumc.nl	4
Manevska	Katerina	k.manevska@fm.ru.nl	2
Marques	José P.	j.marques@donders.ru.nl	1
Matthews	Jon	jon.matthews@radboudumc.nl	6
McQueen	James M.	j.mcqueen@donders.ru.nl	1
Medendorp	Pieter	p.medendorp@donders.ru.nl	1
Melis	René	Rene. Melis@radboudumc.nl	4
Meyer	Antje	antje.meyer@mpi.nl	7
Oosterman	Joukje	j.oosterman@donders.ru.nl	1
Overbeek	Lucy	lucy.overbeek@radboudumc.nl	4
Peelen	Marius	m.peelen@donders.ru.nl	1
Popma	Jean	j.popma@ru.nl	11
Postma	Geert	g.j.postma@science.ru.nl	12
Roelofs	Karin	k.roelofs@donders.ru.nl	1-5
Rossenberg, van	Yvonne G.T.	y.vanrossenberg@fm.ru.nl	2
Schaap	Gabi	g.schaap@maw.ru.nl	5

Scheepers	Paul	paul.scheepers@radboudumc.nl	4
Selen	Luc	l.selen@donders.ru.nl	1
Starren	Marianne	m.starren@let.ru.nl	9
Swinkels	Dorine W.	dorine.swinkels@radboudumc.nl	4
Tendolkar	Indira	indira.tendolkar@radboudumc.nl	6
Thijssen	Dick	dick.thijssen@radboudumc.nl	4
Timmerman	Hans	h.timmerman02@umcg.nl	13
Tutunji	Rayyan	r.toutounji@donders.ru.nl	6
Tuladhar	Anil	anil.tuladhar@radboudumc.nl	6
Veling	Harm	h.veling@psych.ru.nl	5
Verhagen	Maaike	m.verhagen@pwo.ru.nl	5
Verkroost	Jasper	jasper.verkroost@radboudumc.nl	4
Vriezekolk	Vivian	vivian.vriezekolk@radboudumc.nl	4
Vrijsen	Janna	janna.vrijsen@radboudumc.nl	6
Vyrastekova	Jana	j.vyrastekova@fm.ru.nl	2
Wal, van der	Selina	selina.vanderWal@radboudumc.nl	4
Willems	Roel	r.willems@donders.ru.nl	1-9
Willemsen	Arthur	arthur.willemsen@radboudumc.nl	4

Affiliations:

Donders Institute for Brain, Cognition and Behavior, Radboud University, Nijmegen, The Netherlands Insitute for Management Research, Radboud University, Nijmegen, The Netherlands Johannes Gutenberg-University Mainz, Germany Radboud University Medical Center, Nijmegen, The Netherlands Behavioural Science Institute, Radboud University, Nijmegen, The Netherlands Donders Institute for Brain, Cognition and Behavior, Radboud University Medical Center, Nijmegen, The Netherlands Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands Center for Brain and Cognition, University Pompeu Fabra, Barcelona, Spain Centre for Language Studies, Radboud University, Nijmegen, The Netherlands

807	10	School of Psychology and Artificial Intelligence, Radboud University, Nijmegen, the
808		Netherlands
809	11	Interdisciplinary Hub for Security, Privacy and Data Governance, Radboud University,
810		Nijmegen, the Netherlands
811	12	Institute for Molecules and Materials, Radboud University, Nijmegen, the Netherlands
812	13	University Medical Center Groningen, Groningen, The Netherlands