

The search for continuous gravitational waves from small-ellipticity sources at low frequencies

Vladimir Dergachev^{1,2, a} and Maria Alessandra Papa^{1,2,3, b}

¹Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Callinstraße 38, 30167 Hannover, Germany

²Leibniz Universität Hannover, D-30167 Hannover, Germany

³University of Wisconsin Milwaukee, 3135 N Maryland Ave, Milwaukee, WI 53211, USA

We present the results of an all-sky search for continuous gravitational wave signals with frequencies in the 20-500 Hz range from neutron stars with ellipticity of $\approx 10^{-8}$. This frequency region is particularly hard to probe because of the quadratic dependence of signal strength on frequency. The search employs the Falcon analysis pipeline [9] on LIGO O2 public data. Compared to previous Falcon analyses the coherence length has been quadrupled, with a corresponding increase in sensitivity. This enables us to search for small ellipticity neutron stars in this low frequency region up to 44 pc away. The frequency derivative range is up to 3×10^{-13} Hz/s easily accommodating sources with ellipticities of 10^{-7} and a corresponding factor of 10 increase in reach. New outliers are found, many of which we are unable to associate with any instrumental cause.

I. INTRODUCTION

Detectable continuous gravitational waves are expected from fast rotating neutron stars if they present some sort of non-axially symmetric deformation. The deformation in this context is usefully described by the *ellipticity* of the object, defined as $I_{zz}/(I_{xx} - I_{yy})$, where I is the moment of inertia tensor of the star and \hat{z} is along the star's rotation axis [1].

In our previous papers [2, 3] we searched for gravitational wave emission in the 500-2000 Hz range, targeting objects with ellipticity of 10^{-8} . We found a number of outliers, some corresponding to known instrumental artifacts, some in data with pristine frequency spectrum.

Ellipticities of 10^{-8} are interesting: they are well within the range of what the neutron star crusts can sustain [4] and observational indications exist that millisecond pulsars might have a *minimum* ellipticity of $\approx 10^{-9}$ [5].

There have been two recent papers [6, 7] with models of populations of potentially detectable neutron stars.

In [6] the authors conclude that the most likely detection will happen in the frequency range covered by this search. Their conclusions are somewhat pessimistic due in part to a number of simplifying assumptions. In particular they assume an exponential model of frequency evolution, rather than a power law that one expects when spin down is dominated by gravitational or electromagnetic emission. We want to highlight that it is reasonable to make simplifications when starting work on such a complicated question, and with further study the predictions will get refined.

In [7] the authors favor higher frequencies and higher spindown rates than we have searched. This is due to the focus of paper [7] on “probing” neutron stars at a given

ellipticity, without taking into account the frequency evolution of neutron stars. This approach is also reasonable, because finding a high-ellipticity source would be very interesting and thus it is important to know which parameter space to search.

However, finding a high-ellipticity and high frequency source is difficult. A source with a given ellipticity will lose energy due to the emission of gravitational waves and electromagnetic braking. The gravitational wave luminosity grows with the square of the rotation frequency. An isolated high frequency (~ 1000 Hz) and thus high ellipticity source ($\sim 10^{-6}$) would have a fairly rapid frequency drift ($\sim -10^{-8}$ Hz/s) just due to emission of gravitational waves alone. At such spindown rate its frequency will decrease by 100 Hz in the span of only 300 years. Thus we would need to be very lucky to be observing a high-ellipticity isolated neutron star at the very beginning of its spin down evolution.

Sources with smaller ellipticities and smaller spin rates, on the other hand, evolve more slowly. In this paper we explore these sources: with ellipticities of 10^{-8} and 10^{-7} and spin frequencies below 250 Hz, corresponding to the $\ell = m = 2$ gravitational wave mode emission at twice the rotation frequency. In addition, the distribution of known radio pulsars has many more sources in this range.

The spindown of the population of known pulsars has a considerably wider spread in our target frequency range, compared to our previous low-ellipticity searches [2, 3]. While there are known pulsars with frequency drift on the order of 10^{-10} Hz/s, most are well within 3×10^{-13} Hz/s which is the frequency drift tolerance used by this search.

Similarly to [8], we make a tradeoff between breadth and depth of search. Compared to [14, 15], this search concentrates on low ellipticity/low spindown objects and achieves a factor of 2 and 30% better sensitivity (Fig. 1), respectively, than [14] and [15]. Compared to [8], that explored 1 Hz, our frequency range covers the entire 20–500 Hz space. The improvements to the Falcon pipeline in computing speed allow 2 day coherence length for the first analysis stage, a factor of 4 longer than used in our

^a vladimir.dergachev@aei.mpg.de

^b maria.alessandra.papa@aei.mpg.de

previous searches [2, 3].




FIG. 1. Gravitational wave intrinsic amplitude h_0 upper limits at 95% confidence as a function of signal frequency. The upper limits are a measure of the sensitivity of the search. We compare with the latest LIGO/Virgo all-sky results in this frequency range [14], which are a factor $\gtrsim 2$ less constraining than ours, albeit able to detect sources with much larger distortions.

II. THE SEARCH AND THE PARAMETER SPACE

We use LIGO O2 open data [17, 18]. Our search looks for signals that are phase coherent over a certain timespan, called the coherence-length, and that evolve slowly over longer timescales. The search detects with highest efficiency IT2 signals. In [3] we have introduced the notion of “IT2 model” continuous wave signals because in the literature IT2 signals are used as a reference to measure search performance, and for upper limits. These are signals with constant amplitude and a frequency evolution that can be described by a Taylor expansion around nominal frequency and frequency derivative values, see for example Section II of [1]. The “I” indicates the source being isolated and “T2” indicates an intrinsic frequency evolution given by a Taylor polynomial of degree 2.

Our target frequency range of 20-500 Hz is particularly difficult to analyze because of a multitude of detector artifacts, including combs below 100 Hz. Furthermore, for the same physical distortion of the neutron star, the putative signals are weaker at these frequencies, compared to higher frequencies.

In order to overcome this, we increase the sensitivity of the Falcon pipeline by using long coherence lengths




FIG. 2. Reach of the search for stars with ellipticity of 10^{-8} . The search is also sensitive to sources with ellipticities of 10^{-7} with a distance from Earth that is 10 times higher. The X axis is the gravitational wave frequency, which is twice the pulsar rotation frequency for emission due to an equatorial ellipticity. R-modes and other emission mechanisms give rise to emission at different frequencies. The top curve (purple) shows the reach for a population of circularly polarized sources; The middle curve (cyan) holds for a population of sources with random orientations; The bottom curve (blue) holds for linearly polarized sources.

at all stages of the search, as shown in Table I. Regions of parameter space associated with high SNR results are searched again with progressively increasing coherence length.

With these choices we would be able to see sources with ellipticity of 10^{-8} up to 44 pc away (Figure 2). The frequency derivative search range ($\pm 3 \times 10^{-13}$ Hz/s) has been chosen to accommodate sources with ellipticities of 10^{-7} . These sources could be seen up to a distance of 440 pc.

Stage	Coherence length (days)	Minimum SNR
1	2	6
2	3	9
3	4	11
4	6	12
5	9	16
6	16	16
7	16	16

TABLE I. Parameters for each stage of the search. Stage 7 refines outlier parameters, and then subjects them to an additional consistency check.

FIG. 3. One of anomalies in our upper limits due to errors in public data. At these high frequencies the shot noise is dominant. A 60 Hz power line harmonic can add to it, but a dip below the shot noise is completely unphysical. We suspect that the origin of this type of feature is the cleaning procedure applied to the publicly released data. We would very much like the opportunity to understand and fully characterize this effect, as we explain in the text.

III. ANOMALIES IN O2 DATA

The O2 LIGO data have been cleaned. The procedure removes a number of instrumental artifacts by regressing against instrumental channels. The original implementation [19] was fairly conservative and was designed for searches of transient signals.

The procedure was then applied more aggressively using a significantly higher number of instrumental channels. In this type of procedure overfitting is a danger. One of the signs of overfitting is the appearance of artificially low values in the cleaned data. We see examples of this in [14] in the upper limit and in the power spectral density plots. And we find such systematic anomalies also in the upper limits from this search, Figure 3.

At high frequencies the main contribution to the noise floor is shot noise, and regressing against auxiliary channels recording measurements from, say, magnetometers and microphones, cannot reduce the noise below this level. In these specific regions we hence know that the calibration is too low, but we do not know by how much. Even though our pipeline establishes robust upper limits, it cannot correct for input data with incorrect amplitude values.

A systematic underestimation of the noise level is also problematic when considering the search algorithms. Gravitational wave searches usually combine separate

measurements with weights that down-weight noisier periods. Data with artificially low noise skew the weights and give more importance to unphysical data. If this happens in connected narrow bands, for broad-band surveys of continuous waves the overall impact should be marginal. The consistent recovery of the hardware injections (see for example [15]) shows that the issue probably does not significantly impact all frequencies. The effect shown in Figure 3 is small and, if that is all that happens, we could simply excise affected bands.

We are concerned that whatever is causing the low-noise anomalies, has not been fully characterized and understood and hence could be affecting the data in more subtle ways, without leaving such a visible signature, but still impacting the search results.

If the auxiliary channels used to produce the cleaned data were publicly available, we would try and characterize features and magnitude of this effect, and derive corrections for our upper limits. In the absence of this, we present our upper limits with the caveat that they rely on the data with imperfect calibration.

Our expectation is that in the majority of bands the amplitude is artificially low by at most a few percent and varies smoothly. In these bands we do not expect a large effect on phase and thus on our ability to detect signals. It would be very interesting to perform a hardware injection at a frequency where we observe the cleaning procedure to produce low strain data, and check the recovery of the signal parameters in that situation.

IV. RESULTS

The search produces upper limits that bound the strength of possible signals [20]. The upper limits also hold if a signal is present at a detectable level. In fact, in the band hosting loud outliers, for example, due to the hardware injections, the upper limit values are higher than elsewhere. In order to present the results in a concise way, the reported upper limits are produced by maximization over large subsets of parameter space. Internally, however, we compute a much more detailed picture. We show the "worst-case" upper limits - where the maximum is taken over all polarizations, and the "best-case" upper limits where the maximum is taken among circularly polarized upper limits only. The latter correspond to sources for which the search is the most sensitive. In order to compare with population-average upper limits common in the literature, we also estimate those.

The gravitational wave amplitude upper limits from this search are shown in Figure 1 and are available in numerical form in [21].

Our upper limits can be translated into limits on gravitational waves from boson condensates around black holes [22, 23], which are expected to emit monochromatic continuous wave signals [24] with very small frequency drift. We leave it to the interested reader to constrain from our upper limits physical quantities of in-

SNR	f Hz	\dot{f} pHz/s	RA _{J2000} degrees	DEC _{J2000} degrees	Comment
7417.8	265.57535	-1.0	71.591	-56.223	Injection ip0
6383.7	31.42473	-0.5	285.078	-58.269	Injection ip11
3712.5	52.80832	-0.2	302.538	-83.843	Injection ip5
1895.0	26.33722	-0.9	234.716	41.244	Injection ip10
1497.4	108.85716	0.0	178.369	-33.439	Injection ip3
27.1	83.43998	0.3	233.838	-41.127	
27.0	59.99418	0.6	211.421	-21.567	60 Hz line
25.2	60.02252	-0.2	25.723	37.383	60 Hz line
25.1	80.40734	-0.9	225.641	77.846	Line in H1
21.9	465.58924	0.6	87.318	50.980	
21.9	197.87552	-0.8	276.984	-62.431	
21.6	400.01133	0.0	66.157	89.153	Coincident lines
21.6	31.84219	0.6	272.835	67.366	
21.5	449.50129	0.7	308.074	-32.371	
21.3	55.60198	0.7	218.116	-68.806	
21.2	160.49660	0.5	90.941	32.384	Line in L1
21.0	140.26640	0.7	179.620	-19.718	Line in H1
21.0	357.19349	-0.6	202.263	-27.984	
21.0	153.87728	-0.2	29.374	-53.913	Line in H1
21.0	440.03835	0.5	301.795	23.622	
20.8	31.70025	-1.1	101.926	-66.112	Line in H1
20.8	260.10055	-0.5	85.537	17.565	
20.7	180.10461	-0.8	122.697	-77.297	60 Hz harmonic
20.6	307.28973	1.1	330.582	28.601	Near L1 line
20.6	441.11365	-0.1	343.747	-58.935	
20.6	442.38421	-0.0	20.788	-50.078	
20.5	312.31246	0.6	284.115	-42.329	
20.5	427.73851	0.8	259.995	-44.007	
20.5	456.66685	0.8	161.618	67.993	Broad line in H1
20.4	457.11691	-0.0	147.143	44.056	Broad line in H1
20.4	245.06138	1.0	49.632	12.941	
20.3	66.89584	0.8	177.962	-21.334	
20.2	448.02463	0.8	263.539	46.220	
20.2	153.97881	-0.5	131.683	-58.036	Line in H1
20.2	459.31715	0.0	341.763	20.999	
20.2	192.60299	-0.6	340.182	50.884	
20.1	394.38609	-0.1	257.670	-82.979	
20.1	478.98702	0.3	151.593	-67.970	Line in L1
20.1	317.99684	-1.0	26.269	-22.945	
20.0	436.12088	-1.3	342.703	-40.723	

TABLE II. High SNR outliers produced by the detection pipeline. We list outliers with SNR above 20, and exclude those near ecliptic poles and associated with stationary lines. Only the highest-SNR outlier is shown for each 0.1 Hz frequency region. Outliers marked “ipX” are due to a simulated signals “hardware-injected” during the science run for validation purposes. Their parameters are listed in Table III. Outliers marked with “line” have strong narrowband disturbances near the outlier frequency. Signal frequencies refer to GPS epoch 1176425033. Full list of outliers down to SNR > 16 is available in [21].

terest, based on the specific model they wish to consider (for example, an ensemble signal of [25]).

The search also identifies small areas of parameter space containing waveforms that are unusually consistent with the data. These are the outliers of the search. The outliers are often caused by signals of terrestrial origin due to technical or environmental noise sources; the weaker ones may also be fluctuations. A key result of our search is a number of interesting outliers (Table II), many of which are located in bands with clean frequency spectrum. Table II only lists outliers with SNR above 20

and excludes outliers near ecliptic pole most likely associated with instrumental lines, in particular the outliers from the 0.5 Hz frequency comb. The full list of outliers down to SNR 16 is available in [21].

Monte Carlo simulations show that we detect IT2 signals within 8×10^{-6} Hz of the signal frequency f , within 10^{-12} Hz/s of its frequency derivative, and within $0.12 \text{ Hz}/f$ radians from its sky location, the latter calculated after projection on the ecliptic plane (“ecliptic distance”). For non-IT2 signals, or signals with frequency derivative outside search range, the tolerances should be

Label	Frequency Hz	Spindown pHz/s	RA _{J2000} degrees	DEC _{J2000} degrees
ip0	265.575343	-4.15	71.55193	-56.21749
ip3	108.857159	-1.46×10^{-5}	178.37257	-33.4366
ip5	52.808324	-4.03×10^{-6}	302.62664	-83.83914
ip6	145.860492	-6730	358.75095	-65.42262
ip8	190.634274	-8650	351.38958	-33.41852
ip10	26.338016	-85	221.55565	42.87730
ip11	31.424735	-0.507	285.09733	-58.27209
ip12	38.764787	-6250	331.85267	-16.97288

TABLE III. Parameters of the hardware-injected simulated continuous wave signals during the O2 data run (GPS epoch 1176425033).

widened. In particular, there is a degeneracy between sky location and frequency derivative, so a mismatch in the latter results in outlier offset from its true position on the sky.

Because the coherence length of our followup stage has been extended to 16 days, we have much tighter tolerance for the signals that pass our pipeline compared with previous searches. While we see 5 simulated signals clearly (ip0, ip3, ip5, ip10 and ip11, Table III), the injections ip6 and ip12 did not produce any outliers as their frequency derivative smaller than -5×10^{-9} Hz/s is well beyond our target parameter space.

V. CONCLUSIONS

We have performed the most sensitive search for continuous gravitational waves in the frequency band of 20-500 Hz. Just as in our previous papers [2, 3], we have a number of interesting outliers. Our constraints are tighter than before, the signal-to-noise ratio is higher. Are they due to some sort of noise? Or are we seeing astrophysical sources? The latter is unlikely if we assume a classic IT2 source with ellipticity of 10^{-8} - we simply do not expect that many sources within 44 pc. Even widening the reach to 440 pc by including ellipticities of 10^{-7} , it would be surprising that this many outliers would be of astrophysical origin.

We look forward to more data for further analysis.

ACKNOWLEDGMENTS

The authors thank the scientists, engineers and technicians of LIGO, whose hard work and dedication produced the data that made this search possible.

The search was performed on the ATLAS cluster at AEI Hannover. We thank Bruce Allen, Carsten Aulbert and Henning Fehrmann for their support.

This research has made use of data, software and/or web tools obtained from the LIGO Open Science Center (<https://losc.ligo.org>), a service of LIGO Labora-

tory, the LIGO Scientific Collaboration and the Virgo Collaboration. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by Polish and Hungarian institutes.

-
- [1] P. Jaranowski, A. Krolak and B. F. Schutz, “Data analysis of gravitational - wave signals from spinning neutron stars. 1. The Signal and its detection,” *Phys. Rev. D* **58** (1998), 063001
- [2] V. Dergachev, M. A. Papa, “Results from the first all-sky search for continuous gravitational waves from small-ellipticity sources”, *Phys. Rev. Lett.* **125**, no.17, 171101 (2020)
- [3] V. Dergachev, M. A. Papa, “Results from high-frequency all-sky search for continuous gravitational waves from small-ellipticity sources”, *Phys. Rev. D* **103**, 063019 (2021)
- [4] F. Gittins, N. Andersson and D. I. Jones, “Modelling neutron star mountains,” [arXiv:2009.12794 [astro-ph.HE]].
- [5] G. Woan, M. D. Pitkin, B. Haskell, D. I. Jones, and P. D. Lasky, Evidence for a Minimum Ellipticity in Millisecond Pulsars, *ApJL* **863** L40 (2018)
- [6] Detectability of continuous gravitational waves from isolated neutron stars in the Milky Way: the population synthesis approach, Marek Cieřlar, Tomasz Bulik, Małgorzata Curyło, Magdalena Sieniawska, Neha Singh, Michał Bejger, arXiv:2102.08854
- [7] Modeling the Galactic Neutron Star Population for Use in Continuous Gravitational Wave Searches Brendan T. Reed, Alex Deibel, C. J. Horowitz, arXiv:2104.00771
- [8] Karl Wette, Liam Dunn, Patrick Clearwater, Andrew Melatos, Deep exploration for continuous gravitational waves at 171–172 Hz in LIGO second observing run data, arXiv:2103.12976
- [9] V. Dergachev and M. A. Papa, “Sensitivity improvements in the search for periodic gravitational waves using O1 LIGO data,” *Phys. Rev. Lett.* **123**, no.10, 101101 (2019) doi:10.1103/PhysRevLett.123.101101 [arXiv:1902.05530 [gr-qc]].
- [10] V. Dergachev, On blind searches for noise dominated signals: a loosely coherent approach, *Class. Quantum Grav.* **27**, 205017 (2010).
- [11] V. Dergachev, Loosely coherent searches for sets of well-modeled signals, *Phys. Rev. D* **85**, 062003 (2012)
- [12] V. Dergachev, Loosely coherent searches for medium scale coherence lengths, arXiv:1807.02351
- [13] LIGO Data Management Plan textttt <https://dcc.ligo.org/public/0009/M1000066/025/LIGO-M1000066-v25.pdf> (2017)
- [14] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), All-sky search for continuous gravitational waves from isolated neutron stars using Advanced LIGO O2 data, *Phys. Rev. D* **100** 024004 (2019).
- [15] B. Steltner, M. A. Papa, H.-B. Eggenstein, B. Allen, V. Dergachev, R. Prix, B. Machenschalk, S. Walsh, S. J. Zhu, S. Kwang, Einstein@Home all-sky search for continuous gravitational waves in LIGO O2 public data, arXiv:2009.12260
- [16] P. Nguyen, R. M. S. Schofield, A. Effler, C. Austin, V. Adya, *et al.*, Environmental Noise in Advanced LIGO Detectors, arXiv:2101.09935 [astro-ph.IM]
- [17] M. Vallisneri *et al.* The LIGO Open Science Center, proceedings of the 10th LISA Symposium, University of Florida, Gainesville, 2015 *J. Phys.: Conf. Ser.* 610 012021
- [18] LIGO Open Science Center, <https://doi.org/10.7935/CA75-FM95>, (2019)
- [19] J.C. Driggers *et al.* (LIGO Scientific Collaboration Instrument Science Authors), Improving astrophysical parameter estimation via offline noise subtraction for Advanced LIGO, *Phys. Rev. D* **99**, 042001 (2019).
- [20] V. Dergachev, A Novel Universal Statistic for Computing Upper Limits in Ill-behaved Background, *Phys. Rev. D* **87**, 062001 (2013).
- [21] See EPAPS Document No. [number will be inserted by publisher] for numerical values of upper limits, outlier tables and hardware injection parameters. Also at <https://www.aei.mpg.de/continuouswaves/02Falcon20-500>
- [22] M. Baryakhtar, R. Lasenby, M. Teo, Black Hole Super-radiance Signatures of Ultralight Vectors, *Phys. Rev. D* **96**, 035006s (2017)
- [23] A. Arvanitaki, M. Baryakhtar, R. Lasenby, S. Dimopoulos, S. Dubovsky, Black Hole Mergers and the QCD Axion at Advanced LIGO, *Phys. Rev. D* **95**, 043001 (2017)
- [24] A. Arvanitaki, M. Baryakhtar, and X. Huang, Discovering the QCD axion with black holes and gravitational waves, *Phys. Rev. D* **91**, 084011 (2015)
- [25] S. J. Zhu, M. Baryakhtar, M. A. Papa, D. Tsuna, N. Kawanaka and H. Eggenstein, Characterizing the continuous gravitational-wave signal from boson clouds around Galactic isolated black holes, *Phys. Rev. D* **102**, 063020 (2020)