


# Search for Gravitational Waves from the Coalescence of Substellar-Mass Binaries in the First Half of Advanced LIGO and Virgo's Third Observing Run

Alexander H. Nitz<sup>1</sup> and Yi-Fan Wang (王一帆)<sup>2</sup>

*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany  
and Leibniz Universität Hannover, D-30167 Hannover, Germany*

 (Received 18 June 2021; revised 17 July 2021; accepted 31 August 2021; published 6 October 2021)

We present a search for gravitational waves from the coalescence of substellar-mass black hole binaries using data from the first half of Advanced LIGO and Virgo's third observing run. The observation of a substellar-mass black hole merger may be an indication of primordial origin; primordial black holes may contribute to the dark matter distribution. We search for black hole mergers where the primary mass is  $0.1\text{--}7 M_{\odot}$  and the secondary mass is  $0.1\text{--}1 M_{\odot}$ . A variety of models predict the production and coalescence of binaries containing primordial black holes; some involve dynamical assembly, which may allow for residual eccentricity to be observed. For component masses  $> 0.5 M_{\odot}$ , we also search for sources in eccentric orbits, measured at a reference gravitational-wave frequency of 10 Hz, up to  $e_{10} \sim 0.3$ . We find no convincing candidates and place new upper limits on the rate of primordial black hole mergers. The merger rate of  $0.5\text{--}0.5$  ( $1.0\text{--}1.0$ )  $M_{\odot}$  sources is  $< 7100(1200) \text{ Gpc}^{-3} \text{ yr}^{-1}$ . Our limits are  $\sim 3\text{--}4$  times more constraining than prior analyses. Finally, we demonstrate how our limits can be used to constrain arbitrary models of the primordial black hole mass distribution and merger rate.

DOI: [10.1103/PhysRevLett.127.151101](https://doi.org/10.1103/PhysRevLett.127.151101)

*Introduction.*—Gravitational-wave astronomy has entered an era of routine observations. The Advanced LIGO [1] and Virgo [2] observatories have now completed three observing runs (O1, O2, and O3); each was accompanied by significant increases in sensitivity [3]. To date, over 50 binary black hole mergers have been reported [4–6]. These observations have had significant impact on the study of the merger rate and population of compact objects [7]; notable events confirm the likely existence of black holes with component masses in the pair-instability gap ( $> 50 M_{\odot}$ ) [4,8] or in the region  $3\text{--}5 M_{\odot}$  [9]. The possibility that these extremal parts of the distribution [10–13] or a fraction of the bulk of observed mergers [14–22] may be due to the coalescence of primordial black holes (PBHs) is under active investigation.

Currently, there is no clear observational evidence for the existence of PBHs. However, in addition to providing an explanation for some of the observed LIGO and Virgo mergers [11], primordial black holes may be the origin of some observed microlensing incidents [23], excess cross-correlation between cosmic x-ray and cosmic microwave background [24], the current excess in gravitational-wave

background observed by NANOGrav [25], and the seeds for galaxy and supermassive black hole formation [26–28]. Many of these observations are also consistent with more mundane explanations and standard stellar formation scenarios [7]. In contrast, there are no known mechanisms through standard stellar evolution to produce substellar-mass black holes; the observation of a single substellar-mass black hole would be decisive for the existence of primordial black holes or for even more exotic scenarios such as dark matter triggered formation of black holes [29–31].

Several searches for gravitational waves from the coalescence of substellar-mass mergers have already been conducted using data from LIGO's first two observing runs (O1, O2); these include searches for comparable mass binary black holes [32–34], eccentric mergers [35], and high-mass-ratio sources [36]. No likely candidates have been found. In this Letter, we report a search for gravitational waves from the coalescence of black holes with primary mass  $0.1\text{--}7 M_{\odot}$  and secondary mass  $0.1\text{--}1 M_{\odot}$  using the open data from the first half of the third observing (O3a) run of Advanced LIGO and Virgo. The parameter space for our search region is shown in Fig. 1. The most significant candidate in our search has a false alarm rate of 1 per O(month). Given the time observed, we consider our results consistent with a null observation and place new limits on the rate of substellar-mass mergers that are  $3\text{--}4$  times more stringent than prior analyses due to the significant improvement of sensitivity and detector robustness of O3 compared to O1 and O2 [4].

---

*Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Open access publication funded by the Max Planck Society.*




FIG. 1. The regions searched by recent gravitational-wave analyses of the LIGO and Virgo data as a function of detector-frame primary and secondary mass. The region we search for noneccentric sources (green) and the region sensitive to sources with eccentricity up to  $e_{10} \sim 0.3$  (red) are shown. For comparison, we include the search region of the most recent subsolar-mass search by the LIGO-Virgo Collaboration (LVC) [33] (dashed), the search for high-mass-ratio mergers [36], and searches for standard stellar-mass sources [4,5], labeled by 3-OGC (3-Open Gravitational-wave Catalog) or GWTC-2 (Gravitational-Wave Transient Catalog-2). The axes are truncated at  $200 M_{\odot}$ .

Our limits on the rate of subsolar-mass mergers can be related to constraints on the fraction of dark matter composed of PBHs; this requires a model of the binary’s formation to predict the PBH abundance from the observed merger rate. Current astrophysical models have large uncertainties in their predictions of both the black hole mass function and binary formation rate [18–20,37–40]. Primordial black holes may form binaries in the early Universe if they can decouple from the cosmic expansion. However, it is under investigation what fraction of binaries would be disrupted in the following evolution. Reference [41] shows with  $N$ -body simulations that a significant fraction would be disrupted if the fraction that primordial black holes contribute to the dark matter is  $f_{\text{PBH}} = 100\%$ . Primordial black holes can also form binaries in the late Universe by dynamical capture due to gravitational-wave dissipation [18,42,43]; this scenario may also lead to residual eccentricity by the time it is observable by gravitational-wave detectors [44]. However, the event rate of binaries formed in the early Universe is expected to be dominant compared with the late Universe channel, depending on the intensity of binary disruption [41].

Because of the wide variety of models, in this Letter, we consider a fiducial model that assumes a monochromatic distribution of primordial black hole mass. The same model was used in past subsolar-mass searches [32–36], and we include it for comparison purposes. We also consider the uncertainty on the rate estimates arising from the fraction of binaries that are disrupted after formation [41]. Constraints

for specific models with broad mass distributions can be derived from our observational constraints.

*Search.*—We conduct our analysis in a similar manner to our previously presented search of the first two observing runs [35]; however, for the first time we include data from the Virgo observatory. We use the open-source PyCBC toolkit [45,46] to conduct a matched-filtering-based search [47]. Matched filtering allows us to extract a potential signal using the predicted gravitational waveform as a template. Potential candidates are assessed for consistency between the operating observatories [48] and against the expected morphology of the gravitational waveform [49,50]. Each candidate is assigned a ranking statistic value that takes into account these factors in addition to the measured noise variance [51,52].

The statistical significance of each candidate is assessed by comparing it to an empirically measured distribution of false alarms [45]. The distribution is measured by conducting numerous fictitious analyses, whereby we offset detectors’ data in time. This procedure purposefully violates the time-of-flight constraints between the detectors to remove coincident astrophysical sources and create analyses containing only false alarms [53–55].

As matched filtering requires accurate models of the expected gravitational-wave signal, we use a combination of the TaylorF2 post-Newtonian approximant accurate to 3.5PN [56–59] and the TaylorF2e model [60–62]. TaylorF2e is an extension of TaylorF2, which includes corrections for moderate eccentricity. Both TaylorF2 and TaylorF2e model only the inspiral portion of a gravitational-wave signal and do not account for the phase where the binary finally merges. The merger can be safely neglected as we search for sources only up to a total mass of  $8 M_{\odot}$ . For these sources, the merger occurs at a frequency above the most sensitive band of the instruments.

To search for a broad region, we use the stochastic algorithm [63] to create a discrete bank of templates designed to ensure that we recover  $> 95\%$  of a signal’s signal-to-noise ratio if it has parameters within the boundaries of our search. Our bank is designed to recover binaries in quasicircular orbits where the primary mass is  $0.1\text{--}7 M_{\odot}$  and the secondary mass is  $0.1\text{--}1 M_{\odot}$ . In addition, for sources with component masses  $> 0.5 M_{\odot}$ , the bank is designed to recover sources with eccentric orbits up to  $e_{10} \sim 0.3$ , where  $e_{10}$  is the eccentricity at the fiducial dominant-mode gravitational-wave frequency of 10 Hz. We assume that PBHs will have negligible spin; this is consistent with the predictions of PBH spin distributions [64–68]. To save on computational cost, we limit the starting frequency of each template so that its duration is  $< 512$  s; otherwise a cutoff at a gravitational-wave frequency of 20 Hz is used. The template bank was also constructed with this lower frequency criteria. These choices result in a bank with  $\sim 7.8 \times 10^6$  templates, where

50% of the templates have nonzero eccentricity and use the TaylorF2e model.

*Observational results.*—We search for gravitational waves from the coalescence of subsolar-mass compact binaries using the public LIGO and Virgo data from the first half of the third observing run (O3a) [69,70]; data from the second half of the observing run is not yet available. We analyze the nearly 150 days of data, where at least two observatories were operating; the twin LIGO observatories were operating for  $\sim 100$  days of this period. In comparison to the previous observing run, the LIGO instruments had  $\sim 30\%$ – $50\%$  greater range [3].

The most significant candidates from our search are identified at a false alarm rate of one per O(month) and shown in Table I; this is consistent with a null observation and our expectation that the noise candidates follow a Poisson distribution. Under the assumption of a null detection, we place limits on the rate of binary mergers at 90% confidence using the loudest event method of Ref. [71]. The limit on the merger rate  $R_{90}$  is given as

$$R_{90} = \frac{2.3}{VT}, \quad (1)$$

where  $V$  is the estimated sensitive volume of the analysis assessed at the false alarm rate of the most significant observed candidate and  $T$  is the duration of the observation period. We estimate the surveyed volume-time of our analysis by measuring our analysis' response to a simulated population of  $O(10^5)$  sources. We assume that sources are isotropically distributed in their orientations and sky location in addition to a uniform distribution in volume. We measure the mass dependence of our sensitive volume using separate simulation sets each with fixed-source masses. For simulations that include eccentricity, we assume a uniform distribution where  $e_{10} \in [0, 0.3)$ .

The resulting upper limit on the merger rate, combined with the limit from the prior analysis of O1 and O2 [35], is shown as a function of chirp mass in Fig. 2, where chirp mass is defined as  $\mathcal{M} = (m_1 m_2)^{3/5} / (m_1 + m_2)^{1/5}$  and  $m_{1,2}$  are the masses of the two components of a binary. For sources within our target parameter space, this limit also holds for sources with varied mass ratios, but the same

TABLE I. The top five candidates in our search with the highest inverse false alarm rates (IFARs). The Global Positioning System (GPS) time of each candidate, the component mass  $m_{1/2}$ , and the eccentricity of the template chosen by the search are shown.

GPS time	IFAR (yr)	$m_1/M_\odot$	$m_2/M_\odot$	$e_{10}$
1245411568.354	0.084	0.69	0.21	0.00
1242817372.434	0.079	0.86	0.11	0.00
1246418221.718	0.075	0.13	0.13	0.00
1252963276.322	0.062	1.05	0.52	0.28
1240000657.632	0.057	3.04	0.10	0.00

chirp mass; a similar conclusion was noted in Ref. [33]. Our results limit the merger rate for  $0.1$ – $0.1$ ,  $0.5$ – $0.5$ , and  $1.0$ – $1.0 M_\odot$  sources to  $< 670\,000$ ,  $7100$ , and  $1200 \text{ Gpc}^{-3} \text{ yr}^{-1}$ , respectively. This is an improvement between 3 and 4 times over previous analyses, which only used data from the first two observing runs.

*Implications for primordial black hole abundance.*—We use the observational upper limits on the subsolar-mass compact binary merger rate to constrain models of primordial black hole binary formation. Existing models have significant uncertainties on the primordial black hole mass distribution and the binary formation rate [10,18–20,37–40]. Constraints on models can be derived from the observational event rate upper limits given a specific primordial black hole mass distribution. We consider a fiducial mass distribution, a delta distribution following Refs. [32,33], and our previous work [35,36] to allow for a consistent comparison.

For the binary formation rate, we consider the mechanism initially proposed in [37] and developed in [20,22,39–41,72], where primordial black holes form bounded binaries in the early Universe and merge recently. The binaries formed in the late Universe is subdominant [44]; therefore we use the search results from circular binaries to constrain the early Universe formation model. The binary merger rate is given by Refs. [39,40] for a general mass distribution  $P(m)$




FIG. 2. Upper limit on the rate of mergers at 90% confidence ( $R_{90}$ ) for our search (purple). For comparison, we show the previous limits from our search of the O1 and O2 data (blue dashed) [35] in addition to the most recent results from subsolar-mass searches conducted by the LVC of the O2 data (black dotted) [33]. These all assume that sources are quasicircular; constraints assuming a uniform distribution of  $e_{10} \in [0, 0.3)$  are also shown (green). As expected, the limit for eccentric sources closely matches that for quasicircular binaries where the component masses are  $> 0.5 M_\odot$ . Shaded regions show the one sigma uncertainty on the rate due to the Monte Carlo estimation of the search's surveyed volume-time.

$$\begin{aligned}
 R(\tilde{f}_{\text{PBH}}, m_1, m_2) &= 3 \times 10^6 \tilde{f}_{\text{PBH}}^2 (0.7 \tilde{f}_{\text{PBH}}^2 + \sigma_{\text{eq}}^2)^{-\frac{21}{4}} (m_1 m_2)^{\frac{3}{37}} \\
 &\times \left( m_1 + m_2 \right)^{\frac{36}{37}} \min \left( \frac{P(m_1)}{m_1}, \frac{P(m_2)}{m_2} \right) \\
 &\times \left( \frac{P(m_1)}{m_1} + \frac{P(m_2)}{m_2} \right) \text{Gpc}^{-3} \text{yr}^{-1}, \quad (2)
 \end{aligned}$$

where  $R dm_1 dm_2$  is the event rate at the binary component masses  $m_{1/2} M_\odot$ . The normalization for mass distribution is  $\int P(m) dm = 1$ . The parameter  $\sigma_{\text{eq}} = 0.005$  is the variance of dark matter density perturbation at the matter radiation equality epoch [40], and we keep this factor separate from other binary disruption effects to perform consistent comparison with previous work. Reference [41] has used an  $N$ -body simulation to show most binaries after formation would be disrupted by their environment if  $f_{\text{PBH}} = 100\%$  and thus introduced a suppression factor  $S$  with value  $< 1$ , which accounts for the disruption as a function of primordial black hole mass and fraction. For a relatively narrow mass distribution, Ref. [22] shows  $S$  (referred to as  $S_2$  in Ref. [22]) is estimated to be 1% for  $f_{\text{PBH}} = 100\%$  and  $\sim 1$  for  $f_{\text{PBH}} < 1\%$  and the suppression is only a function of  $f_{\text{PBH}}$ . However, underestimation by one order of magnitude may exist for  $f_{\text{PBH}} = 100\%$  if the merger of binaries perturbed by the environment after formation is included [22,34,72]. To account for this uncertainty, we define an effective fraction that relates to the true fraction by  $\tilde{f}_{\text{PBH}}^{53/37} = S f_{\text{PBH}}^{53/37}$ . Constraints in our past work [35,36] on  $f_{\text{PBH}}$  should also be understood as constraints on  $\tilde{f}_{\text{PBH}}$ ; they implicitly assume the suppression factor is unity. The true fraction  $f_{\text{PBH}}$  can be recovered with a well-understood estimation of  $S$ .

For a delta distribution of mass, Eq. (3) is reduced to

$$R(\tilde{f}_{\text{PBH}}, m) = 3 \times 10^6 \tilde{f}_{\text{PBH}}^2 (0.7 \tilde{f}_{\text{PBH}}^2 + \sigma_{\text{eq}}^2)^{-\frac{21}{4}} m^{-\frac{32}{37}}. \quad (3)$$

Using the observational rate limit given in Fig. 2, the upper limit on the effective fraction of primordial black hole with a delta distribution of mass is shown in Fig. 3. A comparison with previous search results for subsolar-mass compact binaries is also plotted. The results should be interpreted as constraints on the combined effect on the binary merger rate from the abundance of primordial black holes and the merger rate suppression due to environmental interaction. Future improvements on theoretical modeling and observational results can resolve the entanglement. Overall, results in Fig. 3 show that our constraints are  $\sim 2$ – $3$  times tighter than [35] using O1 and O2 data.

Constraints can also be derived on primordial black hole scenarios that predict broad mass distributions. By extending Eq. (1), the corresponding 90% upper limits can be obtained by requiring




FIG. 3. The upper limits on the effective fraction of the primordial black hole contribution to dark matter  $\tilde{f}_{\text{PBH}}$ , from the search for subsolar-mass compact binaries in LVC O1–O3a data. As a comparison, we also plot the constraints from O1–O2 data [35] for the monochromatic mass distribution and for the two-point mass distribution of high-mass-ratio (HMR) binaries [36], where the primary mass is fixed to the average of the 2-OGC events  $\approx 37 M_\odot$ . Note that this latter limit requires the additional assumption that the primary mass abundance be consistent with accounting for the majority of LIGO-observed binary black hole mergers.

$$\int R(\vec{\theta}, m_1, m_2) VT(m_1, m_2) dm_1 dm_2 = 2.3, \quad (4)$$

where  $R$  is the model-predicted merger rate density as a function of the component masses and may include additional model parameters  $\vec{\theta}$ .  $VT$  is the surveyed volume-time of our analysis as a function of the component masses; this is available as part of our data release.

To illustrate, we consider the scenario that primordial black holes are produced in the early Universe QCD phase transition era [10]. This model is shown to have a peak at  $\sim 1 M_\odot$  and to be able to account for the origin of the extremal parts of the already observed binary black hole merger distribution. We choose the mass distribution  $P(m)$  by following Ref. [10] from the early Universe scalar perturbation spectral index 0.96 and use the binary formation rate of Eq. (2). We assume a mass-independent suppression factor may be applicable here given the narrow peak of  $P(m)$ . The observational upper limits on event rate from our search require  $\tilde{f}_{\text{PBH}} \leq 1\%$  in this scenario. For more general mass distributions, Eq. (4) is capable of constraining the PBH abundance or other model parameters given a specific primordial black hole mass distribution and prescription that can predict the resulting merger rate density  $R(\vec{\theta}, m_1, m_2)$ .

*Conclusions.*—We conduct a search for gravitational waves from the coalescence of subsolar-mass black holes using data from the first half of the third observing run of Advanced LIGO and Virgo. We find no clear detections and



so place new limits on the rate of mergers. The increased sensitivity of the O3a data allows us to improve upon the state-of-the-art limits by 3–4 times. The second half of the O3 data (O3b) would be expected to improve these limits by another factor of 1.5–2 times.

We apply our rate limits to a fiducial monochromatic mass distribution and compare our results to the limits from prior analyses. Overall, the constraints on the effective fraction of primordial black holes in dark matter are 2–3 tighter than the previous results. Our observational results can also help resolve the event rate modeling uncertainties by constraining model parameters. If we assume that dark matter consisted entirely of black holes, we constrain the suppression factor to  $S \leq 0.1\%$  for  $0.5 - 0.5 M_{\odot}$  mergers.

Lastly, we demonstrate how to apply our limits to models that can predict the rate of PBH mergers.

To aid in comparing our rate limits to the various formation scenarios leading to subsolar-mass mergers, we make the detailed constraints available [73]. In addition, we make available the configuration files and template bank necessary to reproduce the analysis.

We acknowledge the Max Planck Gesellschaft. We thank the computing team from AEI Hannover for their significant technical support. This research has made use of data from the Gravitational Wave Open Science Center ([74]), a service of LIGO Laboratory, the LIGO Scientific Collaboration, and the Virgo Collaboration. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN), and the Dutch Nikhef, with contributions by Polish and Hungarian institutes.

---

\* alex.nitz@aei.mpg.de

- [1] J. Aasi *et al.* (LIGO Scientific Collaboration), Advanced LIGO, *Classical Quantum Gravity* **32**, 115012 (2015).
- [2] F. Acernese *et al.* (VIRGO Collaboration), Advanced Virgo: A second-generation interferometric gravitational wave detector, *Classical Quantum Gravity* **32**, 024001 (2015).
- [3] A. Buikema *et al.* (aLIGO Collaboration), Sensitivity and performance of the Advanced LIGO detectors in the third observing run, *Phys. Rev. D* **102**, 062003 (2020).
- [4] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run, *Phys. Rev. X* **11**, 021053 (2021).
- [5] A. H. Nitz, C. D. Capano, S. Kumar, Y.-F. Wang, S. Kasta, M. Schäfer, R. Dhurkunde, and M. Cabero, 3-OGC: Catalog of gravitational waves from compact-binary mergers, [arXiv:2105.09151](https://arxiv.org/abs/2105.09151).
- [6] T. Venumadhav, B. Zackay, J. Roulet, L. Dai, and M. Zaldarriaga, New binary black hole mergers in the second observing run of Advanced LIGO and Advanced Virgo, *Phys. Rev. D* **101**, 083030 (2020).
- [7] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), Population properties of compact objects from the second LIGO-Virgo gravitational-wave transient catalog, *Astrophys. J. Lett.* **913**, L7 (2021).
- [8] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), GW190521: A Binary Black Hole Merger with a Total Mass of  $150 M_{\odot}$ , *Phys. Rev. Lett.* **125**, 101102 (2020).
- [9] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), GW190814: Gravitational waves from the coalescence of a 23 solar mass black hole with a 2.6 solar mass compact object, *Astrophys. J. Lett.* **896**, L44 (2020).
- [10] B. Carr, S. Clesse, J. García-Bellido, and F. Kühnel, Cosmic conundra explained by thermal history and primordial black holes, *Phys. Dark Universe* **31**, 100755 (2021).
- [11] S. Clesse and J. Garcia-Bellido, GW190425, GW190521 and GW190814: Three candidate mergers of primordial black holes from the QCD epoch, [arXiv:2007.06481](https://arxiv.org/abs/2007.06481).
- [12] K. Vattis, I. S. Goldstein, and S. M. Koushiappas, Could the  $2.6 M_{\odot}$  object in GW190814 be a primordial black hole? *Phys. Rev. D* **102**, 061301(R) (2020).
- [13] V. De Luca, V. Desjacques, G. Franciolini, P. Pani, and A. Riotto, GW190521 Mass Gap Event and the Primordial Black Hole Scenario, *Phys. Rev. Lett.* **126**, 051101 (2021).
- [14] G. Franciolini, V. Baibhav, V. De Luca, K. K. Y. Ng, K. W. K. Wong, E. Berti, P. Pani, A. Riotto, and S. Vitale, Evidence for primordial black holes in LIGO/Virgo gravitational-wave data, [arXiv:2105.03349](https://arxiv.org/abs/2105.03349).
- [15] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, Bayesian evidence for both astrophysical and primordial black holes: Mapping the GWTC-2 catalog to third-generation detectors, *J. Cosmol. Astropart. Phys.* **05** (2021) 003.
- [16] K. Jedamzik, Consistency of Primordial Black Hole Dark Matter with LIGO/Virgo Merger Rates, *Phys. Rev. Lett.* **126**, 051302 (2021).
- [17] J. García-Bellido, J. F. Nuño Siles, and E. R. Morales, Bayesian analysis of the spin distribution of LIGO/Virgo black holes, *Phys. Dark Universe* **31**, 100791 (2021).
- [18] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz, A. Raccanelli, and A. G. Riess, Did LIGO Detect Dark Matter? *Phys. Rev. Lett.* **116**, 201301 (2016).
- [19] S. Clesse and J. García-Bellido, The clustering of massive primordial black holes as dark matter: Measuring their mass distribution with Advanced LIGO, *Phys. Dark Universe* **15**, 142 (2017).
- [20] M. Sasaki, T. Suyama, T. Tanaka, and S. Yokoyama, Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914, *Phys. Rev. Lett.* **117**, 061101 (2016).
- [21] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, Primordial black holes confront LIGO/Virgo data: Current situation, *J. Cosmol. Astropart. Phys.* **06** (2020) 044.
- [22] G. Hütsi, M. Raidal, V. Vaskonen, and H. Veermäe, Two populations of LIGO-Virgo black holes, *J. Cosmol. Astropart. Phys.* **03** (2021) 068.
- [23] H. Niikura, M. Takada, S. Yokoyama, T. Sumi, and S. Masaki, Constraints on Earth-mass primordial black holes from OGLE 5-year microlensing events, *Phys. Rev. D* **99**, 083503 (2019).

- [24] G. Hasinger, Illuminating the dark ages: Cosmic backgrounds from accretion onto primordial black hole dark matter, *J. Cosmol. Astropart. Phys.* **07** (2020) 022.
- [25] V. De Luca, G. Franciolini, and A. Riotto, Nanograv Data Hints at Primordial Black Holes as Dark Matter, *Phys. Rev. Lett.* **126**, 041303 (2021).
- [26] S. G. Rubin, A. S. Sakharov, and M. Yu. Khlopov, The Formation of primary galactic nuclei during phase transitions in the early universe, *J. Exp. Theor. Phys.* **92**, 921 (2001).
- [27] M. Yu. Khlopov, S. G. Rubin, and A. S. Sakharov, Strong primordial inhomogeneities and galaxy formation, *arXiv: astro-ph/0202505*.
- [28] M. Yu. Khlopov, S. G. Rubin, and A. S. Sakharov, Primordial structure of massive black hole clusters, *Astropart. Phys.* **23**, 265 (2005).
- [29] S. Shandera, D. Jeong, and H. S. G. Gebhardt, Gravitational Waves from Binary Mergers of Subsolar Mass Dark Black Holes, *Phys. Rev. Lett.* **120**, 241102 (2018).
- [30] D. Singh, M. Ryan, R. Magee, T. Akhter, S. Shandera, D. Jeong, and C. Hanna, A gravitational-wave limit on the Chandrasekhar mass of dark matter, *Phys. Rev. D* **104**, 044015 (2021).
- [31] B. Dasgupta, R. Laha, and A. Ray, Low Mass Black Holes from Dark Core Collapse, *Phys. Rev. Lett.* **126**, 141105 (2021).
- [32] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), Search for Subsolar-Mass Ultracompact Binaries in Advanced LIGO's First Observing Run, *Phys. Rev. Lett.* **121**, 231103 (2018).
- [33] B. P. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), Search for Subsolar Mass Ultracompact Binaries in Advanced LIGO's Second Observing Run, *Phys. Rev. Lett.* **123**, 161102 (2019).
- [34] K. S. Phukon, G. Baltus, S. Caudill, S. Clesse, A. Depasse, M. Fays, H. Fong, S. J. Kapadia, R. Magee, and A. J. Tanasijczuk, The hunt for sub-solar primordial black holes in low mass ratio binaries is open, *arXiv:2105.11449*.
- [35] A. H. Nitz and Y.-F. Wang, Search for gravitational waves from the coalescence of sub-solar mass and eccentric compact binaries, *Astrophys. J.* **915**, 54 (2021).
- [36] A. H. Nitz and Y.-F. Wang, Search for Gravitational Waves from High-Mass-Ratio Compact-Binary Mergers of Stellar Mass and Subsolar Mass Black Holes, *Phys. Rev. Lett.* **126**, 021103 (2021).
- [37] T. Nakamura, M. Sasaki, T. Tanaka, and K. S. Thorne, Gravitational waves from coalescing black hole macho binaries, *Astrophys. J.* **487**, L139 (1997).
- [38] H. Nishikawa, E. D. Kovetz, M. Kamionkowski, and J. Silk, Primordial-black-hole mergers in dark-matter spikes, *Phys. Rev. D* **99**, 043533 (2019).
- [39] Z.-C. Chen and Q.-G. Huang, Merger rate distribution of primordial-black-hole binaries, *Astrophys. J.* **864**, 61 (2018).
- [40] Y. Ali-Haïmoud, E. D. Kovetz, and M. Kamionkowski, Merger rate of primordial black-hole binaries, *Phys. Rev. D* **96**, 123523 (2017).
- [41] M. Raidal, C. Spethmann, V. Vaskonen, and H. Veermäe, Formation and evolution of primordial black hole binaries in the early Universe, *J. Cosmol. Astropart. Phys.* **02** (2019) 018.
- [42] S. Fakhry, J. T. Firouzjaee, and M. Farhoudi, Primordial black hole merger rate in ellipsoidal-collapse dark matter halo models, *Phys. Rev. D* **103**, 123014 (2021).
- [43] S. Fakhry, M. Naseri, J. T. Firouzjaee, and M. Farhoudi, Primordial black hole merger rate in self-interacting dark matter halo models, *arXiv:2106.06265*.
- [44] Y.-F. Wang and A. H. Nitz, Prospects for detecting gravitational waves from eccentric sub-solar mass compact binaries, *Astrophys. J.* **912**, 53 (2021).
- [45] S. A. Usman *et al.*, The PyCBC search for gravitational waves from compact binary coalescence, *Classical Quantum Gravity* **33**, 215004 (2016).
- [46] A. H. Nitz, I. W. Harry, J. L. Willis, C. M. Biwer, D. A. Brown, L. P. Pekowsky, T. Dal Canton, A. R. Williamson, T. Dent, C. D. Capano, T. J. Massinger, A. K. Lenon, A. B. Nielsen, and M. Cabero, PyCBC Software, <https://github.com/gwastro/pycbc> (2018).
- [47] B. Allen, W. G. Anderson, P. R. Brady, D. A. Brown, and J. D. E. Creighton, FINDCHIRP: An Algorithm for detection of gravitational waves from inspiraling compact binaries, *Phys. Rev. D* **85**, 122006 (2012).
- [48] A. H. Nitz, T. Dent, T. D. Canton, S. Fairhurst, and D. A. Brown, Detecting binary compact-object mergers with gravitational waves: Understanding and improving the sensitivity of the PyCBC search, *Astrophys. J.* **849**, 118 (2017).
- [49] A. H. Nitz, Distinguishing short duration noise transients in LIGO data to improve the PyCBC search for gravitational waves from high mass binary black hole mergers, *Classical Quantum Gravity* **35**, 035016 (2018).
- [50] B. Allen, A  $\chi^2$  time-frequency discriminator for gravitational wave detection, *Phys. Rev. D* **71**, 062001 (2005).
- [51] G. S. Davies, T. Dent, M. Tápai, I. Harry, C. McIsaac, and A. H. Nitz, Extending the PyCBC search for gravitational waves from compact binary mergers to a global network, *Phys. Rev. D* **102**, 022004 (2020).
- [52] S. Mozzon, L. K. Nuttall, A. Lundgren, T. Dent, S. Kumar, and A. H. Nitz, Dynamic normalization for compact binary coalescence searches in non-stationary noise, *Classical Quantum Gravity* **37**, 215014 (2020).
- [53] M. Was, M.-A. Bizouard, V. Brisson, F. Cavalier, M. Davier *et al.*, On the background estimation by time slides in a network of gravitational wave detectors, *Classical Quantum Gravity* **27**, 015005 (2010).
- [54] C. Capano, T. Dent, C. Hanna, M. Hendry, C. Messenger, Y.-M. Hu, and J. Veitch, Systematic errors in estimation of gravitational-wave candidate significance, *Phys. Rev. D* **96**, 082002 (2017).
- [55] B. P. Abbott *et al.* (Virgo, LIGO Scientific Collaborations), Characterization of transient noise in Advanced LIGO relevant to gravitational wave signal GW150914, *Classical Quantum Gravity* **33**, 134001 (2016).
- [56] B. S. Sathyaprakash and S. V. Dhurandhar, Choice of filters for the detection of gravitational waves from coalescing binaries, *Phys. Rev. D* **44**, 3819 (1991).
- [57] S. Droz, D. J. Knapp, E. Poisson, and B. J. Owen, Gravitational waves from inspiraling compact binaries: Validity of

- the stationary-phase approximation to the fourier transform, *Phys. Rev. D* **59**, 124016 (1999).
- [58] L. Blanchet, Gravitational radiation from post-Newtonian sources and inspiralling compact binaries, *Living Rev. Relativity* **5**, 3 (2002).
- [59] G. Faye, S. Marsat, L. Blanchet, and B. R. Iyer, The third and a half post-Newtonian gravitational wave quadrupole mode for quasi-circular inspiralling compact binaries, *Classical Quantum Gravity* **29**, 175004 (2012).
- [60] B. Moore and N. Yunes, A 3PN Fourier domain waveform for non-spinning binaries with moderate eccentricity, *Classical Quantum Gravity* **36**, 185003 (2019).
- [61] B. Moore and N. Yunes, Data analysis implications of moderately eccentric gravitational waves, *Classical Quantum Gravity* **37**, 225015 (2020).
- [62] B. Moore, T. Robson, N. Loutrel, and N. Yunes, Towards a Fourier domain waveform for non-spinning binaries with arbitrary eccentricity, *Classical Quantum Gravity* **35**, 235006 (2018).
- [63] I. W. Harry, B. Allen, and B. S. Sathyaprakash, A stochastic template placement algorithm for gravitational wave data analysis, *Phys. Rev. D* **80**, 104014 (2009).
- [64] T. Chiba and S. Yokoyama, Spin distribution of primordial black holes, *Prog. Theor. Exp. Phys.* (2017), 083E01.
- [65] V. De Luca, V. Desjacques, G. Franciolini, A. Malhotra, and A. Riotto, The initial spin probability distribution of primordial black holes, *J. Cosmol. Astropart. Phys.* **05** (2019) 018.
- [66] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, The evolution of primordial black holes and their final observable spins, *J. Cosmol. Astropart. Phys.* **04** (2020) 052.
- [67] M. Mirbabayi, A. Gruzinov, and J. Noreña, Spin of primordial black holes, *J. Cosmol. Astropart. Phys.* **03** (2020) 017.
- [68] K. Postnov, A. Kuranov, and N. Mitichkin, Spins of black holes in coalescing compact binaries, *Phys. Usp.* **62**, 1153 (2019).
- [69] M. Vallisneri, J. Kanner, R. Williams, A. Weinstein, and B. Stephens, The LIGO open science center, *J. Phys. Conf. Ser.* **610**, 012021 (2015).
- [70] R. Abbott *et al.* (LIGO Scientific, Virgo Collaborations), Open data from the first and second observing runs of Advanced LIGO and Advanced Virgo, *SoftwareX* **13**, 100658 (2021).
- [71] R. Biswas, P. R. Brady, J. D. E. Creighton, and S. Fairhurst, The loudest event statistic: General formulation, properties and applications, *Classical Quantum Gravity* **26**, 175009 (2009).
- [72] V. Vaskonen and H. Veermäe, Lower bound on the primordial black hole merger rate, *Phys. Rev. D* **101**, 043015 (2020).
- [73] A. H. Nitz and Y.-F. Wang, Data release for this paper, v1.0, 2021 <https://github.com/gwastro/subsolar-o3a-search>.
- [74] <https://www.gw-openscience.org>.