

# EXISTENCE OF MINIMAL HYPERSURFACES WITH NON-EMPTY FREE BOUNDARY FOR GENERIC METRICS

ZHICHAO WANG

ABSTRACT. For almost all Riemannian metrics (in the  $C^\infty$  Baire sense) on a compact manifold with boundary  $(M^{n+1}, \partial M)$ ,  $3 \leq (n+1) \leq 7$ , we prove that, for any open subset  $V$  of  $\partial M$ , there exists a compact, properly embedded free boundary minimal hypersurface intersecting  $V$ .

## 1. INTRODUCTION

In 1960s, Almgren [1, 2] initiated a variational theory to find minimal submanifolds in any compact manifolds with boundary. For a closed manifold  $M^{n+1}$ , the regularity of such hypersurfaces was improved by Pitts [20] for  $n \leq 5$ , and Schoen-Simon [21] for  $n = 6$ . Very recently, Li and Zhou finished this program for a general compact manifold with nonempty boundary in [13], in which they proved that every compact manifold with boundary admits a smooth compact minimal hypersurface with (possibly empty) free boundary. This result left widely open the following well-known question::

**Question 1.1.** *Does every compact manifold with non-empty boundary admit a minimal hypersurface with non-empty free boundary?*

We point out that there are similar questions in any free boundary variational theory. In particular, in the mapping approach by Fraser [6], Lin-Sun-Zhou [14], and Lauren-Petradis [12], it was not known whether their free boundary minimal surfaces have nontrivial boundary.

In this paper, we solve this problem in generic scenarios and prove a much stronger property:  $M$  admits infinitely many embedded minimal hypersurfaces with non-empty free boundary.

**Theorem 1.2.** *Let  $(M^{n+1}, \partial M)$  be a compact manifold of dimension  $3 \leq (n+1) \leq 7$ . Then for a  $C^\infty$ -generic Riemannian metric  $g$  on  $(M, \partial M)$ , the union of boundaries of all smooth, embedded, free boundary minimal hypersurfaces is dense in  $\partial M$ .*

We remark that a compact manifold with non-negative Ricci curvature and strictly convex boundary has no closed minimal hypersurface by [5, Lemma 2.2]. Therefore, by Marques-Neves [17] and Li-Zhou [13], it is known that there exist infinitely many properly embedded free boundary minimal hypersurfaces in such ambient manifolds.

For a generic metric on  $(M, \partial M)$ , the author together with Guang, Li and Zhou proved the density of free boundary minimal hypersurfaces in [8, Theorem 1.3]. Making

use of a maximum principle by White [24], such denseness gives that  $M$  contains minimal hypersurfaces with non-empty boundary by merely assuming strict mean convexity at one point of the boundary  $\partial M$  for a generic metric; see [8]. However, without any topological or curvature assumptions, it is in general very difficult to prevent the free boundary components from degenerating in the limit process (see e.g. [3, 9]). Our theorem in this paper greatly improves this result by dropping off mean convexity assumption at one point.

The denseness result in [8, Theorem 1.3] can be seen as a natural free boundary analog of [11]. The key ingredient of [11] by Irie, Marques and Neves is the Weyl law for the volume spectrum proved by Liokumovich, Marques and Neves in [15]. The volume spectrum of a compact Riemannian manifold with boundary  $(M^{n+1}, g)$  is a nondecreasing sequence of numbers  $\{\omega_k(M; g) : k \in \mathbb{N}\}$  defined variationally by performing a min-max procedure for the area functional over multiparameter sweepouts. The first estimates for these numbers were proven by Gromov [7] in the late 1980s (see also [10]). A direct corollary of Weyl Law they used is that, for  $k$  large enough,  $\omega_k(M; g) \neq \omega_k(M; g')$  whenever  $\text{Vol}(M, g) \neq \text{Vol}(M, g')$ .

Another observation by Irie, Marques and Neves is that such spectrum depends continuously on the metrics of  $M$ ; see [11, Lemma 2.1] and [19, Lemma 1]. Applying this, they showed that continuous perturbations in an open set must create new minimal hypersurfaces intersecting that set.

In this paper, we also borrow the idea from Irie-Marques-Neves [11]. However, the original perturbation would only produce new free boundary minimal hypersurfaces intersecting *an open set*, but not an  $n$ -dimensional subset, that we need to consider here. To overcome this new challenge, we perturb the metric  $g$  around a boundary point in *a special way* so that a hypersurface whose boundary does not intersect the prescribed subset of  $\partial M$  can also be regarded a hypersurface in  $(M, \partial M, g)$ . Recall that Weyl law in [15] gives that for large  $k$ ,  $\omega_k$  will change continuously if the volume of  $M$  is changed under the perturbation. From these two observations, we are able to prove that such a special perturbation would produce new minimal hypersurfaces with free boundary intersecting the prescribed subset of  $\partial M$ .

We finish the introduction with the idea of the construction of the special perturbation. Making use of the cut-off trick, the unit inward normal vector field of  $\partial M$  can be extended to the whole  $M$ . Also, by multiplying another cut-off function, we can always construct a vector field whose support is close to our prescribed open set of  $\partial M$ . Such a vector field would give a one-parameter family of diffeomorphisms (not surjective) of  $M$ . Then the pull back metric given by such family is the desired perturbation since it is isometric to a subset of  $M$  with the original metric. We refer to Proposition 3.1 for more details.

**Acknowledgement.** The author would like to thank Prof. Xin Zhou for bringing this problem to us and many helpful discussion.

## 2. PRELIMINARIES

Let  $(M^{n+1}, g)$  be a smooth compact connected Riemannian manifold with nonempty boundary  $\partial M$  and  $3 \leq (n+1) \leq 7$ . Moreover,  $M$  can always be embedded to a closed Riemannian manifold  $\widetilde{M}$  which has the same dimension with  $M$ . We can also assume that  $\widetilde{M}$  is isometrically embedded in some  $\mathbb{R}^L$  for  $L$  large enough.

**2.1. Geometric measure theory.** We now recall some basic notations in geometric measure theory; see [13].

We use  $\mathcal{V}_k(M)$  to denote the closure of the space of  $k$ -dimensional rectifiable varifolds in  $\mathbb{R}^L$  with support contained in  $M$ . Let  $\mathcal{R}_k(M; \mathbb{Z}_2)$  (resp.  $\mathcal{R}_k(\partial M; \mathbb{Z}_2)$ ) be the space of  $k$ -dimensional modulo two flat chains of finite mass in  $\mathbb{R}^L$  which are supported in  $M$  (resp. in  $\partial M$ ). Denote by  $\text{spt } T$  the support of  $T \in \mathcal{R}_k(M; \mathbb{Z}_2)$ . Given any  $T \in \mathcal{R}_k(M; \mathbb{Z}_2)$ , denote by  $|T|$  and  $\|T\|$  the integer rectifiable varifold and the Radon measure in  $M$  associated with  $T$ , respectively. The mass norm and the flat metric on  $\mathcal{R}_k(M; \mathbb{Z}_2)$  are denoted by  $\mathbf{M}$  and  $\mathcal{F}$  respectively; see [4]. Set

$$Z_k(M, \partial M; \mathbb{Z}_2) = \{T \in \mathcal{R}_k(M; \mathbb{Z}_2) : \text{spt}(\partial T) \subset \partial M\}.$$

We say that  $T, S \in Z_k(M, \partial M; \mathbb{Z}_2)$  are equivalent if  $T - S \in \mathcal{R}_k(\partial M; \mathbb{Z}_2)$ . We use  $\mathcal{Z}_k(M, \partial M; \mathbb{Z}_2)$  to denote the space of all such equivalent classes; see [8, Section 3] for the equivalence with the formulation using integer rectifiable currents in [13].

The flat metric and the mass norm in the space of relative cycles are defined, respectively, as

$$\mathcal{F}(\tau_1, \tau_2) = \inf\{\mathcal{F}(T) : T \in \tau\}, \quad \mathbf{M}(\tau) = \inf\{\mathbf{M}(T) : T \in \tau\}.$$

The connected component of  $\mathcal{Z}_n(M, \partial M; \mathbb{Z}_2)$  containing 0 is weakly equivalent to  $\mathbb{R}\mathbb{P}^\infty$  by Almgren [1] (see also [15, §2.5] and [8, Section 3]). Denote by  $\bar{\lambda}$  the generator of  $H^1(\mathcal{Z}_n(M, \partial M; \mathbb{Z}_2); \mathbb{Z}_2) = \mathbb{Z}_2$ .

**2.2. Auxiliary Lemmas.** In this part, we introduce some Lemmas in [8, 11, 19].

Let  $X$  be a finite dimensional simplicial complex. A continuous map  $\Phi : X \rightarrow \mathcal{Z}_n(M, \partial M; \mathbb{Z}_2)$  is called a  $k$ -sweepout if

$$\Phi^*(\bar{\lambda}^k) \neq 0 \in H^k(X; \mathbb{Z}_2).$$

We denote by  $\mathcal{P}_k(M)$  the set of all  $k$ -sweepouts that have *no concentration of mass*, meaning that

$$\limsup_{r \rightarrow 0} \{\mathbf{M}(\Phi(x) \cap B_r(p)) : x \in X, p \in M\} = 0.$$

**Definition 2.1.** The  $k$ -width of  $(M, \partial M; g)$  is defined as

$$\omega_k(M) := \inf_{\Phi \in \mathcal{P}_k(M)} \sup\{\mathbf{M}(\Phi(x)) : x \in \text{dmn}(\Phi)\},$$

where  $\text{dmn}(\Phi)$  is the domain of  $\Phi$ .

For any compact Riemannian manifold with boundary  $(M, \partial M, g)$ , the sequence  $\{\omega_p(M)\}$  satisfies Weyl Law, which is proven by Liokumovich, Marques and Neves.

**Theorem 2.2** (Weyl Law for the Volume Spectrum; [15]). *There exists a constant  $\alpha(n)$  such that, for every compact Riemannian manifold  $(M^{n+1}, \partial M, g)$  with (possibly empty) boundary, we have*

$$\lim_{k \rightarrow \infty} \omega_k(M; g) k^{-\frac{1}{n+1}} = \alpha(n) \text{Vol}(M, g)^{\frac{n}{n+1}}.$$

Irie-Marques-Neves [11, Lemma 2.1] proved that  $\omega_k(M; g)$  depends continuously on metrics. The following is an improved version given by Marques, Neves and Song.

**Lemma 2.3** ([11, Lemma 2.1; 19, Lemma 1]). *Let  $g_0$  be a  $C^2$  Riemannian metric on  $(M, \partial M)$ , and let  $C_1 < C_2$  be positive constants. Then there exists  $K = K(g_0, C_1, C_2) > 0$  such that*

$$|p^{-\frac{1}{n+1}} \omega_p(M; g) - p^{-\frac{1}{n+1}} \omega_p(M; g')| \leq K \cdot |g - g'|_{g_0}$$

for any  $C^2$  metric  $g, g' \in \{h; C_1 g_0 \leq h \leq C_2 g_0\}$  and any  $p \in \mathbb{N}$ .

Inspired by Marques-Neves [16], the author with Guang, Li, and Zhou (see [8, Theorem 2.1]) gave a general index estimate for min-max minimal hypersurfaces with free boundary. Combining with a compactness theorem in [9] by the author and Guang and Zhou, we also proved in [8] that the  $k$ -width is realized by the area (counting multiplicities) of min-max free boundary minimal hypersurfaces.

**Proposition 2.4** ([8, Proposition 7.3; 11, Proposition 2.2]). *Suppose  $3 \leq (n+1) \leq 7$ . Then for each  $k \in \mathbb{N}$ , there exist a finite disjoint collection  $\{\Sigma_1, \dots, \Sigma_N\}$  of almost properly embedded free boundary minimal hypersurfaces in  $(M, \partial M, g)$ , and integers  $\{m_1, \dots, m_N\} \subset \mathbb{N}$ , such that*

$$\omega_k(M; g) = \sum_{j=1}^N m_j \text{Area}_g(\Sigma_j) \quad \text{and} \quad \sum_{j=1}^N \text{index}(\Sigma_j) \leq k.$$

*Remark 2.5.* In a recent exciting work [25], X. Zhou proved that, for a bumpy metric on a closed manifold, each  $m_j$  equals to 1, which is conjectured by Marques and Neves in [16]. Based on this Multiplicity One Theorem, Marques-Neves [18] proved that the index is in fact equals to  $k$  for min-max minimal hypersurfaces realizing  $\omega_k$ .

### 3. PROOF OF THEOREM 1.2

Let  $(M^{n+1}, \partial M)$  be a compact manifold with boundary and  $3 \leq (n+1) \leq 7$ . Let  $\mathcal{M}$  be the space of all smooth Riemannian metrics on  $M$ , endowed with the smooth topology. Suppose that  $V \subset \partial M$  is a non-empty open set. Let  $\mathcal{M}_V$  be the set of metrics  $g \in \mathcal{M}$  such that there exists a non-degenerate, properly embedded free boundary minimal hypersurface  $\Sigma$  in  $(M, \partial M, g)$  whose boundary intersects  $V$ .

We approach the theorem by proving the following proposition.

**Proposition 3.1.** *For any compact manifold  $(M, \partial M)$  and any open subset  $V \subset \partial M$ ,  $\mathcal{M}_V$  is open and dense in  $\mathcal{M}$  in the smooth topology.*

*Proof.* Let  $g \in \mathcal{M}_V$  and  $\Sigma$  be like in the statement of the proposition. Following the step by Irie-Marques-Neves in [11], we first show the openness of  $\mathcal{M}_V$ . Note that  $\Sigma$

is a properly embedded, then the Structure Theorem of White [23, Theorem 2.1] (see [3, Theorem 35] for a version on free boundary minimal hypersurfaces) also gives that for every Riemannian metric  $g'$  sufficiently close to  $g$ , there exists a unique nondegenerate properly embedded free boundary minimal hypersurface  $\Sigma'$  close to  $\Sigma$ . This implies  $\mathcal{M}_V$  is open.

It remains to show the set  $\mathcal{M}_V$  is dense. Let  $g$  be an arbitrary smooth Riemannian metric on  $(M, \partial M)$  and  $\mathcal{V}$  be an arbitrary neighborhood of  $g$  in the  $C^\infty$  topology. By the Bumpy Metrics Theorem ([3, Theorem 9; 23, Theorem 2.1]), there exists  $g' \in \mathcal{V}$  such that every compact, almost properly embedded free boundary minimal hypersurface with respect to  $g'$  is nondegenerate.

Since  $g'$  is bumpy, then by [8, Proposition 5.3] (see also [9, 22]), the space of almost embedded free boundary minimal hypersurfaces with  $\text{Area} \leq \Lambda$  and  $\text{index} \leq I$  is countable with respect to  $g'$  for all  $\Lambda > 0$  and  $I \geq 0$ . Therefore, the set

$$\mathcal{C} := \left\{ \sum_{j=1}^N m_j \text{Area}_{g'}(\Sigma_j) \mid \begin{array}{l} N \in \mathbb{N}, \{m_j\} \subset \mathbb{N}, \{\Sigma_j\} \text{ disjoint collection of almost} \\ \text{properly embedded free boundary minimal} \\ \text{hypersurfaces in } (M, \partial M, g') \end{array} \right\}$$

is countable.

Let  $U$  be an open set of  $M$  such that  $\bar{U} \cap \partial M \subset V$  is non-empty. Let  $X$  be a vector field on  $M$  so that  $\text{spt } X \subset U$  and for  $x \in \partial M$  satisfying  $X(x) \neq 0$ ,  $X(x)/|X(x)|$  is the outward unit normal vector of  $\partial M$ . Denote by  $(F_t)_{0 \leq t \leq 1}$  a family of diffeomorphisms of  $M$  generated by  $X$ . Set

$$g_t = F_t^* g' \quad \text{and} \quad M_t = F_t(M).$$

Then  $(M, \partial M, g_t)$  is isometric to  $(M_t, \partial M_t, g')$ . Note that we can take  $\delta > 0$  so that  $g_t \in \mathcal{V}$  for all  $t \in [0, \delta]$ .

**Claim 1.** *Let  $\Gamma$  be an integral varifold in  $M$  whose support is a free boundary minimal hypersurface  $\Sigma$  (possibly disconnected) in  $(M, \partial M, g_t)$ . Assuming that  $\partial \Sigma \cap V = \emptyset$ , then  $\mathbf{M}(\Gamma) \in \mathcal{C}$ .*

*Proof of Claim 1.* By the definition of  $g_t$ ,  $\Sigma$  can be seen as a free boundary minimal hypersurface in  $(M_t, \partial M_t, g')$  so that  $\partial \Sigma \cap F_t(V) = \emptyset$ . Thus,  $\Sigma$  is also a free boundary minimal hypersurface in  $(M, \partial M, g')$ . It follows that  $\mathbf{M}(\Gamma) \in \mathcal{C}$  (counted with multiplicities).  $\square$

**Claim 2.** *There exist  $t_1 \in [0, \delta]$  and an almost properly embedded free boundary minimal hypersurface  $(\Sigma_1, \partial \Sigma_1) \subset (M, \partial M, g_{t_1})$  satisfying  $\partial \Sigma_1 \cap V \neq \emptyset$ .*

*Proof of Claim 2.* Suppose not, then for all  $t \in [0, \delta]$ , all the almost properly embedded minimal hypersurfaces in  $(M_t, \partial M_t, g')$  have no boundaries in  $V$ . Recall that Proposition 2.4 gives that  $\omega_k(M; g_t)$  is realized by the area of such hypersurfaces. Together with Claim 1, we conclude that

$$\omega_k(M_t; g') \in \mathcal{C} \quad \text{for all } t \in [0, \delta] \text{ and } k \in \mathbb{N}.$$

On the other hand, the Weyl law (see Theorem 2.2) implies that  $\omega_k(M; g_\delta) < \omega_k(M; g')$  for  $k$  large enough. The Lemma 2.3 deduces that  $\omega_k(M; g_t)$  is continuous, which leads to a contradiction with that  $\omega_k(M; g_t)$  lies in a countable set. The proof is finished.  $\square$

Thus we have proved that for some  $t_1 \in [0, \delta]$ , there exists an almost properly embedded free boundary minimal hypersurface  $(\Sigma_1, \partial\Sigma_1) \subset (M, \partial M; g_{t_1})$  satisfying  $\partial\Sigma_1 \cap V \neq \emptyset$ . Then by [11, Proposition 2.3] (see also [8, Proposition 7.6; 19, Lemma 4]),  $g_{t_1}$  can be perturbed to  $g'' \in \mathcal{V}$  so that  $(M, \partial M, g'')$  contains an almost properly embedded, non-degenerate, free boundary minimal hypersurfaces  $\Sigma''$  whose boundary intersects  $V$ . Finally, [8, Proposition 7.7] would allow us to perturb  $g''$  to  $\tilde{g} \in \mathcal{V}$  and  $\Sigma''$  is a properly embedded free boundary minimal hypersurface in  $(M, \partial M, \tilde{g})$ . This implies that  $\tilde{g} \in \mathcal{M}_V$  and we are done.  $\square$

Now we are ready to prove Theorem 1.2. The proof is the same with that of [11, Main theorem].

*Proof of Theorem 1.2.* Let  $\{V_i\}$  be a countable basis of  $\partial M$ . Since, by Proposition 3.1, each  $\mathcal{M}_{V_i}$  is open and dense in  $\mathcal{M}$ , and hence the set  $\bigcap_i \mathcal{M}_{V_i}$  is  $C^\infty$  Baire-generic in  $\mathcal{M}$ . This finishes the proof.  $\square$

#### REFERENCES

- [1] Frederick Justin Almgren Jr., *The homotopy groups of the integral cycle groups*, *Topology* **1** (1962), 257–299. MR0146835
- [2] ———, *The theory of varifolds*, Mimeographed notes, Princeton (1965), 12.
- [3] Lucas Ambrozio, Alessandro Carlotto, and Ben Sharp, *Compactness analysis for free boundary minimal hypersurfaces*, *Calc. Var. Partial Differential Equations* **57** (2018), no. 1, 57:22. MR3740402
- [4] Herbert Federer, *Geometric measure theory*, *Die Grundlehren der mathematischen Wissenschaften*, Band 153, Springer-Verlag New York Inc., New York, 1969. MR0257325
- [5] Ailana Fraser and Martin Man-chun Li, *Compactness of the space of embedded minimal surfaces with free boundary in three-manifolds with nonnegative Ricci curvature and convex boundary*, *J. Differential Geom.* **96** (2014), no. 2, 183–200. MR3178438
- [6] Ailana M. Fraser, *On the free boundary variational problem for minimal disks*, *Comm. Pure Appl. Math.* **53** (2000), no. 8, 931–971. MR1755947
- [7] Mikhael Gromov, *Isoperimetry of waists and concentration of maps*, *Geom. Funct. Anal.* **13** (2003), no. 1, 178–215. MR1978494
- [8] Qiang Guang, Martin Man-chun Li, Zhichao Wang, and Xin Zhou, *Min-max theory for free boundary minimal hypersurfaces II—General Morse index bounds and applications*, arXiv preprint arXiv:1907.12064 (2019).
- [9] Qiang Guang, Zhichao Wang, and Xin Zhou, *Compactness and generic finiteness for free boundary minimal hypersurfaces (I)*, arXiv preprint arXiv:1803.01509 (2018).
- [10] Larry Guth, *Minimax problems related to cup powers and Steenrod squares*, *Geom. Funct. Anal.* **18** (2009), no. 6, 1917–1987. MR2491695
- [11] Kei Irie, Fernando Marques, and André Neves, *Density of minimal hypersurfaces for generic metrics*, *Ann. of Math. (2)* **187** (2018), no. 3, 963–972. MR3779962
- [12] Paul Laurain and Romain Petrides, *Existence of min-max free boundary disks releasing the width of a manifold*, to appear in *Advances in Mathematics*, arXiv preprint arXiv:1806.04665 (2018).
- [13] Martin Li and Xin Zhou, *Min-max theory for free boundary minimal hypersurfaces I-regularity theory*, to appear in *JDG*, arXiv:1611.02612 (2016).
- [14] Longzhi Lin, Ao Sun, and Xin Zhou, *Min-max minimal disks with free boundary in Riemannian manifolds*, to appear in *Geom. Topol.*, arXiv preprint arXiv:1806.04664 (2018).
- [15] Yevgeny Liokumovich, Fernando C. Marques, and André Neves, *Weyl law for the volume spectrum*, *Ann. of Math. (2)* **187** (2018), no. 3, 933–961. MR3779961
- [16] Fernando C. Marques and André Neves, *Morse index and multiplicity of min-max minimal hypersurfaces*, *Camb. J. Math.* **4** (2016), no. 4, 463–511. MR3572636

- [17] ———, *Existence of infinitely many minimal hypersurfaces in positive Ricci curvature*, Invent. Math. **209** (2017), no. 2, 577–616. MR3674223
- [18] ———, *Morse index of multiplicity one min-max minimal hypersurfaces*, arXiv preprint arXiv:1803.04273 (2018).
- [19] Fernando C. Marques, André Neves, and Antoine Song, *Equidistribution of minimal hypersurfaces for generic metrics*, Invent. Math. **216** (2019), no. 2, 421–443. MR3953507
- [20] Jon T. Pitts, *Existence and regularity of minimal surfaces on Riemannian manifolds*, Mathematical Notes, vol. 27, Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981. MR626027
- [21] Richard Schoen and Leon Simon, *Regularity of stable minimal hypersurfaces*, Comm. Pure Appl. Math. **34** (1981), no. 6, 741–797. MR634285 (82k:49054)
- [22] Zhichao Wang, *Compactness and generic finiteness for free boundary minimal hypersurfaces (II)*, arXiv preprint arXiv:1906.08485 (2019).
- [23] Brian White, *The space of minimal submanifolds for varying Riemannian metrics*, Indiana Univ. Math. J. **40** (1991), no. 1, 161–200. MR1101226 (92i:58028)
- [24] ———, *The maximum principle for minimal varieties of arbitrary codimension*, Comm. Anal. Geom. **18** (2010), no. 3, 421–432. MR2747434
- [25] Xin Zhou, *On the multiplicity one conjecture in min-max theory*, arXiv preprint arXiv:1901.01173 (2019).

MAX-PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, 53111 BONN, GERMANY  
E-mail address: wangzhichaonk@gmail.com