ELSEVIER

Contents lists available at ScienceDirect

NeuroImage



journal homepage: www.elsevier.com/locate/neuroimage

Mapping the principal gradient onto the corpus callosum

Patrick Friedrich^{a,b,*}, Stephanie J. Forkel^{a,b,c}, Michel Thiebaut de Schotten^{a,b}

^a Brain Connectivity and Behaviour Laboratory, Sorbonne Universities, Paris, France

^b Groupe d'Imagerie Neurofonctionelle, CEA, Univ. Bordeaux, CNRS, IMN, UMR 5293, Bordeaux F-33000, France

^c Centre for Neuroimaging Sciences, Department of Neuroimaging, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

ABSTRACT

Gradients capture some of the variance of the resting-state functional magnetic resonance imaging (rsfMRI) signal. Amongst these, the principal gradient depicts a functional processing hierarchy that spans from sensory-motor cortices to regions of the default-mode network. While the cortex has been well characterised in terms of gradients little is known about its underlying white matter. For instance, comprehensive mapping of the principal gradient on the largest white matter tract, the corpus callosum, is still missing. Here, we mapped the principal gradient onto the midsection of the corpus callosum using the 7T human connectome project dataset. We further explored how quantitative measures and variability in callosal midsection connectivity relate to the principal gradient values. In so doing, we demonstrated that the extreme values of the principal gradient are located within the callosal genu and the posterior body, have lower connectivity variability but a larger spatial extent along the midsection of the corpus callosum than mid-range values. Our results shed light on the relationship between the brain's functional hierarchy and the corpus callosum. We further speculate about how these results may bridge the gap between functional hierarchy, brain asymmetries, and evolution.

1. Introduction

Cognitive neuroscience assumes that cognition is embedded within and constraint by the organization of the brain. The advent of neuroimaging methods granted novel insights into the functional organization of the living human brain. Resting-state functional connectivity (rsfMRI), in particular, has proven useful for mapping multiple functional networks without relying on participants to perform functional tasks in the scanner (Smith et al., 2009; Power et al., 2011; Biswal et al., 1995). rsfMRI networks reveal different time courses and spatially correspond to tasks-related functional activations (Tavor et al., 2016). This spatial correspondence suggests an underlying organization that adheres to fundamental principles. For instance, Mesulam (1998) proposes that brain regions can be organised along a gradient ranging from sensorymotor to higher-order brain processes. In line with Mesulam, most of the variance of rsfMRI resembles a gradient that spans from sensorymotor cortices to regions of the default-mode network (i.e. the principal gradient, Margulies et al., 2016). Sensory-motor cortices are primary cortices (i.e. idiotypic areas Mesulam, 1998) and the default-mode network has been associated with transmodal higher-order functions (Margulies et al., 2016; Alves et al., 2019). The principal gradient may thus represent one fundamental principle driving a hierarchical organisation of cognitive functions (Huntenburg et al., 2018).

While this principal gradient is putatively the same for both hemispheres (Margulies et al., 2016) little is known on the connections between the left and the right principal gradient. With 200 to 300 millions axons (Tomasch 1954), the largest white matter commissure in the primate brain is the corpus callosum (Burdach, 1819). Other routes of interhemispheric interactions are also possible (e.g. anterior commissure, Risse et al., 1978), however surgically severing callosal connections severely interrupts interhemispheric functional connectivity in monkeys (Johnston et al., 2008; O'Reilly et al., 2013) and humans (Roland et al., 2017). Concordantly, studies in split-brain patients present with severe behavioural manifestations such as Alien hand syndrome (Akelaitis 1945; Bogen, 1993), hemispatial neglect (Kashiwagi et al., 1990; Heilman and Adams, 2003; Lausberg et al., 2003; Park et al., 2006; Tomaiuolo et al., 2010), optic aphasia (Freund, 1889; Luzzatti et al., 1998) or pure alexia (Dejerine, 1892; Geschwind and Fusillo, 1966) amongst others. The corpus callosum is hence considered critical for the collaborative integration of functions relying on both hemispheres. Despite its crucial role in functions of various hierarchical order, current parcellation approaches divide the corpus callosum based on its cytoarchitecture, geometry and topological projections, with little regard to the callosums functional organization.

Several methods have been proposed to parcellate or segment the corpus callosum. Amongst these methods, four main classifications are still applied. Post mortem studies identified anatomical features along the midsagittal section of the corpus callosum, including the rostrum, genu, anterior and posterior body, isthmus and splenium (Crosby et al., 1962; Fig. 1A). While this classification is still widely used today, it does not define clear boundaries between the different segments. A second tentatively sharper parcellation was based on histological measures of the midsagittal axonal diameter but it fails to systematically generalize across individuals (Aboitiz and Montiel, 2003; Aboitiz et al., 1992, Fig. 1B). The enforcement of a geometrical grid of seven parcels has been used to somewhat circumvent the limitations of generalisation (Wittelson, 1989; Fig. 1C). All these anatomical, histological and

* Corresponding author.

E-mail address: patrick.friedrich@rub.de (P. Friedrich).

https://doi.org/10.1016/j.neuroimage.2020.117317

Received 27 February 2020; Received in revised form 31 July 2020; Accepted 26 August 2020 Available online 31 August 2020

1053-8119/© 2020 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)



Fig. 1. Parcellations of the corpus callosum along the midsection (right view). A) Anatomical division: depicts the anatomical classification (rostrum, genu, anterior and posterior body, isthmus, splenium) based on gross anatomical features. B) Histological division: the dots in this schematic diagram indicate the range of axonal density and associate them with cortical projections to cortical areas (F: Frontal, M: Motor, SS: Somato-Sensory, A: Auditory, T/P: Temporo-Parietal and V: Visual (from Aboitiz and Montiel, 2003). C) Geometrical division: This parcellation generates seven parcels by equally diving the midsection (from Wittelson, 1989). ACC: most anterior point of the corpus callosum, PCC: most posterior point of the corpus callosum. D: Connectional division: Shows the parcellation of the cortex based on diffusion tractography (from Catani and Thiebaut de Schotten, 2012).

geometrical parcellations of the midsection of the corpus callosum do however not consider the functional organisation of the brain areas it connects.

Since the functions of the corpus callosum are defined by the areas that it connects to, tract-tracing in non-human primates (Schmahmann et al., 2009) or diffusion weighted imaging tractography in humans (Huang et al., 2005; Hofer and Frahm, 2006; Fig. 1D), is likely the most promising approach to investigate the functional organisation of the corpus callosum. However, a comprehensive mapping of the hierarchical organization of cognitive functions on the corpus callosum is still missing. Such a mapping would help to better understand the functional impact of brain lesions that disconnect the corpus callosum.

In this study we propose a novel topography of the corpus callosum based on the projection of the principal gradient onto the callosal midsection. We also explored how variability in connectivity and quantitative measures of this connectivity relate to the principal gradient values along the callosal midsection.

2. Methods

2.1. Dataset

All 7T rsfMRI and diffusion-weighted data used in this study were based on the Human Connectome Project (HCP; Van Essen et al., 2013), which is publicly available and anonymized. For rsfMRI data, we used an averaged map of the principal gradient published in Margulies et al. (2016), which can be found under: https://neurovault.org/collections/1598/. The minimal preprocessed diffusion-weighted data consisted of 164 participants (age: mean = 29.44 (3.29); females: 100) from the HCP. Participant recruitment procedures and informed consent forms, including consent to share

de-identified data, were previously approved by the Washington University Institutional Review Board.

2.2. Gradient percentage maps

We used an averaged map of the principal gradient published (Margulies et al., 2016). In Margulies et al. (2016), functional connectivity matrices were calculated from the HCP dataset and the gradient extracted using nonlinear dimensionality reduction via diffusion embedding (Coifman et al., 2005). In order to assess callosal connections to cortical regions along the principal gradient, we segregated voxels within the cortex based on their location along the principal gradient. We parcellated the principal gradient map into 100 units where each unit represents one percentage of the principal gradient. All operations were conducted using FSL (www.fmrib.ox.ac.uk/fsl). The gradient intensities range between -7.52 and +10.07. A central value of 0 denoted areas not defined along the gradient. For computational reasons, we transposed the gradient values by multiplying the minimum intensity value by -1and adding the result to all values, thus shifting all values into positive values. We subsequently masked the results with a binarized map of the principal gradient. The resulting image was masked with a cortical grey matter template (i.e. FAST applied to MNI152 template). This resulted in 100 distinct gradient percentage maps, each one presenting a 1% range of the gradient (see Fig. 2B).

2.3. Mapping the principal gradient onto the corpus callosum

In order to map the gradient along the white matter connections, we employed the *Disconnectome* within the BCBtoolkit (Foulon et al., 2018) using a study-specific HCP dataset of 164 participants to generate the underlying white matter tractograms (Karolis et al., 2019). The scanning parameters have been described previously (Vu et al., 2015). In brief,



Fig. 2. Mapping the principal gradient on the corpus callosum. A) Visualisation of the principal gradient on the cortical surface. B) Examples for percentage maps (20–40–60%) depicting associated cortical areas. C) Percentage-specific tractograms as derived from the percentage gradients maps shown in B. D) Midsagittal display of callosal connectivity based on the tractograms for the 20th, 40th and 60th gradient percentages. Callosal areas with low probabilities (red) indicate that fewer participants showed connections to the particular gradient percentage in this area. The prevalence across gradient values for 12 exemplary voxels is depicted in supplementary figure S1-A. E) Callosal gradient map (left), which was computed as the weighted average of callosal connectivity across all percentage maps of the gradient with the 7 regions of the Wittelson parcellation superimposed. The distribution of gradient values along the Wittelson parcellation scheme is shown on the right. For a hard-border classification of voxels to gradient values please see supplementary figure S1-B. F) Scatterplots showing the relation between callosal connectivity (left) or number of non-zero connectivity voxels (right) and the residual between the gradient corrected for the number of voxels in the respective gradient percentage map. By callosal connectivity, we mean interindividual probability of connection.

diffusion-weighted imaging consisted of a total of 132 near-axial slices acquired with an acceleration factor of 3 (Moeller et al., 2010), isotropic (1.05 mm3) resolution and whole head coverage with a TE of 71.2 ms and a TR of 7000 ms. These images were acquired with 65 uniformly distributed gradients in multiple Q-space shells (Caruyer et al., 2013) with 6 b0 images. The acquisition was repeated four times with a b-value of 1000 and 2000s mm-2 in pairs with left-to-right and right-to-left phaseencoding directions. The default HCP preprocessing pipeline (v3.19.0) was applied to the data (Andersson et al., 2012; Sotiropoulos et al., 2013), which consists of susceptibility-induced distortion correction (Andersson et al., 2003), TOPUP (Smith et al., 2004) and subsequent motion and geometrical distortion correction via the EDDY tool implemented in FSL.

For each gradient percentage map, the *Disconnectome* filtered tractography of the white matter tractograms. The resulting filtered tractography (i.e. streamline passing by the gradient percentage maps) were visually inspected and transformed into visitation maps (i.e. binary masque of the voxels including more than 1 streamline, Thiebaut De Schotten et al., 2011). To generate a group-level visitation map, all normalised individual visitation maps were averaged voxel-by-voxel. This group-level visitation map represents the connection probability per voxel, and thus accounts for the interindividual variability of the anatomical representation of tracts for the respective gradient percentage (see Fig. 2C).

This study focussed on the relationship between the functional gradient and the interhemispheric commissural connections of the corpus callosum. We manually delineated the corpus callosum midsection on the midsagittal slice of the MNI152 template. To project the cortical gradient onto the corpus callosum, we subsequently created a weighted average map based on the following equation:

weighted average connectivity =
$$\sum \left(\frac{\text{connectivity map}}{\sum \text{connectivity maps}} \right) * \text{ gradient \%}$$

Each of the divided visitation maps was multiplied by the respective level of the gradient (1% = times 1, 2% = times 2, etc.). This resulted in 100 weighted connectivity maps, one for each gradient percentage. To obtain one single unified callosal connectivity gradient map, we summed the weighted connectivity maps and masked the result with the midsection of the corpus callosum. The resulting weighted average callosal connectivity map is shown in Fig. 2E. The complete weighted average tractogram is available here: https://neurovault.org/collections/7797/. In addition, we provide a hard-border segmentation approach based on similarity of voxels in their connection profile across the gradient in the supplementary material. For this, we used UMAP (McInnes et al., 2018) to reduce the dimensions of the data. Subsequently, we clustered callosal voxels to separate groups via HDBSCAN (McInnes et al., 2017). Results are available as supplementary figure S1.

2.4. Statistical analyses

Statistical analyses were performed in SPSS (IBM SPSS Statistics, Version 24.0. Armonk, NY: IBM Corp.). In order to avoid spurious measurements, we truncated the distribution of the gradient percentage maps at either end to avoid including maps with less than 300 voxels (supplementary figure S2). Included percentage maps ranged from 13% to 82% (equivalent original gradient values: -5.23 to +6.90). We explored how variability in connectivity and quantitative measures of callosal connectivity relates to the principal gradient values. The connectivity probability between the corpus callosum and the gradient percentage was calculated by averaging each callosal connectivity map. The callosal space occupied by each percentage was defined as the number of non-zero voxels for each callosal connectivity map. To explore the relationship between the corpus callosum and the principal gradient, we plotted the connection probability against the residuals of the gradient value and the number of voxels of the respective gradient percentage map. To assess the relationship between the value of the gradient and the spatial extent on the corpus callosum unbiased by the number of voxels for each percentage value we regressed out the number of voxels in the scatterplot. The same procedure was applied to plot the callosal space occupied by each percentage. The two distributions were visually inspected and scrutinised with either linear or quadratic regression as applicable.

3. Results

3.1. Corpus callosum gradient map

The principal gradient projections on the corpus callosum are displayed below (Fig. 2E). Results indicate a structured distribution of the principal gradient along the midsection of the corpus callosum where the anterior corpus callosum is more likely connected to areas of higher principal gradient values compared to posterior areas. Accordingly, the negative spectrum of the principal gradient peaked in the isthmus and the posterior body, whereas the positive spectrum occupied the anterior part of the body, the genu and the rostrum. In some parts of the splenium a small peak of positive values was also evident.

3.2. Relationship between the principal gradient and callosal connectivity

Visual inspection of the data clearly indicated a U-shaped distribution. Using a curve-fit estimation, we investigated the possibility for either a linear or quadratic relationship between connection probability and the principal gradient. A linear regression model was fitted onto the distribution (Y_i = 0.229 + -0.002*X_i), which did not match the data (F_(1, 68) = 0.698, r² = 0.010, *p* = 0.406). However, a quadratic regression revealed a highly significant (r² = 0.739, *p* < 0.001; (Y_i = 0.174 + -0.006*X_i + 0.004*X²_i)) relationships between callosal connection probability and the principal gradient as illustrated in Fig. 2F, left side.

3.3. Relationship between the principal gradient and associated callosal area

Similar to the previous analysis, a curve-fit estimation was employed to investigate a linear or quadratic relationship between the number of callosal voxels that may connect to a certain gradient value. A linear regression model ($Y_i = 840.029 + 5.280^*X_i$) failed to reach significance ($F_{(1,68)} = 3.147$, $r^2 = 0.044$, p = 0.081), while the quadratic model reached significance ($Y_i = 996.792 + 10.801^*X_i + -5.387^*X_i^2$ with $F_{(2,67)} = 41.708$, $r^2 = 0.555$, p < 0.001). The inverted U-shaped distribution of associated callosal areas across the principal gradient is depicted in Fig. 2F, right side, right.

4. Discussion

The current study investigated the relationship between the principal gradient and cortical projections of the corpus callosum. We reveal three main results: First, the apexes of the principal gradient span from the genu to the posterior body. Second, the individual variability of callosal connectivity is lowest at the apexes of the gradient. Lastly, the number of connected voxels in the callosal midsection showed a decrease at both ends of the gradient, while gradient values in the middle appeared more similar. We discuss these results in light of evolutionary theories, thus potentially shedding light onto evolutionary processes involved in changes of callosal connectivity.

Previous research investigated the topography of callosal connections solely based on atlases that parcellate the cortex into areas (Chao et al., 2009) or gyri (Park et al., 2008) that are not defined by their functional role nor their hierarchy in the brain. Our study circumvents this limitation by utilizing the principal gradient that has been linked to functional hierarchy (Margulies et al., 2016). To depict the principal gradient on the callosal midsection, we mapped each percentage of the gradient independently. The callosal gradient (Fig. 2E, left) indicates that the negative spectrum of the principal gradient peaks in the posterior body, whereas the positive spectrum peaks in the genu. Previous research demonstrated that the posterior midbody contains fibers that project to lower-level processing areas (e.g. somatosensory areas; Schmahmann et al., 2009). The genu connects to higher-level associative regions (i.e. prefrontal and orbitofrontal cortices; Schmahmann et al., 2009). Gradient values in the splenium were heterogeneous with highest values for a region previously shown as connecting the angular gyrus (Park et al., 2008; Chao et al., 2009) well known for its high values in the principal gradient (Margulies et al., 2016) and central structure of the default mode network (Zuo et al., 2012). Hence the spatial layout of the callosal gradient map is concordant with a previously hypothesised anatomical-functional organisation of the corpus callosum (Catani and Thiebaut de Schotten, 2012). Gradient values in the different portions of the Wittelson parcellation (Fig. 2E, right) indicated a heterogeneous distribution of the gradient along the corpus callosum. With some regions (3,2, 1) being highly consistent in terms of gradient values and region 7 to 4 being more distributed. A higher distribution of values might be related to a wider range of neuropsychological syndromes associated with disconnections of the same regions of the corpus callosum (Catani and Thiebaut de Schotten, 2012). Additionally, the callosal gradient resembles previous work mapping the diameter and myelination of callosal fibres (Aboitiz et al., 1992; Aboitiz and Montiel 2003). Accordingly, the genu contains poorly myelinated fibers of thin diameter compared to callosal fibers that connect to the posterior midbody. Diameter and myelination are contingent on conduction speed (Hursh, 1939; Waxman and Bennett, 1972). Furthermore, a study comparing conduction delay in various callosal segments showed a fastest speed in unimodal areas (motor, somatosensory and premotor areas) but a longer delay in transmodal temporal, parietal and occipital areas (Caminiti et al., 2013). This would suggest a different interhemispheric conduction speeds of unimodal (i.e. faster) and transmodal areas (i.e. slower). As such, the callosal gradient may putatively be related to different conduction speeds, which needs to be addressed by future research.

There is reason to believe that the general principle demonstrated at the group level might hold on the individual level, given that interindividual variability in callosal connectivity was lowest for the apexes of the gradient. These findings feed into a recent hypothesis in which structural variability is associated with evolutionary stability (Croxson et al., 2018). The observed callosal connectivity results may thus be linked to the phylogenetic age of cortical areas. Based on the dual origin theory, it has been suggested that multimodal, prefrontal and paralimbic cortices may have evolved in parallel to sensory and motor cortices (Pandy, Petrides, and Cipolloni, 2015). With regards to the gradient architecture, support for this assumption comes from non-human primate studies. For instance, a comparison between resting-state based gradients in human and structural connectivity-based gradients in macaque monkeys displayed similar spatial layouts spanning from primary sensory-motor to fronto-parietal regions (Oligschläger et al., 2019). In humans, this transmodal end of the gradient concords with the default mode network (DMN) that is also present in chimpanzees (Barks et al., 2015) and macaque monkeys (Mantini et al., 2011). Hence several pieces of evidence converge

towards phylogenetic older apexes of the gradient. While these arguments may support the interpretation that callosal connectivity may be a proxy for evolutionary stability, comparative studies may compare the callosal gradient architecture across primate species and derive phylogenetic models in order to test this assumption.

In addition, our results complement the latest model on the DMN anatomy (Alves et al., 2019), since the positive end of the principal gradient was highly connected with the contralateral hemisphere via the corpus callosum. This suggests that the core of the DMN (i.e. located at the positive end of the principal gradient, Waymel et al., 2020) could be phylogenetically more ancient than its periphery as areas highly connected by the corpus callosum might have appeared earlier on the evolutionary tree (Karolis et al., 2019). Clinical studies further hint at a link between the DMN and the corpus callosum, given that disrupted callosal white matter leads to decreases in DMN functional connectivity (Sharp et al., 2011). This link may appear unsurprising, since interhemispheric functional connectivity is lost shortly after complete transection of the corpus callosum (Johnston et al., 2008). In contrast, other studies suggest a preserved bilateral resting state networks (e.g. the DMN) in split-brain patients several years after surgery (Uddin et al., 2008; for a comprehensive overview see Mancuso et al., 2019). Together these findings point towards a contribution of the corpus callosum to the DMN, which, in split brain patients may recover with time.

With regards to functional laterality, a more symmetrical functional organization is found for higher probability in callosal connectivity (Karolis et al., 2019) and corresponds to the apexes of the gradient in the present study. This is in line with studies showing that stronger callosal connections are associated with a decrease in perceptual processing asymmetries. The corpus callosum has a long history as an underlying factor of functional hemispheric asymmetries (Ocklenburg et al., 2016) and appears to decrease asymmetries in perceptual processes (e.g. Friedrich et al., 2017). However, this result is not generalizable to all cognitive functions as for example studies investigating speech processes (i.e.: word fluency and semantic decision making) indicate that a 'stronger' corpus callosum increases asymmetries between the two hemispheres (Westerhausen et al., 2006; Josse et al., 2008). Hence, the callosal gradient map may be a useful new framework to clarify contrasting results in the role of the corpus callosum in functional hemispheric asymmetries.

A decrease in the variability of callosal connectivity at either end of the gradient was in part accompanied by a decrease of the number of voxels connected in the callosal midsagittal area on both ends of the gradient (Fig. 2F, right side). Inversely, callosal connectivity was more variable and involved more voxels in the corpus callosum when distant from the gradient's apexes. According to the tethering hypothesis, evolutionary expansion of the human cortex may have untethered large portions of the cortex from the constraints of early sensory activation cascades and molecular gradients (Buckner and Krienen, 2013). Similarly, the dual origin theory suggests that evolution progressively expended two cortical trends, one of which emerges from the hippocampocentric division (Pandya et al., 2015) and forms the limbic system that is cortically represented by the default mode network (Alves et al., 2019). In other words, evolutionary expansion would have created regions less functionally restricted than the apexes of the gradient. This, in turn, could have led to a more diverse pattern of connectivity (i.e. higher number of voxels in the associated callosal area) which is in accordance with the observed higher individual variability, the tethering hypothesis, as well as the dual origin theory.

Mapping the principal gradient onto the corpus callosum grants insight into the distribution of callosal connections across functions of various hierarchical levels. Contrary to categorical segmentation methods, our approach visualizes a transient gradient of areas in the callosal midsection based on the connectivity profile of unimodal and transmodal cortical areas. Therefore, this map may guide clinical investigation of brain lesions that disconnect the corpus callosum with a novel gradientbased perspective. It is important to stress that the current work is not a new parcellation of the CC. In the current work we aimed to avoid a categorical parcellation in favour of a map that shows smooth transitions of gradient with high and low values. As we work in the group average space, a continuum of values would be more accurate than a categorical parcellation (for a hard-border clustering-based solution, please see the supplementary material). Future research might explore the variability of division of the corpus callosum using gradients values and advanced tractography at the individual level. However, some limitations need to be noted. While we describe the first corpus callosum principal gradient map, this map depicts the weighted average of structural connectivity, based on an averaged map of the principal gradient. The weighted average of structural connectivity allowed us to assess - to some extent the variability of callosal connectivity, however, individual patterns of the principal gradient are yet unavailable. Therefore, the effect of individual variability of the principal gradient's spatial layout on callosal projections could not be assessed in this study. Addressing the extent to which the principal gradient and its callosal connections differ across individuals for example via new softwares (e.g. Vos de Wael et al., 2020) will be a matter of future research. Furthermore, revealing the individual variance of the gradient will allow us to investigate the relationship between individual variability and behaviour, similarly to other studies (e.g. Marquand et al., 2017).

Another limitation comes from the tractography approach. Tractography analyses can produce inaccurate results (Jones and Cercignani, 2010; Maier-Hein et al., 2017). To avoid these caveats, we employed methods that have previously demonstrated high anatomical reliability when compared to axonal tracing (Thiebaut de Schotten et al. 2011a; 2012) and with post-mortem dissections (Catani and Thiebaut de Schotten, 2012; Thiebaut de Schotten et al., 2011, 2017; Vergani et al., 2014). In the disconnectome tool, all image matrices were binarized depending on the existence of absence of streamlines connecting the corpus callosum with a given gradient percentage masque. In line with other studies (Gong et al., 2009; Shu et al., 2011), we chose a binarization threshold of 1 because the number of streamlines does not reflect the connectivity strength or the true number of axonal projections between two brain regions (Gong et al., 2009; Jones et al., 2013) and previous work shows that changing streamline count threshold for binarization does not change the overall results of network connectivity analyses (Shu et al., 2011). However, this approach does not prevent that each streamline may cross voxels of different gradient values, leading to potential overlap, especially if gradient values are partially asymmetric. The current approach does not allow to assess gradient asymmetries because of methodological limitations inherent to the gradient estimation (including for instance asymmetries of the seeding regions, usage of an asymmetrical template, contamination of the functional connectivity by the other hemisphere, and white matter asymmetries). Future studies may wish to address these limitations to investigate hemispheric asymmetries of the principal gradient along the callosal connectivity.

Conclusion

Our study provides the first comprehensive map of the callosal crosssection that displays the of callosal connections according to a hierarchical organisation of cognitive functions, without enforced boundaries. Our findings align with previous considerations about the evolutionary changes in the brain. Future studies may utilize the map of the callosal gradient to bridge the gap between brain asymmetries and functional hierarchy.

Data and code availability statement

The processed data including the callosal gradient map, the hardborder segmentation and the weighted and averaged tractogram can be found in https://neurovault.org/collections/7797/. The code for creating the hard-border segmentation of the corpus callosum is available in the following repository: https://github.com/PatFriedrich/ccGradient_ hardBorder. For further requests please contact the corresponding author.

Author credit statement

Dr. Patrick Friedrich designed the study, wrote the code for data processing and wrote and revised the manuscript.

Dr. Michel Thiebaut de Schotten, designed the study, processed the DTI data and wrote and revised the manuscript.

Dr. Stephanie Forkel, wrote and revised the manuscript.

All authors were enganged in discussions and interpretation of the results.

Acknowledgements

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant agreement No. 818521).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.neuroimage.2020.117317.

References

- Aboitiz, F., Montiel, J., 2003. One hundred million years of interhemispheric communication: the history of the corpus callosum. Braz. J. Med. Biol. Res. 36 (4), 409–420.
- Aboitiz, F., Scheibel, A.B., Fischer, R.S., Zaidel, E., 1992. Fiber composition of the human corpus-callosum. Brain Research 1–2, 143–153.
- Akelaitis, A.J., 1945. Studies on the corpus callosum: IV. Diagonistic dyspraxia in epileptics following partial and complete section of the corpus callosum. Am. J. Psychiatry 101 (5), 594–599.
- Alves, P.N., Foulon, C., Karolis, V., Bzdok, D., Margulies, D.S., Volle, E., Thiebaut de Schotten, M., 2019. An improved neuroanatomical model of the default-mode network reconciles previous neuroimaging and neuropathological findings. Commun. Biol. 2 (1), 1–14.
- Andersson, J.L.R., Xu, J., Yacoub, E., Auerbach, E., Moeller, S., Ugurbil, K., 2012, May. A comprehensive Gaussian process framework for correcting distortions and movements in diffusion images. In: Proceedings of the 20th Annual Meeting of ISMRM, 20, p. 2426.
- Andersson, J.L., Skare, S., Ashburner, J., 2003. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. Neuroimage 20 (2), 870–888.
- Barks, S.K., Parr, L.A., Rilling, J.K., 2015. The default mode network in chimpanzees (Pan troglodytes) is similar to that of humans. Cereb. Cortex 25 (2), 538–544.
- Biswal, B., Zerrin Yetkin, F., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magn. Reson. Med. 34 (4), 537–541.
- Bogen, J.E., 1993. The callosal syndromes. In: Heilman, K.M., Valenstein, E. (Eds.), Clinical Neuropsychology. Oxford University Press, pp. 337–407.
- Buckner, R.L., Krienen, F.M., 2013. The evolution of distributed association networks in the human brain. Trends Cogn. Sci. (Regul. Ed.) 17 (12), 648–665.
- Burdach, K.F.Royal College of Physicians of London, 1819. Vom Baue und Leben des Gehirns. Dy'sche Buchhandlung, Leipzig (Medical Heritage Library.).
- Caminiti, R., Carducci, F., Piervincenzi, C., Battaglia-Mayer, A., Confalone, G., Visco-Comandini, F., Innocenti, G.M., 2013. Diameter, length, speed, and conduction delay of callosal axons in macaque monkeys and humans: comparing data from histology and magnetic resonance imaging diffusion tractography. J. Neurosci. 33 (36), 14501–14511.
- Caruyer, E., Lenglet, C., Sapiro, G., Deriche, R., 2013. Design of multishell sampling schemes with uniform coverage in diffusion MRI. Magn. Reson. Med. 69 (6), 1534–1540.
- Catani, M., Robertsson, N., Beyh, A., Huynh, V., de Santiago Requejo, F., Howells, H., ... Krug, K., 2017. Short parietal lobe connections of the human and monkey brain. Cortex 97, 339–357.
- Catani, M., Thiebaut de Schotten, M., 2012. Atlas of Human Brain Connections. Oxford University Press.
- Chao, Y.P., Cho, K.H., Yeh, C.H., Chou, K.H., Chen, J.H., Lin, C.P., 2009. Probabilistic topography of human corpus callosum using cytoarchitectural parcellation and high angular resolution diffusion imaging tractography. Hum. Brain Mapp. 30 (10), 3172–3187.
- Coifman, R.R., Lafon, S., Lee, A.B., Maggioni, M., Nadler, B., Warner, F., Zucker, S.W., 2005. Geometric diffusions as a tool for harmonic analysis and structure definition of data: diffusion maps. In: Proceedings of the National Academy of Sciences, 102, pp. 7426–7431.
- Crosby, E.C., Humphrey, T., Lauer, E.W., 1962. White Matter of the Hemisphere, Pt 3. Correlative Anatomy of the Nervous System. Macmillan, New York.

- Croxson, P.L., Forkel, S.J., Cerliani, L., Thiebaut de Schotten, M., 2018. Structural variability across the primate brain: a cross-species comparison. Cereb. Cortex 28 (11), 3829–3841.
- Dejerine, J., 1892. Contribution à l'étude anatomopathologique et clinique des différents variétés de cécité verbale. Mém. Soc. Biol. 4, 61–90.
- Foulon, C., Cerliani, L., Kinkingnehun, S., Levy, R., Rosso, C., Urbanski, M., Thiebaut de Schotten, M., 2018. Advanced lesion symptom mapping analyses and implementation as BCBtoolkit. Gigascience 7 (3), giy004.
- Freund, C.S., 1889. II. Heber optische Aphasie und Seelenblindheit. Arch. Psychiatr. Nervenkr. 20 (1), 276–297.
- Friedrich, P., Ocklenburg, S., Heins, N., Schlüter, C., Fraenz, C., Beste, C., Genç, E., 2017. Callosal microstructure affects the timing of electrophysiological left-right differences. Neuroimage 163, 310–318.
- Geschwind, N., Fusillo, M., 1966. Color-naming defects in association with alexia. Arch. Neurol. 15 (2), 137–146.
- Gong, G., He, Y., Concha, L., Lebel, C., Gross, D.W., Evans, A.C., Beaulieu, C., 2009. Mapping anatomical connectivity patterns of human cerebral cortex using in vivo diffusion tensor imaging tractography. Cereb. Cortex 19 (3), 524–536.
- Heilman, K.M., Adams, D.J., 2003. Callosal neglect. Arch. Neurol. 60 (2), 276-279.
- Hofer, S., Frahm, J., 2006. Topography of the human corpus callosum revisited—Comprehensive fiber tractography using diffusion tensor magnetic resonance imaging. Neuroimage 32 (3), 989–994.
- Huang, H., Zhang, J., Jiang, H., Wakana, S., Poetscher, L., Miller, M.I., Mori, S., 2005. DTI tractography based parcellation of white matter: application to the mid-sagittal morphology of corpus callosum. Neuroimage 26 (1), 195–205.
- Huntenburg, J.M., Bazin, P.L., Margulies, D.S., 2018. Large-scale gradients in human cortical organization. Trends Cogn. Sci. (Regul. Ed.) 22 (1), 21–31.
- Hursh, J.B., 1939. Conduction velocity and diameter of nerve fibers. Am. J. Physiol. Legacy Content 127 (1), 131–139.
- Jones, D.K., Knösche, T.R., Turner, R., 2013. White matter integrity, fiber count, and other fallacies: the do's and don'ts of diffusion MRI. Neuroimage 73, 239–254.
- Jones, D.K., Cercignani, M., 2010. Twenty-five pitfalls in the analysis of diffusion MRI data. NMR Biomed. 23 (7), 803–820.
- Johnston, J.M., Vaishnavi, S.N., Smyth, M.D., Zhang, D., He, B.J., Zempel, J.M., ... Raichle, M.E., 2008. Loss of resting interhemispheric functional connectivity after complete section of the corpus callosum. J. Neurosci. 28 (25), 6453–6458.
- Josse, G., Seghier, M.L., Kherif, F., Price, C.J., 2008. Explaining function with anatomy: language lateralization and corpus callosum size. J. Neurosci. 28 (52), 14132–14139.
- Karolis, V.R., Corbetta, M., De Schotten, M.T., 2019. The architecture of functional lateralisation and its relationship to callosal connectivity in the human brain. Nat. Commun. 10 (1), 1–9.
- Kashiwagi, A., Kashiwagi, T., Nishikawa, T., Tanabe, H., Okuda, J, 1990. Hemispatial neglect in a patient with callosal infarction. Brain 113 (4), 1005–1023.
- Lausberg, H., Kita, S., Zaidel, E., Ptito, A., 2003. Split-brain patients neglect left personal space during right-handed gestures. Neuropsychologia 41 (10), 1317–1329.
- Luzzatti, C., Vecchi, T., Agazzi, D., Cesa-Bianchi, M., Vergani, C., 1998. A neurological dissociation between preserved visual and impaired spatial processing in mental imagery. Cortex 34 (3), 461–469.
- Maier-Hein, K.H., Neher, P.F., Houde, J.C., Côté, M.A., Garyfallidis, E., Zhong, J., Reddick, W.E., 2017. The challenge of mapping the human connectome based on diffusion tractography. Nat. Commun. 8 (1), 1–13.
- Mantini, D., Gerits, A., Nelissen, K., Durand, J.B., Joly, O., Simone, L., Vanduffel, W., 2011. Default mode of brain function in monkeys. J. Neurosci. 31 (36), 12954–12962.
- Mancuso, L., Uddin, L.Q., Nani, A., Costa, T., Cauda, F., 2019. Brain functional connectivity in individuals with callosotomy and agenesis of the corpus callosum: a systematic review. Neurosci. Biobehav. Rev. 105, 231–248.
- Margulies, D.S., Ghosh, S.S., Goulas, A., Falkiewicz, M., Huntenburg, J.M., Langs, G., Jefferies, E., 2016. Situating the default-mode network along a principal gradient of macroscale cortical organization. Proceed. Natl. Acad. Sci. 113 (44), 12574–12579.
- Marquand, A.F., Haak, K.V., Beckmann, C.F., 2017. Functional corticostriatal connection topographies predict goal-directed behaviour in humans. Nat. Hum. Behav. 1 (8), 1–9. McInnes, L., Healy, J., Astels, S., 2017. hdbscan: Hierarchical density based clustering. J.
- Open Source Softw. 2 (11), 205. McInnes, L., Healy, J., & Melville, J. (2018). Umap: uniform manifold approximation and
- projection for dimension reduction. arXiv preprint arXiv:1802.03426.

Mesulam, M.M., 1998. From sensation to cognition. Brain: J. Neurol. 121 (6), 1013–1052. Moeller, S., Yacoub, E., Olman, C.A., Auerbach, E., Strupp, J., Harel, N., Uğurbil, K., 2010.

- Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. Magn. Reson. Med. 63 (5), 1144–1153.
- Ocklenburg, S., Friedrich, P., Güntürkün, O., Genç, E., 2016. Intrahemispheric white matter asymmetries: the missing link between brain structure and functional lateralization? Rev. Neurosci. 27 (5), 465–480.
- Oligschläger, S., Xu, T., Baczkowski, B.M., Falkiewicz, M., Falchier, A., Linn, G., Margulies, D.S., 2019. Gradients of connectivity distance in the cerebral cortex of the macaque monkey. Brain Struct. Funct. 224 (2), 925–935.
- O'Reilly, J.X., Croxson, P.L., Jbabdi, S., Sallet, J., Noonan, M.P., Mars, R.B., Rushworth, M.F., 2013. Causal effect of disconnection lesions on interhemispheric functional connectivity in rhesus monkeys. Proceed. Natl. Acad. Sci. 110 (34), 13982–13987.
- Pandya, D., Petrides, M., Cipolloni, P.B., 2015. Cerebral Cortex: Architecture, Connections, and the Dual Origin Concept. Oxford University Press.
- Park, H.J., Kim, J.J., Lee, S.K., Seok, J.H., Chun, J., Kim, D.I., Lee, J.D., 2008. Corpus callosal connection mapping using cortical gray matter parcellation and DT-MRI. Hum. Brain Mapp. 29 (5), 503–516.

- Park, K.C., Lee, B.H., Kim, E.J., Shin, M.H., Choi, K.M., Yoon, S.S., Kwon, S.U., ChungCS, Lee KH, Heilman, K.M., et al., 2006. Deafferentation-disconnection neglect induced by posterior cerebral artery infarction. Neurology. 66 (1), 56–61.
- Power, J.D., Cohen, A.L., Nelson, S.M., Wig, G.S., Barnes, K.A., Church, J.A., Petersen, S.E., 2011. Functional network organization of the human brain. Neuron 72 (4), 665–678.
- Risse, G.L., LeDoux, J., Springer, S.P., Wilson, D.H., Gazzaniga, M.S., 1978. The anterior commissure in man: functional variation in a multisensory system. Neuropsychologia 16 (1), 23–31.
- Roland, J.L., Snyder, A.Z., Hacker, C.D., Mitra, A., Shimony, J.S., Limbrick, D.D., Leuthardt, E.C., 2017. On the role of the corpus callosum in interhemispheric functional connectivity in humans. Proceed. Natl Acad. Sci. 114 (50), 13278–13283.
- Sharp, D.J., Beckmann, C.F., Greenwood, R., Kinnunen, K.M., Bonnelle, V., De Boissezon, X., Leech, R., 2011. Default mode network functional and structural connectivity after traumatic brain injury. Brain 134 (8), 2233–2247.
- Schmahmann, J.D., Schmahmann, J., Pandya, D., 2009. Fiber Pathways of the Brain. OUP, USA.
- Shu, N., Liu, Y., Li, K., Duan, Y., Wang, J., Yu, C., ... He, Y., 2011. Diffusion tensor tractography reveals disrupted topological efficiency in white matter structural networks in multiple sclerosis. Cereb. cortex 21 (11), 2565–2577.
- Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., ... Beckmann, C.F., 2009. Correspondence of the brain's functional architecture during activation and rest. Proceed. Natl. Acad. Sci. 106 (31), 13040–13045.
- Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E., Johansen-Berg, H., ... Niazy, R.K., 2004. Advances in functional and structural MR image analysis and implementation as FSL. Neuroimage 23, S208–S219.
- Sotiropoulos, S.N., Jbabdi, S., Xu, J., Andersson, J.L., Moeller, S., Auerbach, E.J., ... Feinberg, D.A., 2013. Advances in diffusion MRI acquisition and processing in the human Connectome project. Neuroimage 80, 125–143.
- Tavor, I., Jones, O.P., Mars, R.B., Smith, S.M., Behrens, T.E., Jbabdi, S., 2016. Task-free MRI predicts individual differences in brain activity during task performance. Science 352 (6282), 216–220.
- Thiebaut de Schotten, M., Bizzi, A., Dell'Acqua, F., Allin, M., Walshe, M., Murray, R., ... Catani, M., 2011. Atlasing location, asymmetry and inter-subject variability of white matter tracts in the human brain with MR diffusion tractography. Neuroimage 54 (1), 49–59.
- Thiebaut De Schotten, M., M., Dell'Acqua, Dell'Acqua, M., Forkel, S.J., Simmons, A., Vergani, F., Murphy, D.G.M, Catani, M., 2011. A lateralized brain network for visuo-spatial attention. Nature Neuroscience 14 (10), 1245–1246.

- Thiebaut de Schotten, M., Dell'Acqua, F., Valabregue, R., Catani, M., 2012. Monkey to human comparative anatomy of the frontal lobe association tracts. Cortex 48 (1), 82–96
- Tomaiuolo, F., Voci, L., Bresci, M., Cozza, S., Posteraro, F., Oliva, M., Doricchi, F., 2010. Selective visual neglect in right brain damaged patients with splenial interhemispheric disconnection. Exp. Brain Res. 206 (2), 209–217.
- Tomasch, J., 1954. Size, distribution, and number of fibres in the human corpus callosum. Anat. Rec. 119 (1), 119–135.
- Uddin, L.Q., Mooshagian, E., Zaidel, E., Scheres, A., Margulies, D.S., Kelly, A.C., ... Milham, M.P., 2008. Residual functional connectivity in the split-brain revealed with resting-state fMRI. Neuroreport 19 (7), 703.
- Van Essen, D.C., Smith, S.M., Barch, D.M., Behrens, T.E., Yacoub, E., Ugurbil, K., Consortium, Wu-Minn HCP, 2013. The WU-Minn human connectome project: an overview. Neuroimage 80, 62–79.
- Vergani, F., Lacerda, L., Martino, J., Attems, J., Morris, C., Mitchell, P., Dell'Acqua, F., 2014. White matter connections of the supplementary motor area in humans. J. Neurol. Neurosurg. Psychiatry 85 (12), 1377–1385.
- Vos de Wael, R., Benkarim, O., Paquola, C., Lariviere, S., Royer, J., Tavakol, S., ... Misic, B., 2020. BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. Commun. Biol. 3 (1), 1–10.
- Vu, A.T., Auerbach, E., Lenglet, C., Moeller, S., Sotiropoulos, S.N., Jbabdi, S., ... Ugurbil, K., 2015. High resolution whole brain diffusion imaging at 7 T for the Human Connectome Project. Neuroimage 122, 318–331.
- Waxman, S.G., Bennett, M.V., 1972. Relative conduction velocities of small myelinated and non-myelinated fibres in the central nervous system. Nat. New Biol. 238 (85), 217–219.
- Waymel, A., Friedrich, P., Bastian, P.A., Forkel, S.J., De Schotten, M.T., 2020. Anchoring the human olfactory system to a functional gradient. Neuroimage, 116863.
- Westerhausen, R., Kreuder, F., Sequeira, S.D.S., Walter, C., Woerner, W., Wittling, R.A., ... Wittling, W., 2006. The association of macro-and microstructure of the corpus callosum and language lateralisation. Brain Lang. 97 (1), 80–90.
- Witelson, S.F., 1989. Hand and sex differences in the isthmus and genu of the human corpus callosum: a postmortem morphological study. Brain 112 (3), 799–835.
- Zuo, X.N., Ehmke, R., Mennes, M., Imperati, D., Castellanos, F.X., Sporns, O., Milham, M.P., 2012. Network centrality in the human functional connectome. Cereb. Cortex 22 (8), 1862–1875.