










Estimation of Forest Water Potential From Ground-Based L-Band Radiometry

Thomas Jagdhuber , Senior Member, IEEE, Anne-Sophie Schmidt, Anke Fluhrer , Member, IEEE, David Chaparro , Member, IEEE, François Jonard , Senior Member, IEEE, María Piles , Senior Member, IEEE, Natan Holtzman, Alexandra G. Konings , Andrew F. Feldman, Martin J. Baur, Susan Steele-Dunne , Member, IEEE, Konstantin Schellenberg , and Harald Kunstmann 

Abstract—Monitoring the water status of forests is paramount for assessing vegetation health, particularly in the context of increasing duration and intensity of droughts. In this study, a

Received 22 May 2024; revised 29 September 2024 and 20 December 2024; accepted 14 January 2025. Date of publication 23 January 2025; date of current version 18 February 2025. This work was supported in part by the MIT-Germany Seed Fund “Global Water Cycle and Environmental Monitoring using Active and Passive Satellite-based Microwave Instruments,” and in part by the MIT-Belgium Seed Fund “Early Detection of Plant Water Stress Using Remote Sensing.” The work of David Chaparro was supported in part by the Ramón Areces Foundation postdoctoral grant, in part by the projects “la Caixa” Junior Leader Fellowship under Grant LCF/BQ/PI23/11970013 (lead: O. Binks) and Grant H2020 FORGENIUS (Improving access to FORest GENetic resources Information and services for end-USers) #862221, and in part by the Ramón Areces Postdoctoral Fellowship. The work of Andrew F. Feldman was supported in part by the NASA ECOSTRESS science team and in part by the NASA Terrestrial Ecology scoping study for a dryland field campaign (ARID). (Corresponding author: Thomas Jagdhuber.)

Thomas Jagdhuber and Anke Fluhrer are with the Microwaves and Radar Institute, German Aerospace Center, 82234 Wessling, Germany, and also with the Institute of Geography, Augsburg University, 86159 Augsburg, Germany (e-mail: Thomas.jagdhuber@dlr.de; anke.fluhrer@dlr.de).

Anne-Sophie Schmidt is with the Microwaves and Radar Institute, German Aerospace Center, 82234 Wessling, Germany (e-mail: anne-sophie.schmidt@dlr.de).

David Chaparro is with the Microwaves and Radar Institute, German Aerospace Center, 82234 Wessling, Germany, and also with the Center for Ecological Research and Forestry Applications, 08197 Cerdanyola del Vallès, Spain (e-mail: d.chaparro@creaf.uab.cat).

François Jonard is with the Earth Observation and Ecosystem Modelling Laboratory, SPHERES Research Unit, Université de Liège (ULiège), 4000 Liège, Belgium (e-mail: francois.jonard@uliege.be).

María Piles is with the Image Processing Lab, University of Valencia, 46980 Paterna, Spain (e-mail: maria.piles@uv.es).

Natan Holtzman and Alexandra G. Konings are with the Stanford University, School of Earth & Environmental Sciences, Stanford, CA 94305-2210 USA (e-mail: nholtzma@stanford.edu; konings@stanford.edu).

Andrew F. Feldman is with the NASA Goddard Spaceflight Center, Biospheric Sciences Laboratory, Greenbelt, MD 20771 USA, and also with the Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20742 USA (e-mail: andrew.feldman@nasa.gov).

Martin J. Baur is with the Department of Geography, University of Cambridge, CB2 3EN Cambridge, U.K. (e-mail: mjb311@cam.ac.uk).

Susan Steele-Dunne is with the Department of Geoscience and Remote Sensing, Delft University of Technology, 2628 CD Delft, The Netherlands (e-mail: s.c.steele-dunne@tudelft.nl).

Konstantin Schellenberg is with the Microwaves and Radar Institute, German Aerospace Center, 82234 Wessling, Germany, also with the Department for Earth Observation, Friedrich Schiller University Jena, 07743 Jena, Germany, and also with the Department Biogeochemical Processes, Max Planck Institute for Biogeochemistry, 07745 Jena, Germany (e-mail: konstantin.schellenberg@uni-jena.de).

Harald Kunstmann is with the Institute of Geography, Augsburg University, 86159 Augsburg, Germany (e-mail: harald.kunstmann@geo.uni-augsburg.de).

This article has supplementary downloadable material available at <https://doi.org/10.1109/JSTARS.2025.3533567>, provided by the authors.

Digital Object Identifier 10.1109/JSTARS.2025.3533567

methodology was developed for estimating forest water potential at the canopy scale from ground-based L-band radiometry. The study uses radiometer data from a tower-based experiment of the SMAPVEX 19-21 campaign from April to October 2019 at Harvard Forest, MA, USA. The gravimetric and the relative water content of the forest stand was retrieved from radiometer-based vegetation optical depth. A model-based methodology was adapted and assessed to transform the relative water content estimates into values of forest water potential. A comparison and validation of the retrieved forest water potential was conducted with in situ measurements of leaf and xylem water potential to understand the limitations and potentials of the proposed approach for diurnal, weekly and monthly time scales. The radiometer-based water potential estimates of the forest stand were found to be consistent in time with r_{Pearson} correlations up to 0.6 and similar in value, down to RMSE = 0.14 [MPa], compared to their in situ measurements from individual trees in the radiometer footprint, showing encouraging retrieval capabilities. However, a major challenge was the bias between the radiometer-based estimates and the in situ measurements over longer times (weeks & months). Here, an approach using either air temperature or soil moisture to update the minimum water potential of the forest stand (FWP_{min}) was developed to adjust the mismatch. These results showcase the potential of microwave radiometry for continuous monitoring of plant water status at different spatial and temporal scales, which has long been awaited by forest ecologists and tree physiologists.

Index Terms—L-band, microwave radiometry, soil moisture active-passive (SMAP), soil plant atmosphere system, SPAC, transmissivity, vegetation, vegetation water potential.

I. INTRODUCTION

IN PLANTS, water moves due to suction (and tension) created by a potential energy gradient from the soil into the atmosphere and by capillary action within plant vessels [4], [15], [26] and [27], [71]). Thus, water is driven by a pressure gradient which allows plants to take up water from the soils, transfer it through the xylem to the leaves, and finally release it into the atmosphere by transpiration through stomata.

Plant water potential is often described for individual plant components (e.g., leaf water potential) or referred to by the physical process to be analyzed, like matric and gravimetric water potential [4]. Since this study focuses on vegetation at the canopy scale, it will refer generically to forest water potential (FWP). The FWP depends on the soil water availability: a drier soil has a lower soil water potential, which can limit the plant water uptake and reduce the FWP [77]. Finally, at the leaf canopy–atmosphere interface, the dryness of the atmosphere (i.e., vapor pressure deficit) contributes to the pressure gradient

in the plant. A drier atmosphere can induce higher transpiration and decreases the FWP which, in turn, may induce stomatal closure in order to reduce water losses at the cost of reducing plant carbon uptake [35], [47]. This way, FWP can control transpiration and photosynthesis rates [78] by changing stomatal conductance. FWP is important for plant hydraulic behavior and is a key variable influencing the water, energy, and carbon cycles [42], [47]. Including FWP information as a predictor can lead to improved estimates of stomatal conductance responses to drought [2], [6], enhance the capacity to understand transpiration responses to increased vapor pressure deficit under climate change [45], and allow enhanced land surface models by accounting for plant hydraulics [12], [21], [37], [49]. Detecting reductions in water potential is also essential to assess the risk of tree mortality [10], [50], [57], [80] and of fire [56], as well as to explain variations in tree growth and in forest phenology [81].

FWP monitoring helps to predict an ecosystems' response to actual and future climate extremes. However, current FWP measurements are mainly limited to in situ efforts, meaning FWP is sparsely measured in space and time, given destructive and expensive techniques [59]. Specifically, manual plant water potential observations are mostly taken with pressure chambers [64], [70] and psychrometers [19], among other techniques [3], [17], [51].

Due to these limitations, only a few a databases of plant water potential measurements exist (e.g., [9]). Still, they are site-specific and are not continuously sampled in time. Data on plant water potential are rarely collected (PsiNet) and not standardized in databases and networks. Moreover, scaling water potential measurements from the individual plant to entire forests or ecosystems remains a challenge ([41]; Novick et al., 2022). Satellite sensors have the potential to measure vegetation conditions and fulfill the scientific need for global water potential estimates with a high temporal resolution (e.g., less than a week). So far, optical vegetation indices (VIs) from satellite and airborne platforms have shown promising results even with commercial applicability in croplands [28], [44], [85] and in some natural landscapes ([60] and [32], [61]), at local scales. To the authors' knowledge, however, regional-scale or larger scale studies are lacking so far. In addition, optical VIs are limited by clouds and sun-glint effects and, importantly, are only sensitive to the vegetation properties at the top of the canopy leaves [40], [84]. Moreover, VIs that measure pigment (chlorophyll) variations and therefore photosynthetic activity are even less sensitive to vegetation water content dynamics directly.

To overcome these issues, the global ecological remote sensing community has recently explored the capacity of passive microwave measurements to sense the water potential of vegetation. Such measurements have the advantage of being sensitive to plants' water content, structure, and biomass. In passive microwave radiometry, this information is integrated in a parameter named vegetation optical depth (VOD). VOD refers to the attenuation that the vegetation canopy exerts over the land emissions in the microwave range of the electromagnetic spectrum [52], [76]. At higher microwave frequencies (i.e., X-band, ~ 10 GHz, and Ku-band, ~ 15 GHz), the VOD is sensitive to the water content in the upper canopy (due to lack of canopy sensing depth

at these frequencies). It has been related to the live fuel moisture content (Forkel et al., 2023; [8]) or to the isohydricity behavior of vegetation [39] at global scale. At lower frequencies (i.e., L-band, ~ 1 GHz, which senses nearly all the canopy layers given its longer wavelength), a recent forest study has revealed that VOD is correlated with the xylem water potential, and that it shows a similar diurnal cycle to in situ xylem and leaf water potential [30].

Ideally, the water component of VOD could be disentangled from the components of biomass and structure to evaluate the water status of plants more precisely. To achieve this goal, a common strategy is to derive the relative water content (RWC) by normalizing VOD in time, and assuming the biomass is constant at yearly time scales [48], [62].

Another strategy consists of estimating vegetation moisture mg (in [kg/kg]) from VOD using information from additional sensors and deriving RWC from minimum and maximum values of mg. This approach has been applied at local [53] and continental scales (Fink et al., 2018, [8]) using active microwaves and lidar to account for the biomass and vegetation structure imprint on VOD. From each of the approaches, the obtained RWC can be used to estimate FWP by applying pressure-volume curves [34], [40], [41].

The goal of this study is therefore to derive FWP dynamics using tower-based, local-scale L-band VOD measurements coupled to pressure-volume curves. These time series of FWP are compared with in situ xylem and leaf water potential measurements. Thus, it is hypothesized that the proposed approach leads to a first ground-based FWP retrieval from VOD. Moreover, the diurnal, weekly and up to monthly dynamics have been assessed to understand the ability to estimate water potential on different time scales. The study sets the path for enabling future retrievals that could be based on passive microwave sensors currently in orbit, like the Advanced Microwave Scanning Radiometer 2 (AMSR-2), and the L-band radiometers on the soil moisture and ocean salinity and the soil moisture active-passive (SMAP) missions. Such FWP retrievals, if provided at larger spatial scales (e.g., regional), would allow to study forest water relations at a scale that was not possible before, and would ease finding forest drought-induced mortality thresholds in space and time to know when and where forests are at risk of die back.

II. TEST SITE AND EXPERIMENTAL DATA

Previous to the study, a ground-based acquisition setup was designed to estimate FWP and established within the research site of Harvard University [30]. The forest test site is located on Prospect Hill within Harvard Forest (42.535° N, 72.174° W), MA, USA. It is a temperate deciduous forest dominated by red Oak (*Quercus rubra*) in a humid continental climate [30]. An overview of the measurement setup is shown in Fig. 1.

As part of the SMAP Validation Experiment 2019-2021 (SMAPVEX 19 -21), in situ measurements for soil, plants, and atmosphere were carried out between 28th of April and 17th of October 2019 [14]. These measurements include soil moisture and soil temperature at 5 and 10 cm depth, tree xylem permittivity and xylem water potential (XWP), leaf wetness and leaf water potential (LWP), and air temperature. Not all




Fig. 1. Overview of the measurement setup at the Harvard Forest, MA test site during the SMAPVEX 19-21 campaign: The tower-based radiometer with its slanted observation geometry sensing into the forest complemented by in situ measurements on air and soil temperature, soil moisture, xylem and leaf water potential, leaf permittivity and leaf wetness [13], [30].

in situ measurements could be conducted continuously along the campaign period. Three intensive measurement periods took place between 9th and 18th of July, 5th and 13th of August, and 7th and 26th of September 2019 with almost all sensors in operation.

An L-band (1.4 GHz) radiometer (Potter Horn, PR-1475, Radiometrics Inc.) was mounted at 28 m height on a tower pointing with an incidence angle of 40° into the forest canopy with about 21 to 23 m height (cf. Fig. 1). This incidence angle is consistent with the current spaceborne SMAP radiometer. The ground footprint dimensions are 25 by 20 m, defined by the half power beam width (-3 dB) of the antenna at 30° opening angle [29]. The instrument conducted hourly measurements of dual-polarimetric, i.e., horizontally (H) and vertically (V) polarized brightness temperatures with an accuracy of 2 K [65]. Only V-polarized measurements were used in this study, as H-polarized data showed unexplained fluctuations, as stated previously in Holtzman et al. [30]. Further details on radiometer operation and in situ measurements are provided in Holtzman et al. [30] and Roy et al. [66].

The VOD-estimates, also included in Holtzman et al. [29], were calculated using a single channel V-polarized (SCA-V) retrieval algorithm based on a zeroth order radiative transfer model [23], [55]. In contrast to the classical SCA-V retrieval approach in the SMAP baseline algorithm [23], in situ-based soil moisture was here input to the algorithm, and the radiative transfer equation solved for VOD, like in Baur et al. [5]. Radiometer measurements were made at times lacking thermal equilibrium between the soil and vegetation (such as appropriately assumed at 6 am for the SMAP baseline algorithm [23]). Therefore, the soil and canopy temperatures were taken from the in situ measurements of Holtzman et al. [29].




Fig. 2. Time series of L-band radiometer-retrieved vegetation optical depth (VOD) [-] and in situ-measured xylem water potential (XWP) [MPa] in the temperate forest stand during the intensive measurement campaign in July 2019 at prospect hill (harvard forest).

For canopy temperature, the air temperature at one-meter height within the forest stand was used, because direct canopy measurements were not available. This temperature substitution was investigated by Holtzman et al. [30] and was shown to be sufficient. For parameterization of soil roughness and scattering albedo, the standard values of Holtzman et al. [30], deduced from the SMAP soil moisture product for temperate broadleaf forests, were taken (soil roughness: $h = 0.16$, scattering albedo: $\omega = 0.05$).

Fig. 2 presents the VOD time series from 10th (noon) to 18th (11 pm) of July 2019 during the first intensive measuring period together with in situ data of the XWP [MPa], measured for one tree in the stand. The VOD time series in Fig. 2 shows a diurnal dynamic of temperature (driven by the solar cycle, cf.




Fig. 3. Time series of *in situ*-measured soil and air temperature [K] and net radiation [W/m²] (top) as well as surface soil moisture [m³/m³], precipitation [mm/h] and depth to ground water table [m] (bottom) in the temperate forest stand during the intensive measurement campaign in July 2019 at Prospect Hill (Harvard Forest).

Fig. 3 top) similar to the XWP dynamics in most days (with few exceptions, e.g., July 17th). An overall decreasing trend was observed similar to the hydro-dynamics of the *in situ* soil moisture (cf. Fig. 3 bottom) along multiple days up to one week. Fig. 3 provides an *in situ*-based overview on the water and energy status and dynamics for the intensive measurement campaign in July 2019 (August and September campaigns in the Supplement). It reveals how the light and water conditions influenced the *in situ* XWP measurements and remote sensing (VOD) plant water dynamics during the intensive campaign period. Diurnal soil and air temperature minima and maxima are opposite to that of the xylem water potential confirming the interplay between light and water.

III. METHODOLOGY OF RETRIEVING FOREST WATER POTENTIAL FROM VEGETATION OPTICAL DEPTH

Since VOD represents the attenuation of the L-band electromagnetic waves by the forest canopy, it is not only sensitive to the water component of the canopy, but also to its biomass and structure characteristics. Here, a simple, physics-based and analytically solvable model of VOD from Jackson and Schmugge [33] is used to disentangle the water component from the radiometer-based VOD.

A. From VOD to Gravimetric Water Content of Vegetation m_g

VOD is linked to the gravimetric water content of vegetation m_g , in kilogram of water per kilogram of wet biomass [kg/kg]. The m_g can be extracted by the comparison of forward modeled VOD with data-retrieved VOD as initially applied in work [53] and further refined in work [8].

The VOD model in (1)–(3) needs information on the vegetation height h_{veg} [m], the vegetation volume fraction δ_i [m³/m³] (volume of plant material per volume of air as seen by microwave sensors), the main plant structure (here: random orientation of plant components), the shape of the major plant component (here: spheroids) and the wavelength of the observing system [m] (here: L-band, $\lambda = 0.21$ [m]) as input variables [8], [33], [53], [69], [75]

$$\text{VOD} = 4 \cdot \pi \cdot \frac{h_{veg}}{\lambda} \cdot \text{Im}[\sqrt{\epsilon_{can}}]. \quad (1)$$

ϵ_{can} is the dielectric constant of the canopy

$$\epsilon_{can} = \epsilon_h + \delta_i \frac{(\epsilon_i - \epsilon_h)(\epsilon^* + \epsilon_i)}{3(\epsilon_i + \epsilon^*)} \quad (2)$$

where ϵ_h [-] and ϵ_i [-] are the permittivity of the host medium and the inclusions, respectively. The host medium is assumed to be air ($\epsilon_h = 1$). In addition, it is assumed that $\epsilon^* = \epsilon_h$, as δ_i is expected to be small ($\delta_i \ll 1$) [75].

Equation (2) is derived from Ulaby and Long [76] assuming that the spheroids accounting for the plant material structure are randomly oriented needles, in agreement with Meyer et al. [53] and Chaparro et al. [7].

The permittivity of the plant material can be linked to m_g by the Debye–Cole dual dispersion dielectric mixing model [75]

$$\epsilon_i = \epsilon_{fw} v_{fw} + \epsilon_{bw} v_{bw} + \epsilon_r \quad (3)$$

where ϵ_{fw} and ϵ_{bw} are the permittivity of free water and bound water within the plants, respectively. The terms v_{fw} , v_{bw} , and ϵ_r are the volume fractions of free and bound water as well as the residual permittivity component. The latter three parameters are a function of m_g , the salinity and the temperature of the plant material (cf. [75] for details). The salinity and the temperature are fixed to 10 [‰] and 22 [°C], as done in previous studies [8], [75]. Details of this dielectric mixing model are described in Ulaby and El-Reyes [75]. The m_g -value minimizing the difference between the modelled VOD in (1)–(3) and the radiometer-derived VOD is chosen as the result.

B. From Gravimetric Water Content of Vegetation m_g to Relative Water Content RWC

Next, m_g is converted into the relative water content (RWC) using dry moisture-minimum ($m_{g_{min}}$) and full turgor moisture-maximum ($m_{g_{max}}$) references [72]:

$$\text{RWC}_{m_g} = \frac{m_g - m_{g_{min}}}{m_{g_{max}} - m_{g_{min}}} \cdot 100 [\%]. \quad (4)$$

The definition of the extremes (min, max) sets the boundaries of the relative water dynamic and therefore the extent of the dynamic when moving to a relative metric. The time period for finding the extreme values may reach from diurnal (24 h)




Fig. 4. Conceptual view on the RWC-FWP relationship for forest under slow, intermediate and rapid change rate, adapted from Zweifel et al. [91]; calculated with (6) using $\text{FWP}_{\min} = 2.5$ [MPa], $k_{1\text{slow}} = 55$ & $k_{2\text{slow}} = 10$, $k_{1\text{intermediate}} = 68$ & $k_{2\text{intermediate}} = 7.5$, $k_{1\text{rapid}} = 81$ & $k_{2\text{rapid}} = 5$.

up to seasonal (e.g., summer, winter) or even up to multiannual scale, but will be set to diurnal ranges for the following analyses. Moreover, VOD can be directly used to calculate RWC in case of stable biomass and canopy structure conditions along the observation period [62]

$$\text{RWC}_{\text{VOD}} = \frac{\text{VOD} - \text{VOD}_{\min}}{\text{VOD}_{\max} - \text{VOD}_{\min}} \cdot 100 \text{ [%]}. \quad (5)$$

RWC_{VOD} will serve as a benchmark to $\text{RWC}_{m,g}$ and as a relative indicator of biomass/structure-bias between (4) and (5).

C. From Relative Water Content RWC to Forest Water Potential FWP

RWC is then converted into FWP by the sigmoidal function of Zweifel et al. [91]

$$\text{FWP}_{m,g} = \frac{\text{FWP}_{\min}}{e^{-\frac{k_1 + \text{RWC}}{k_2}} + 1} \text{ [MPa]} \quad (6)$$

where FWP_{\min} is the minimum forest water potential of the RWC-FWP relationship. Parameters k_1 [-] and k_2 [-] are empirical (site- and plant-specific) representing the inflection point and the rate of change between RWC and FWP. These parameters depend on hydraulic components of the vegetation and local climate within the stand.

The advantage of this functional form compared to alternative ones (e.g., Mirfenderesgi et al. [54]) is it is easy to adapt and has parsimonious character (k_1 , k_2 , FWP_{\min}), to enable water potential estimation. Fig. 4 illustrates the adaptivity of this relationship for a slow ($k_1 = 55$, $k_2 = 10$), an intermediate ($k_1 = 68$, $k_2 = 7.5$), and a rapid ($k_1 = 81$, $k_2 = 5$) rate of RWC-to-FWP change, starting from an FWP_{\min} of -2.5 [MPa].

In this work, the three parameters (k_1 , k_2 , FWP_{\min}) were optimized jointly using the on-site XWP (psychrometer) measurements of the first intensive observation period in July 2019. The optimization results in k_1 and k_2 of 25 and 31, respectively.

Both were fixed throughout the analyses of the remaining time series. This was supported by a sensitivity analysis (not shown) indicating that the most sensitive parameter in the RWC-FWP relationship is FWP_{\min} compared to k_1 and k_2 . FWP_{\min} was first set to the minimum of XWP of the first diurnal cycle and then updated dynamically through time.

D. Dynamization of FWP by Updating FWP_{\min}

The dynamic update of FWP_{\min} was based on two commonly available and continuously recorded in situ measurements: surface soil moisture θ_s [m^3/m^3] and air temperature T [K] at 1 [m] height, as they are key factors in constraining plant water uptake and plant water loss, respectively. In more general terms, these dynamic updates of FWP with soil moisture as well as with air temperature enable the extrapolation of the FWP estimation to longer periods, in terms of weeks and months toward seasons.

For FWP_{\min} -update, θ_s and T were tested individually to understand the benefit of a mass-related variable (θ_s) and an energy-related variable (T). In this study, this FWP_{\min} -update was developed by adapting the numerator of (6). Equations (7) and (8) define the proposed FWP_{\min} -updated equations.

The soil moisture-based FWP_{\min} -update is expressed by the following:

$$\text{FWP} = \frac{\text{FWP}_{\min} + |\text{FWP}_{\min}| * \frac{\partial \theta_s}{\partial t}}{e^{-\frac{k_1 + \text{RWC}}{k_2}} + 1} \text{ [MPa]} \quad (7)$$

where $\frac{\partial \theta_s}{\partial t}$ represents the rate of change in soil moisture, $\partial \theta_s$, along time interval, ∂t , that is multiplied with the increment of $|\text{FWP}_{\min}|$ from the previous time step. Soil moisture-based FWP_{\min} -update is done in steps of five hours using the preceding five-hour time period for calculus. Shortly after a significant rain pulse, $\frac{\partial \theta_s}{\partial t}$ can grow as large as 0.5 [$\frac{\text{m}^3}{\text{m}^3}/\text{h}$] or higher. In these exceptional cases of 50% increase within the preceding time interval, a data-calibrated cap factor of 0.7 was used for the respective rate of change to control updates of FWP_{\min} and prevent overshooting.

Corresponding to (7), (6) can be also adapted for an air temperature-based FWP_{\min} -update

$$\text{FWP} = \frac{\text{FWP}_{\min} + |\text{FWP}_{\min}| * \left(-\frac{\partial T}{\partial t} * k_3\right)}{e^{-\frac{k_1 + \text{RWC}}{k_2}} + 1} \text{ [MPa]} \quad (8)$$

including the inverse rate of change $-\frac{\partial T}{\partial t}$ in air temperature, ∂T , along time interval, ∂t , (since air temperature and water potential are negatively correlated), and a factor k_3 [-] to normalize for the distinctively different value ranges between XWP and T

$$k_3 = \frac{\text{XWP}_{\max} - \text{XWP}_{\min}}{T_{\max} - T_{\min}}. \quad (9)$$

The $-\frac{\partial T}{\partial t} * k_3$ -term in (8) was multiplied with the last increment of $|\text{FWP}_{\min}|$ to update FWP_{\min} . The air temperature-based FWP_{\min} -update was conducted in 24 h-intervals (from one daily mean value to the next). Daily updates were used instead of sub-daily ones to decouple FWP_{\min} -dynamics from the diurnal solar cycle. In some cases, strong alterations in water potential can happen during the day. Then, the short-term change in water potential is so strong that the transient and mild changes in




Fig. 5. Time series of m_g -estimates (blue and red colors) compared to their retrieval input parameter, vegetation optical depth (green color). Variation between the blue and red curves is due to the assumed vegetation volume fraction δ [-]: min = 0.001 (red solid), max = 0.9 (red dashed) and mean (blue) are calculated from minimum to maximum values.




Fig. 6. Comparison of in situ measurements (xylem water potential: brown curve; leaf water potential: blue dots) and radiometer-based estimates (green curve) of FWP, assuming a static FWP_{\min} .

daily air temperature update would not cover it. At these points in time (hereafter named breakpoints), the temperature-based FWP_{\min} was re-calibrated with the minimum of XWP in situ measurements at the day of the breakpoint.

IV. RESULTS AND ASSESSMENT OF FOREST WATER POTENTIAL ESTIMATION

In the following, FWP-estimates are shown for the intensive measurement periods in July, August, and September 2019. First, the different intermediate variables along the retrieval process are presented (cf. Section III). Since all variables were retrieved from tower-based L-band microwave radiometry observing an entire forest canopy (vertical extent), they can only be considered as effective canopy-scale variables compared to their location-based in situ counterparts given their spatial scale mismatch [41].

A. m_g Retrieval Results

The retrieval of m_g is shown in Fig. 5 for the intensive measurement period in July 2019. It includes an average stand






Fig. 7. Comparison of water potentials for the different intensive measuring periods in 2019 (Top: July, Middle: August, Bottom: September): Estimated from remote sensing (FWP in green for static (bright color) and dynamic FWP_{\min} (middle and dark color)), measured in situ water potential XWP (from xylem; in brown); gray dashed bars indicate the breakpoints (cf. Section III).

height of 21 m (top of canopy) provided by airborne Lidar scanning [73]. Since δ is generally not known for an L-band tower-based radiometer and optics-based laser scanning does not inform about δ seen by microwaves, a range of naturally occurring volume fractions covering the full physical range of δ from 0.001 to 0.9 was tested representing the vegetation structure and dry biomass influence on VOD.

In Fig. 5, the mean of m_g for the range of δ is located around 0.1, whereas the extremes of δ led to m_g -values from 0.03 to 0.6. The higher δ is, the more the temporal dynamics of m_g and VOD-dynamics differed significantly. This led to almost no dynamics in m_g for maximum δ , because vegetation structure and dry biomass dominated VOD (dashed red curve in Fig. 5). In contrast, at minimum δ , VOD dynamics are predominantly




Fig. 8. Average daily cycle of water potentials: Estimated from remote sensing (FWP in green for static (bright) and dynamic FWP_{\min} (middle and dark green)) and measured daily cycle of in situ water potentials: LWP (from leaves; black dashed line) and XWP (within the xylem; brown dashed line). Averaging was done over the intensive measuring period in July 2019. The green area shows the range over which the FWP from static FWP_{\min} was averaged. Grey arrows indicate the starting and the ending phase of the diel dynamics of FWP.

caused by water content variations leading to strong dynamics in m_g (solid red curve in Fig. 5). Also, higher values of δ led to lower values of m_g , while lower values of δ increase the estimated vegetation moisture. Importantly, choosing the minimum δ (i.e., $\delta = 0.001$) was appropriate because it led to m_g values between 0.3 and 0.6 [kg/kg].

This is in range with the average vegetation moisture conditions according to in situ data from oak species both at global scale ([82]; $n = 11118$; mean $m_g = 0.49$ [kg/kg]; std. $m_g = 0.07$ [kg/kg]) and, in specific regions such as the (drier) French Mediterranean area (Duché et al., 2017; $n = 2007$; mean $m_g = 0.44$ [kg/kg]; std. $m_g = 0.03$ [kg/kg]). In addition, choosing such low values for microwave observations agrees with literature [11], [69], [74], [79].

B. RWC Retrieval Results

Since δ is arduous to assess with experimental measurements at canopy scales and not known, neither generally nor for this study, RWC was calculated in two ways: first as RWC- m_g according to (4), second as RWC- VOD substituting m_g with VOD as shown in (5). Moreover, the minimum and maximum of the respective time series of the daily (24 h) extreme values is used in (4).

Using VOD instead of m_g in (4) had negligible effects when daily minimum and maximum values were considered (not shown). It is understood that VOD seems to be an equivalent predictor compared to m_g for plant water content for shorter studies (less than one month) [30]. This might be the case since dry biomass and vegetation structure (detectable at L-band) varied relatively little at Prospect Hill (Harvard Forest) during the intensive measurement weeks in July, August, and September, respectively.

C. FWP Retrieval Results

In the final processing step, an effective (as seen by an L-band radiometer system), canopy-scale FWP is calculated from the retrieved RWC-estimates using the individual RWC-FWP relationships presented in (6) to (9). In Fig. 6, the FWP-estimates based on RWC from m_g are compared with in situ measurements of XWP (brown curve) and LWP (blue dots). The temporal (daily) dynamics of the three potentials (FWP, XWP, LWP) show a promising concurrency in time, but a mismatch in absolute values (daily range of potentials). FWP- m_g was calculated in Fig. 6 with (6) assuming a time static FWP_{\min} along the week in July. This resulted in a constant lower boundary of FWP, which never reached values below -0.4 [MPa]. This FWP estimation method is called “static FWP_{\min} ” retrieval hereafter. In this static approximation, the constant FWP_{\min} was set to -2.5 [MPa]. This value was obtained from the models of Zweifel et al. [91], which are based on spruce (*Picea abies*). In the case of oak (*Quercus rubra*), the value chosen is also realistic for a healthy forest. This species has been found to start stomatal closure at LWP = -1.85 [MPa] [1], and change the xylem morphology due to progressive collapse at LWP between -2 [MPa] and -3 [MPa] [87]. Risk of cavitation for this species has been found at XWP = -3.10 [MPa] [46] and, for oaks in general, it occurs at XWP = -4 [MPa] (median value from 14 species in supplement data of [9]).

For dynamization along time, FWP_{\min} is updated continuously over time according to (7)–(9) to serve as a dynamic variable in (6). A time-dynamic FWP_{\min} was essential to guarantee the flexibility of FWP in terms of absolute values and different tension levels. RWC as input variable in (6) is a relative measure and only FWP_{\min} sets the lowest value in the FWP-range transforming into an absolute measure.

The update can be organized by incorporating available auxiliary data that is recorded independently and is easily available from contemporary in situ monitoring, like soil moisture (mass-based update) as in (7) or air temperature (energy-based update) as in (8)–(9).

The analysis of weekly FWP dynamics is shown in Fig. 7. It presents the retrieved FWP using either the static FWP_{\min} (green curve), or the time-variable FWP_{\min} , either updated by soil moisture (light green curve) or air temperature (dark green curve) in situ measurements.

In the July period, the soil moisture adaption led to an increasing daily dynamic of FWP especially around 16th to 18th of July that was not captured by XWP (brown curve). The air temperature adaption led to a similar dynamic of FWP in July, and stayed close to the static FWP dynamics. In contrast, the August period revealed that the temperature adapted FWP was able to trace partially (-1.6 to -0.2 [MPa]) the large dynamic range of the XWP (-1.9 to -0.2 MPa) in the beginning of the week (5th to 7th of August).

There is a distinct precipitation phase (>5 [mm]) in the afternoon of the 7th (see Fig. S1 in the Supplementary material) leading to a soil moisture level close to 40 [vol.%] around midnight. After this period (8th to 12th of August) XWP stayed close to saturation (-0.61 [MPa] to zero [MPa]). Qualitatively,

this was detected by both, the static and the temperature-based, FWP approaches.

The soil moisture-adapted FWP showed larger variation compared to XWP along these days. Additionally, daily variations of soil moisture-adapted FWP increased toward 13th of August due to constantly decreasing soil moisture down to values close to 20 [vol.%] (see Fig. S1 in the Supplementary material).

In contrast, the static FWP stayed between -0.4 [MPa] and 0 [MPa], which is expected, since FWP_{\min} is a constant calibrated once from in situ XWP on 10th of July. The same is true for the period in September and clearly indicates the importance of a time dynamic FWP_{\min} to meet the changing hydrological conditions along the week and throughout the year.

In the September period (Fig. 7 bottom), the daily minima of XWP showed the lowest values (down to -4.5 [MPa]) of all intensive measurement periods. Here, two rain events occurred on 15th and 23rd of September (cf. Fig. S1) leading to reduced (less negative) water potentials afterward. In addition, air temperature rose significantly from 19th to 23rd of September, depicted in Fig. S1. The temperature adapted FWP also demonstrates the capability to reach down to the lowest (most negative) water potentials in that week and indicates the strength of the break point mechanism for the temperature-based update (cf. Section III). However, this is not the case for the soil moisture adapted FWP which stayed between -1.0 and zero [MPa] and followed the hydro-dynamics guided by the two rain events during the intensive period.

D. Evaluation of the FWP Retrievals

For the intensive measurement period in July, results based on the static FWP_{\min} show FWP-values ranging between zero and -0.4 [MPa], like the dynamic range of in situ XWP (zero to -0.5 [MPa]) and shorter in dynamic range than in situ LWP (zero to -2 [MPa]). This suggests that FWP derived at L-band is sensitive to deeper canopy layers (trunk and branches) as well as to leaves (see Section IV-E for detailed analysis), which was expected due to the larger sensing depth at this wavelength and agrees with literature [39], [43], [68]. The dynamic range captured by the FWP-estimates, as well as that of the in situ data, show that the oak stand is generally not affected by hydric stress during July 2019, since there is no gradual decrease in the predawn water potential toward higher moisture deficits (more negative FWP-values). The results demonstrate, however, a constant minimum value of $FWP = -0.4$ [MPa], which means that the static method is not able to capture dynamics at the lower edge of the in situ XWP time-series due to the FWP_{\min} -fixation over time (see FWP-curve in Figs. 6 and 7).

It is confirmed here that FWP_{\min} needed to be updated for each individual time increment to enable the estimation of realistic FWP-values. We further validate the proposed two guiding variables (soil moisture & air temperature) for this estimation. The dynamic FWP_{\min} values were also within the expected range (between -0.5 and -6 [MPa]). For the lower limit case, note that the resulting FWP did not reach this minimum of -6 [MPa], which can only be found in some oak species [9].

At hourly to weekly timescales (cf. Fig. 7 and Table I), results revealed a moderate capacity to track XWP variability using the

TABLE I
SPEARMAN AND PEARSON CORRELATION COEFFICIENTS AND ROOT MEAN SQUARE ERROR (RMSE) FOR COMPARISON OF FWP AND XWP TIME SERIES ALONG THE THREE INTENSIVE PERIODS (JULY, AUGUST, SEPTEMBER) IN 2019

Retrieval Method	Intensive measuring period	Rank Correlation (Spearman)	Linear Correlation (Pearson)	RMSE [MPa]
Static FWP_{\min}	July	0.39	0.49	0.14
	August	0.08	0.13	0.61
	September	0.08	0.26	1.09
Soil moisture based FWP_{\min}	July	0.43	0.51	0.15
	August	0.12	0.14	0.56
	September	0.23	0.48	0.90
Air temperature based FWP_{\min}	July	0.38	0.48	0.14
	August	0.46	0.59	0.48
	September	0.42	0.60	0.83
VOD (no FWP retrieval)	July	0.30	0.32	-
	August	0.54	0.65	-
	September	0.62	0.69	-

air temperature dynamic approach ($0.48 < r < 0.60$), outperforming the capacity of the static and soil moisture approaches ($0.13 < r < 0.51$). Importantly, the temperature-driven approach for FWP was able to capture sudden and large declines (down to -4 [MPa]) of the in situ XWP (20th–24th September, Fig. 7) due to its break point mechanism (cf. Section III). This suggests that at coarser scales the proposed approach could help to capture cavitation-prone hydraulic conditions in forests.

E. Diurnal Dynamics of FWP

After investigating the weekly dynamics in Fig. 7, the study focused on FWP diurnal dynamics in Fig. 8. It shows the average daily cycle of in situ measurements (LWP, XWP) and radiometer-based FWP. In Fig. 8 FWP_{\min} was initially calibrated with XWP. This is why all FWP-curves are within the XWP-range (-0.4 to -0.1 [MPa]) and only partially within the LWP-range (-1.2 to -0.3 [MPa]). Nevertheless, a comparison of the time dynamics in relative changes along the day is appropriate. All FWP-curves indicated from 5 A.M. in the morning (dawning time; gray downward arrow in Fig. 8) an increased water demand by more negative water potential values. This was in phase with in situ LWP dynamics showing an increase at the same time. However, XWP measured at the tree stems reacted with a delay of several hours and the strongest tension increase occurred after 10 A.M. (negative slope of brown curve in Fig. 8). During this time, FWP was at a constant level around -0.3 [MPa].

After the heat maximum of the day (~ 2 P.M.) the water demand led to a decrease in LWP (leaves) first, followed by XWP (stems) at ~ 3 P.M. The XWP curve showed a stronger decrease in water demand from ~ 6 P.M. onward. At the same time FWP also started to move strongly towards reduced potential levels (gray upward arrow in Fig. 8). FWP dynamics (green curves in Fig. 8) coordinated in the morning with LWP (leaf dynamics) and in the evening with XWP (stem dynamics), implying that FWP is guided by leaf as well as by stem dynamics at the same time.




Fig. 9. Comparison of water potentials: FWP estimated from remote sensing with soil moisture-based FWP_{\min} dynamics (in green color), XWP measured in situ (in brown color) and soil matric potential SMP estimated from soil moisture measurements (in red color).

This makes sense since FWP has a sensing volume including leaves, twigs and branches down to stems due to the sensing depth of L-band microwaves.

At daily timescale, results demonstrate how all three methods captured morning drying and post-dusk water rehydration in the forest stand (gray arrows in Fig. 8). Further work should be directed to extend the approach so that it can capture minimum midday XWP, which is a key hydraulic measurement in plant ecology research (e.g., [9]).

F. Weekly and Monthly Dynamics of FWP

Fig. 9 shows the temporal extrapolation of FWP from July to September 2019 using the dynamic FWP_{\min} updated by the measured soil moisture. Here, soil moisture was chosen, since transient changes of FWP_{\min} over weeks from July to September 2019 are more attributed to soil moisture dynamics than to air temperature variations with a strong daily cycle at middle latitudes.

For reasons of unit equivalence, soil moisture was converted to soil matric potential [kPa]. These results are understood in the way that the trends and dynamics of FWP along the weeks match well with the measured XWP, except for large declines to very negative FWPs which are not captured by the proposed method in (7) (e.g., 7th August, or 11th and 23rd September).

This means that a potential extrapolation of the FWP time series and therefore conjunction of the different intensive measurement weeks by a soil moisture-updated FWP_{\min} could extend the FWP estimations to months and seasons, but further

investigations are needed to obtain an appropriate dynamic range of FWP.

V. DISCUSSION

Understanding how plants transport and use water is key to monitor and predict the impact of droughts on vegetation and their exchange with the atmosphere [88]. According to Novick et al. [59], there is a water potential information gap between ground-based observation networks and spatiotemporal scales relevant for Earth system models. This pathfinder study serves as a step towards addressing this gap and tracking water dynamics in the tree canopy over days, weeks and months. L-band radiometry was used for sensing the water dynamics of an oak stand at Harvard forest, USA, in the summer of 2019 as part of the SMAPVEX campaign.

The developed algorithm is based on a series of conversions to transform the radiometer-based, unitless VOD into absolute values of FWP. They include the conversion to relative water content (VOD or m_g to RWC) and then from relative-to-absolute metric (RWC to FWP).

There are still challenges, but also considerable potential to derive and estimate microwave-based FWP. Previous research has shown the ability to track this site's xylem and leaf water potential [30]. This study moved a step further, presenting conversion of VOD (microwave attenuation) to FWP (water potential) for a physically explicit assessment of water hydraulics in a forest stand from microwave radiometry. Results show how correlations between FWP estimates and XWP (in situ) are better (July) or only slightly lower (August and September) compared to the VOD–XWP correlation. This confirmed the potential utility of the method, providing such a conversion and tracking water potential in vegetation. In addition, the fact that FWP did not strongly improve the correlation with in situ water potential (if compared to VOD) was expected due to two shortcomings. First, VOD variations at short timescales are tightly linked to water status in the plants [41] and dry biomass dynamics were little during the study campaign, which qualifies VOD alone as a proxy to track the dynamics of water potentials [24]. In the case of agriculture, like in Jagdhuber et al. [34], the growing cycle with its change in phenology (plant structure, height and biomass) from sowing to harvest required a decomposition of VOD into a plant water and a plant biomass component to effectively extract RWC. Such dynamics would result in a greater change in dry biomass and vegetation structure than occurring at Prospect Hill between July and September 2019.

Second, additional parameterization, needed to convert VOD to FWP, requires several data sources and adds additional uncertainty to the results [89]. A key step during the FWP estimation is the definition of the pressure-volume curves (cf. Fig. 4). The period used to compute maximum and minimum VOD or m_g for RWC calculation includes only three months, which hampers the possibility of accounting for extreme dry and wet periods. This is partially compensated for by the high density of samples, although longer time-periods should be studied to test the sensitivity of the method. Hence, one major drawback of this study is that in situ comparison data from xylem or leaf water potential were not recorded continuously along entire months,

but for individual (unconnected) intensive measurement weeks. The resulting time gaps should be avoided, if possible in future campaigns.

Furthermore, although the pressure–volume curves are to some extent species-specific, the applied Zweifel et al. [91] model was not directly optimized for oak species. The main reason for choosing the applied model relies on its adaptability and easy computation (dependence on a low number of parameters, cf. Section III). This qualifies the proposed RWC-to-FWP model as a candidate for future satellite-scale FWP estimates which will require generalizable, nonspecies-specific and low-parameterized methods.

Next steps for future satellite-based water potential estimates also involve the application of appropriate technologies to track plant water potential. In that sense, the future Copernicus imaging microwave radiometer (CIMR) mission ([18]; expected to launch in 2029+) will provide passive microwave measurements of the land surface at L-, C-, X-, K-, and Ka-bands, offering different sensing depths and therefore different levels of canopy penetration, at an almost daily time scale globally. CIMR's consistent multifrequency microwave measurements will potentially allow sensing water potential from different canopy layers (e.g., leaf water potential from xylem water potential), which is paramount for an appropriate understanding of the plant hydraulics and the soil-plant-atmosphere continuum. In addition, this can ease the validation process of water potential estimates, as in situ measurements are provided separately, for leaves or for xylem, and enable the study of plant hydraulics behavior at short (diel) time scales [41]. For instance, the two in situ measurement networks VODnet [67] and PSInet¹ for water potential as well as ground-based VOD retrieval through GNSS-T are under current construction increasing their number of equipped stations around the world. VODnet is already equipped with site-scale ground-based GNSS receivers for transmissometry assessment [31]. Ultimately, both networks are ideal testbeds to link tree measures to satellite data. In addition, similar approaches could be applied using GNSS reflectometry (GNSS-R) linking ground-based VOD and tree water content retrievals (Rodríguez-Álvarez et al., 2012) to satellite-based estimates of VWC from the cyclone global navigation satellite system mission [86] and to in situ networks. However, it is worth noting that ground reflections in GNSS-R might complicate the estimation of canopy properties.

VI. CONCLUSION AND OUTLOOK

In this research study, a methodology is presented to estimate FWP for an oak stand in the Harvard Forest based on ground-based L-band radiometry. A way was evaluated to generate time series of dynamic FWP starting from VOD and transitioning through m_g and RWC. The Zweifel et al. [91] model was adapted to estimate time series of FWP. It is found that the incremental update of FWP_{\min} in the adapted model [see (7) and (8)] is crucial to obtain a realistic range of values compared to the in

situ observations of LWP and XWP and to estimate a continuous stream of FWP along weekly and monthly time scales.

FWP estimates showed the expected temporal dynamics, following the diurnal cycles and varying consistently with environmental drivers such as temperature. The correlation of the estimates with the in situ values confirmed the capacity of the method to track temporal variations in FWP. Consequently, this study provided a first valid approach to track vegetation water potential in a forest using the synergy between remotely sensed VOD and ancillary data through a model-based retrieval.

The achievement of absolute accuracy in FWP estimation still depends on the availability and quality of the auxiliary information. The needed accuracy will be defined in close coordination with the end users of FWP products. Here, mostly forest ecologists and tree physiologists will benefit from spatio-temporal water potential estimates from plot to ecosystem scales. In this regard, the presented approach builds the basic foundation for transitioning towards a remote sensing-based, up to ecosystem-scale estimation of FWP dynamics with limited dependency on ancillary data. In the context of ecosystem-scale FWP retrievals, the current limitation of a mono-species (homogeneous oak stand) retrieval needs to be overcome. One of the next challenges is transferring the developed approach to the ecosystem scale, including various plant species representing different physiological traits.

Furthermore, FWP retrievals in this study cannot be separated into different vertical forest compartments, like trunks, branches, lower leaf (mostly shaded) canopy, upper leaf (mostly sunlit) canopy, due to the mono-frequency retrieval (only L-band). Upcoming microwave missions like NISAR (NASA-ISRO SAR) from the National Aeronautics and Space Administration and the Indian Space Research Organization, or BIOMASS and CIMR from the European Space Agency, will provide the possibility to acquire VOD globally and for different forest compartments (upper canopy, lower canopy, trunks) due to recordings of multi-frequency (K-, X-, C-, L-, and P-band) microwave data at resolutions of tens of meters. However, research on VOD estimation from SAR (active microwave) techniques is still urgently needed to enable high resolution VOD and FWP products (e.g., [20], [90]).

REFERENCES

- [1] M. D. Abrams, "Adaptations and responses to drought in *Quercus* species of North America," *Tree Physiol.*, vol. 7, pp. 227–238, 1990.
- [2] W. R. L. Anderegg et al., "Plant water potential improves prediction of empirical stomatal models," *PLoS One*, vol. 12, 2017, Art. no. e0185481, doi: [10.1371/journal.pone.0185481](https://doi.org/10.1371/journal.pone.0185481).
- [3] A. Balling and U. Zimmermann, "Comparative measurements of the xylem pressure of *Nicotiana* plants by means of the pressure bomb and pressure probe," *Planta*, vol. 182, pp. 325–338, 1990.
- [4] J. Barceló-Coll, G. Nicolás-Rodrigo, B. Sabater-García, and R. Sánchez-Tamés, *Fisiología Vegetal*, 5th ed. Madrid, Spain: Ediciones Pirámide S.A., 1980.
- [5] M. J. Baur, T. Jagdhuber, A. F. Feldman, R. Akbar, and D. Entekhabi, "Estimation of relative canopy absorption and scattering at L-, C- and X-bands," *Remote Sens. Environ.*, vol. 233, 2019, Art. no. 111384.
- [6] G. B. Bonan, M. Williams, R. A. Fisher, and K. W. Oleson, "Modeling stomatal conductance in the earth system: Linking leaf water-use efficiency and water transport along the soil–plant–atmosphere continuum," *Geoscientific Model Develop.*, vol. 7, no. 5, pp. 2193–2222, 2014.

1.[Online]. Available: <https://psinetrcn.github.io/>

- [7] D. Chaparro et al., "Global L-band vegetation volume fraction estimates for modeling vegetation optical depth," in *Proc. IEEE Int. Geosci. Remote Sens. Symp.*, 2021, pp. 6399–6402.
- [8] D. Chaparro et al., "Vegetation moisture estimation in the Western United States using radiometer-radar-LiDAR synergy," *Remote Sens. Environ.*, vol. 303, 2024, Art. no. 113993.
- [9] B. Choat et al., "Global convergence in the vulnerability of forests to drought," *Nature*, vol. 491, no. 7426, pp. 752–755, 2012.
- [10] B. Choat et al., "Triggers of tree mortality under drought," *Nature*, vol. 558, pp. 531–539, 2018, doi: [10.1038/s41586-018-0240-x](https://doi.org/10.1038/s41586-018-0240-x).
- [11] B. J. Choudhury and P. Pampaloni, Eds., *Passive Microwave Remote Sensing of Land–Atmosphere Interactions*. Boca Raton, FL, USA: CRC Press, 1995.
- [12] B. O. Christoffersen et al., "Linking hydraulic traits to tropical forest function in a size-structured and trait-driven model (TFS v.1-Hydro)," *Geosci. Model Dev.*, vol. 9, pp. 4227–4255, 2016, doi: [10.5194/gmd-9-4227-2016](https://doi.org/10.5194/gmd-9-4227-2016).
- [13] M. A. Clark, J. Choi, and M. Douglas, *Biology 2e*. Houston, TX, USA: Rice Univ., 2018.
- [14] A. Colliander et al., "SMAP detects soil moisture under temperate forest canopies," *Geophysical Res. Lett.*, vol. 47, no. 19, 2020, Art. no. e2020GL089697.
- [15] I. Cowan, "Transport of water in the soil–plant–atmosphere system," *J. Appl. Ecol.*, vol. 2, pp. 221–239, 1965.
- [16] D. Chaparro et al., "Estimation of gravimetric vegetation moisture in the western United States using a multi-sensor approach," in *Proc. Int. Geosci. Remote Sens. Symp.*, 2023, pp. 1505–1508.
- [17] M. A. Dixon and M. T. Tyree, "A new stem hygrometer, corrected for temperature gradients and calibrated against the pressure bomb," *Plant, Cell Environ.*, vol. 7, pp. 693–697, 1984.
- [18] C. J. Donlon, "The copernicus imaging microwave radiometer (CIMR) mission requirements document v5.0," ESA, Paris, France, 2023. [Online]. Available: https://cimr.eu/mrd_v5
- [19] D. R. Edwards and M. A. Dixon, "Mechanisms of drought response in *Thuja occidentalis* LI water stress conditioning and osmotic adjustment," *Tree Physiol.*, vol. 15, no. 2, pp. 121–127, 1995.
- [20] M. El Hajj et al., "First vegetation optical depth mapping from Sentinel-1 C-band SAR data over crop fields," *Remote Sens.*, vol. 11, no. 23, 2019, Art. no. 2769.
- [21] C. B. Eller et al., "Stomatal optimization based on xylem hydraulics (SOX) improves land surface model simulation of vegetation responses to climate," *New Phytol.*, vol. 226, pp. 1622–1637, 2020, doi: [10.1111/nph.16419](https://doi.org/10.1111/nph.16419).
- [22] D. Entekhabi et al., "The soil moisture active passive (SMAP) mission," *Proc. IEEE*, vol. 98, no. 5, pp. 704–716, May 2010.
- [23] D. Entekhabi et al., "SMAP handbook soil moisture active passive: Mapping soil moisture and freeze/thaw from space," 2014.
- [24] A. F. Feldman, D. J. Short Gianotti, A. G. Konings, P. Gentine, and D. Entekhabi, "Patterns of plant rehydration and growth following pulses of soil moisture availability," *Biogeosciences*, vol. 18, pp. 831–847, 2021.
- [25] M. Forkel, L. Schmidt, R. M. Zotta, W. Dorigo, and M. Yebra, "Estimating leaf moisture content at global scale from passive microwave satellite observations of vegetation optical depth," *Hydrol. Earth Syst. Sci. Discuss.*, vol. 27, pp. 39–68, 2023.
- [26] W. R. Gardner, "Dynamic aspects of water availability to plants," *Soil Sci.*, vol. 89, pp. 63–73, 1960.
- [27] W. R. Gardner, "Dynamic aspects of soil-water availability to plants," *Ann. Rev. Plant Physiol.*, vol. 16, pp. 323–342, 1965.
- [28] D. Helman et al., "Using time series of high-resolution planet satellite images to monitor grapevine stem water potential in commercial vineyards," *Remote Sens.*, vol. 10, no. 10, 2018, Art. no. 1615.
- [29] N. Holtzman, A. G. Konings, A. Roy, and A. Colliander, "SMAPVEX19-21 Massachusetts vegetation optical depth, version 1," NASA Nat. Snow Ice Data Center Distributed Active Archive Center, Boulder, CO, USA, Dec. 2020, doi: [10.5067/2PJZJURUJLWF](https://doi.org/10.5067/2PJZJURUJLWF).
- [30] N. M. Holtzman et al., "L-band vegetation optical depth as an indicator of plant water potential in a temperate deciduous forest stand," *Biogeosciences*, vol. 18, no. 2, pp. 739–753, 2021.
- [31] V. Humphrey and C. Frankenberg, "Continuous ground monitoring of vegetation optical depth and water content with GPS signals," *Biogeosciences*, vol. 20, pp. 1789–1811, 2023.
- [32] T. Hwang et al., "Capturing species-level drought responses in a temperate deciduous forest using ratios of photochemical reflectance indices between sunlit and shaded canopies," *Remote Sens. Environ.*, vol. 199, pp. 350–359, 2017.
- [33] T. J. Jackson and T. J. Schmugge, "Vegetation effects on the microwave emission of soils," *Remote Sens. Environ.*, vol. 36, no. 3, pp. 203–212, 1991.
- [34] T. Jagdhuber et al., "Toward estimation of seasonal water dynamics of winter wheat from ground-based L-band radiometry: A concept study," *Biogeosciences*, vol. 19, no. 8, pp. 2273–2294, 2022.
- [35] F. Jonard, F. André, Q. Ponette, C. Vincke, and M. Jonard, "Sap flux density and stomatal conductance of European beech and common oak trees in pure and mixed stands during the summer drought of 2003," *J. Hydrol.*, vol. 409, pp. 371–381, 2011, doi: [10.1016/j.jhydrol.2011.08.032](https://doi.org/10.1016/j.jhydrol.2011.08.032).
- [36] S. Junttila et al., "Terrestrial laser scanning intensity captures diurnal variation in leaf water potential," *Remote Sens. Environ.*, vol. 255, 2021, Art. no. 112274.
- [37] D. Kennedy et al., "Implementing plant hydraulics in the community land model, version 5," *J. Adv. Model. Earth Syst.*, vol. 11, pp. 485–513, 2019, doi: [10.1029/2018MS001500](https://doi.org/10.1029/2018MS001500).
- [38] Y. H. Kerr, P. Waldteufel, J. -P. Wigneron, J. A. M. J. Martinuzzi, J. Font, and M. Berger, "Soil moisture retrieval from space: The soil moisture and ocean salinity (SMOS) mission," *IEEE Trans. Geosci. Remote Sens.*, vol. 39, no. 8, pp. 1729–1735, Aug. 2001.
- [39] A. G. Konings and P. Gentine, "Global variations in ecosystem-scale isohydricity," *Glob. Change Biol.*, vol. 23, no. 2, pp. 891–905, 2017.
- [40] A. G. Konings, K. Rao, and S. C. Steele-Dunne, "Macro to micro: Microwave remote sensing of plant water content for physiology and ecology," *New Phytologist*, vol. 223, no. 3, pp. 1166–1172, 2019.
- [41] A. G. Konings et al., "Detecting forest response to droughts with global observations of vegetation water content," *Glob. Change Biol.*, vol. 27, no. 23, pp. 6005–6024, 2021.
- [42] H. Lambers, F. Stuart Chapin III, and T. L. Pons, *Plant Physiological Ecology*. Berlin, Germany: Springer, 2008.
- [43] X. Li et al., "Global-scale assessment and inter-comparison of recently developed/reprocessed microwave satellite vegetation optical depth products," *Remote Sens. Environ.*, vol. 253, 2021, Art. no. 112208.
- [44] Y. Lin, Z. Zhu, W. Guo, Y. Sun, X. Yang, and V. Kovalsky, "Continuous monitoring of cotton stem water potential using sentinel-2 imagery," *Remote Sens.*, vol. 12, no. 7, 2020, Art. no. 1176.
- [45] Y. Liu, M. Kumar, G. G. Katul, X. Feng, and A. G. Konings, "Plant hydraulics accentuates the effect of atmospheric moisture stress on transpiration," *Nature Climate Change*, vol. 10, pp. 691–695, 2020, doi: [10.1038/s41558-020-0781-5](https://doi.org/10.1038/s41558-020-0781-5).
- [46] H. Maherali, C. F. Moura, M. C. Caldeira, C. J. Willson, and R. B. Jackson, "Functional coordination between leaf gas exchange and vulnerability to xylem cavitation in temperate forest trees," *Plant, Cell Environ.*, vol. 29, pp. 571–583, 2006.
- [47] J. Martínez-Vilalta and N. García-Fórner, "Water potential regulation, stomatal behaviour and hydraulic transport under drought: Deconstructing the iso/anisohydric concept," *Plant Cell Environ.*, vol. 40, pp. 962–976, 2017.
- [48] J. Martínez-Vilalta, W. R. Anderegg, G. Sapes, and A. Sala, "Greater focus on water pools may improve our ability to understand and anticipate drought-induced mortality in plants," *New Phytol.*, vol. 223, pp. 22–32, 2019.
- [49] A. M. Matheny, G. Mirfenderesgi, and G. Bohrer, "Trait-based representation of hydrological functional properties of plants in weather and ecosystem models," *Plant Diversity*, vol. 39, pp. 1–12, 2017.
- [50] N. McDowell et al., "Mechanisms of plant survival and mortality during drought: Why do some plants survive while others succumb to drought?," *New Phytologist*, vol. 178, no. 4, pp. 719–739, 2008.
- [51] P. J. Melcher, F. C. Meinzer, D. E. Yount, G. Goldstein, and U. Zimmermann, "Comparative measurements of xylem pressure in transpiring and non-transpiring leaves by means of the pressure chamber and the xylem pressure probe," *J. Exp. Botany*, vol. 49, pp. 1757–1760, 1998.
- [52] T. Meyer, L. Weihermüller, H. Vereecken, and F. Jonard, "Vegetation optical depth and soil moisture retrieved from L-band radiometry over the growth cycle of a winter wheat t," *Remote Sens.*, vol. 10, no. 10, 2018, Art. no. 1637, doi: [10.3390/rs10101637](https://doi.org/10.3390/rs10101637).
- [53] T. Meyer et al., "Estimating gravimetric water content of a winter wheat field from L-band vegetation optical depth," *Remote Sens.*, vol. 11, no. 20, 2019, Art. no. 2353, doi: [10.3390/rs11202353](https://doi.org/10.3390/rs11202353).
- [54] G. Mirfenderesgi et al., "Tree level hydrodynamic approach for resolving aboveground water storage and stomatal conductance and modeling the effects of tree hydraulic strategy," *J. Geophysical Res., Biogeosciences*, vol. 121, no. 7, pp. 1792–1813, 2016.

- [55] T. Mo, B. J. Choudhury, T. J. Schmugge, J. R. Wang, and T. J. Jackson, "A model for microwave emission from vegetation-covered fields," *J. Geophysical Res., Oceans*, vol. 87, pp. 11229–11237, 1982.
- [56] R. H. Nolan et al., "Linking forest flammability and plant vulnerability to drought," *Forests*, vol. 11, 2020, Art. no. 779, doi: [10.3390/f11070779](https://doi.org/10.3390/f11070779).
- [57] K. A. Novick et al., "The increasing importance of atmospheric demand for ecosystem water and carbon fluxes," *Nature Climate Change*, vol. 6, pp. 1023–1027, 2016, doi: [10.1038/nclimate3114](https://doi.org/10.1038/nclimate3114).
- [58] K. A. Novick, A. G. Konings, and P. Gentine, "Beyond soil water potential: An expanded view on isohydrlicity including land–atmosphere interactions and phenology," *Plant, Cell Environ.*, vol. 42, no. 6, pp. 1802–1815, 2019.
- [59] K. A. Novick et al., "Confronting the water potential information gap," *Nature Geosci.*, vol. 15, no. 3, pp. 158–164, 2022.
- [60] A. R. Palmer et al., "Using pre-dawn leaf water potential and MODIS LAI to explore seasonal trends in the phenology of Australian and southern African woodlands and savannas," *Australian J. Botany*, vol. 56, pp. 557–563, 2008.
- [61] A. R. Palmer et al., "Towards a spatial understanding of water use of several land-cover classes: An examination of relationships amongst pre-dawn leaf water potential, vegetation water use, aridity and MODIS LAI," *Ecohydrogeomorphology*, vol. 3, no. 1, pp. 1–10, 2010.
- [62] K. Rao, W. R. Anderegg, A. Sala, J. Martínez-Vilalta, and A. G. Konings, "Satellite-based vegetation optical depth as an indicator of drought-driven tree mortality," *Remote Sens. Environ.*, vol. 227, pp. 125–136, 2019.
- [63] N. Rodriguez-Alvarez et al., "Vegetation water content estimation using GNSS measurements," *IEEE Geosci. Remote Sens. Lett.*, vol. 9, no. 2, pp. 282–286, Mar. 2012.
- [64] C. M. Rodriguez-Dominguez et al., "Leaf water potential measurements using the pressure chamber: Synthetic testing of assumptions towards best practices for precision and accuracy," *Plant, Cell Environ.*, vol. 45, pp. 2037–2061, 2022.
- [65] T. L. Rowlandson et al., "Capturing agricultural soil freeze/thaw state through remote sensing and ground observations: A soil freeze/thaw validation campaign," *Remote Sens. Environ.*, vol. 211, pp. 59–70, 2018.
- [66] A. Roy et al., "L-band response to freeze/thaw in a boreal forest stand from ground-and tower-based radiometer observations," *Remote Sens. Environ.*, vol. 237, 2020, Art. no. 111542.
- [67] K. Schellenberg et al., "Estimating canopy interception water storage with GNSS-transmissometry," in *Proc. IEEE Int. Geosci. Remote Sens. Symp.*, 2024, pp. 4507–4510.
- [68] L. Schmidt et al., "Assessing the sensitivity of multi-frequency passive microwave vegetation optical depth to vegetation properties," *Biogeosciences*, vol. 20, no. 5, pp. 1027–1046, 2023.
- [69] T. J. Schmugge and T. J. Jackson, "A dielectric model of the vegetation effects on the microwave emission from soils," *IEEE Trans. Geosci. Remote Sens.*, vol. 30, no. 4, pp. 757–760, Jul. 1992.
- [70] P. F. Scholander, E. D. Bradstreet, E. A. Hemmingens, and H. T. Hammel, "Sap pressure in vascular plants: Negative hydrostatic pressure can be measured in plants," *Science*, vol. 148, no. 3668, pp. 339–346, 1965, doi: [10.1126/scien.148.3668.339](https://doi.org/10.1126/scien.148.3668.339).
- [71] R. O. Slatyer and S. A. Taylor, "Terminology in plant-and soil-water relations," *Nature*, vol. 187, pp. 922–924, 1960.
- [72] R. E. Smart and G. E. Bingham, "Rapid estimates of relative water content," *Plant Physiol.*, vol. 53, no. 2, pp. 258–260, 1974.
- [73] F. B. Sullivan et al., "Comparison of LiDAR- and allometry-derived canopy height models in an eastern deciduous forest," *Forest Ecol. Manage.*, vol. 406, pp. 83–94, 2017.
- [74] F. T. Ulaby, M. Razani, and M. C. Dobson, "Effects of vegetation cover on the microwave radiometric sensitivity to soil moisture," *IEEE Trans. Geosci. Remote Sens.*, vol. GE-21, no. 1, pp. 51–61, Jan. 1983.
- [75] F. T. Ulaby and M. A. El-Rayes, "Microwave dielectric spectrum of vegetation-part II: Dual-dispersion model," *IEEE Trans. Geosci. Remote Sens.*, vol. GE-25, no. 5, pp. 550–557, Sep. 1987.
- [76] F. Ulaby and D. Long, *Microwave Radar and Radiometric Remote Sensing*. Ann Arbor, MI, USA: Univ. Michigan Press, 2014.
- [77] T. H. Van den Honert, "Water transport in plants as a catenary process," *Discuss. Faraday Soc.*, vol. 3, pp. 146–153, 1948.
- [78] M. D. Venturas, J. S. Sperry, and U. G. Hacke, "Plant xylem hydraulics: What we understand, current research, and future challenges: Plant xylem hydraulics," *J. Integr. Plant Biol.*, vol. 59, pp. 356–389, 2017, doi: [10.1111/jipb.12534](https://doi.org/10.1111/jipb.12534).
- [79] J. P. Wigneron, Y. Kerr, A. Chanzy, and Y. Q. Jin, "Inversion of surface parameters from passive microwave measurements over a soybean field," *Remote Sens. Environ.*, vol. 46, no. 1, pp. 61–72, 1993.
- [80] A. P. Williams et al., "Temperature as a potent driver of regional forest drought stress and tree mortality," *Nature Climate Change*, vol. 3, pp. 292–297, 2013, doi: [10.1038/nclimate1693](https://doi.org/10.1038/nclimate1693).
- [81] X. Xu, D. Medvigy, J. S. Powers, J. M. Becknell, and K. Guan, "Diversity in plant hydraulic traits explains seasonal and inter-annual variations of vegetation dynamics in seasonally dry tropical forests," *New Phytol.*, vol. 212, pp. 80–95, 2016, doi: [10.1111/nph.14009](https://doi.org/10.1111/nph.14009).
- [82] M. Yebra et al., "Globe-LFMC, a global plant water status database for vegetation ecophysiology and wildfire applications," *Sci. Data*, vol. 6, no. 1, 2019, Art. no. 155.
- [83] Y. Yvon Duché, R. Savazzi, M. Toutchkov, and E. Cabanne, "Multisite and multispecies live fuel moisture content (LFMC) series in the french mediterranean since 1996 [Data set]," *Zenodo*, 2017, doi: [10.5281/zenodo.162978](https://doi.org/10.5281/zenodo.162978).
- [84] P. J. Zarco-Tejada, C. A. Rueda, and S. L. Ustin, "Water content estimation in vegetation with MODIS reflectance data and model inversion methods," *Remote Sens. Environ.*, vol. 85, pp. 109–124, 2003.
- [85] P. J. Zarco-Tejada et al., "A PRI-based water stress index combining structural and chlorophyll effects: Assessment using diurnal narrow-band airborne imagery and the CWSI thermal index," *Remote Sens. Environ.*, vol. 138, pp. 38–50, 2013.
- [86] Y. Zhang, J. Bu, X. Zuo, K. Yu, Q. Wang, and W. Huang, "Vegetation water content retrieval from spaceborne GNSS-R and multi-source remote sensing data using ensemble machine learning methods," *Remote Sens.*, vol. 16, 2024, Art. no. 2793.
- [87] Y. J. Zhang, F. E. Rockwell, A. C. Graham, T. Alexander, and N. M. Holbrook, "Reversible leaf xylem collapse: A potential 'circuit breaker' against cavitation," *Plant Physiol.*, vol. 172, no. 4, pp. 2261–2274, 2016.
- [88] Y. Zhang, S. Zhou, P. Gentine, and X. Xiao, "Can vegetation optical depth reflect changes in leaf water potential during soil moisture dry-down events?," *Remote Sens. Environ.*, vol. 234, 2019, Art. no. 111451.
- [89] M. Zhao, V. Humphrey, A. F. Feldman, and A. G. Konings, "Temperature is likely an important omission in interpreting vegetation optical depth," *Geophysical Res. Lett.*, vol. 51, pp. 1–11, 2024.
- [90] Z. Zhou et al., "Retrieval of high-resolution vegetation optical depth from Sentinel-1 data over a grassland region in the Heihe River Basin," *Remote Sens.*, vol. 14, 2022, Art. no. 5468, doi: [10.3390/rs14215468](https://doi.org/10.3390/rs14215468).
- [91] R. Zweifel, H. Item, and R. Häsler, "Link between diurnal stem radius changes and tree water relations," *Tree Physiol.*, vol. 21, no. 12/13, pp. 869–877, 2001.
- [92] A. Fink et al., "Estimating gravimetric moisture of vegetation using an attenuation-based multisensor approach," *IEEE Int. Geosci. Remote Sens. Symp.*, pp. 353–356, Jul. 2018.



Thomas Jagdhuber (Senior Member, IEEE) received the Diploma degree in physical geography, physics, remote sensing, and geoinformatics from the Ludwig Maximilian University of Munich (LMU), Munich, Germany, and the Technical University (TUM) of Munich, Munich, in 2006, and the Ph.D. degree in hydrology from the Faculty of Science, University of Potsdam, Potsdam, Germany, in 2012.

Since 2007, he has been with the Microwaves and Radar Institute (HR), German Aerospace Center (DLR), Weßling, Germany, and since 2022, he has been leading the signatures research group of HR. From 2014 to 2019 and 2022 to 2023, he was a Yearly Visiting Scientist with the Massachusetts Institute of Technology (MIT), Boston, MA, USA, contributing to the preparation and continuation of the Soil Moisture Active Passive (SMAP) and SMAP/Sentinel-1 missions. He is also a Lecturer with the University of Jena, Jena, Germany, and a habilitation candidate as well as an Associate Lecturer with the University of Augsburg, Augsburg, Germany. His main research interests include physics-based multisensor data integration with a focus on active and passive microwave interaction theory and polarimetric techniques for hydrological, agricultural, ecological, and cryospheric parameter modeling and estimation.

Dr. Jagdhuber was the recipient of the DLR Science Award for his research on polarimetric SAR decomposition techniques in 2014. Together with Prof. Entekhabi (MIT), he was awarded the MIT-MISTI Grant for global water cycle and environmental monitoring using active and passive satellite-based microwave instruments. He also serves as a Reviewer for several international journals and conference boards.



Anne-Sophie Schmidt received the B.S. degree in geography and the M.S. degree in climate and environment science from the University Augsburg, Augsburg, Germany, in 2019 and 2022, respectively. Since 2022, she has been the team leader in industry responsible for installing power and fiber optic lines.



Anke Fluhrer (Member, IEEE) received the B.S. degree in geography from the Julius-Maximilians University Würzburg, Würzburg, Germany, in 2014, and the M.S. degree in physical geography from the Friedrich-Alexander University Erlangen-Nuremberg, Erlangen, Germany, in 2018.

Since 2018, she has been a Research Associate with the prospect of a Ph.D. with the Microwaves and Radar Institute (HR), German Aerospace Center (DLR), Weßling, Germany. Her research interests include terrestrial microwave remote sensing, satellite

image processing, geoinformatics, and data assimilation.



David Chaparro (Member, IEEE) received the B.S. degree in biology from the Universitat Autònoma de Barcelona (UAB), Barcelona, Spain, in 2011, the M.S. degree in remote sensing and geographic information systems from the Centre for Ecological Research and Forestry Applications (CREAF), Cerdanyola del Vallès, Spain, in 2012, the M.S. degree in terrestrial ecology from UAB, in 2013, and the Ph.D. degree in remote sensing (telecommunications engineering) from the Universitat Politècnica de Catalunya (UPC), Barcelona, in 2018.

He has been a Visiting Scientist with the Massachusetts Institute of Technology for four months (2019), Postdoc Researcher with UPC (2018–2021) and the German Aerospace Center, Weßling, Germany (2022–2023). He is currently a Postdoc Researcher with the Ecosystem Physiology and Global Change (EcoPhys) group, CREAM. At CREAM, he aims at bridging research from the plant ecophysiology and the remote sensing communities leading collaboration between them. His research interests include microwave remote sensing, with experience on sensors synergy (microwaves, optical, and lidar) for vegetation moisture retrievals from space.



François Jonard (Senior Member, IEEE) received the M.Sc. and Ph.D. degrees in bioscience engineering from the University of Louvain (UCLouvain), Louvain-la-Neuve, Belgium, in 2002 and 2012, respectively.

From 2009 to 2021, he was a Research Scientist with the Institute of Bio- and Geosciences (Agrosphere), Research Centre Jülich, Jülich, Germany. In 2011, he was a Visiting Scientist with the NASA Goddard Space Flight Center, Greenbelt, MD, USA, where he contributed to the preparation of the Soil

Moisture Active and Passive mission. From 2018 to 2024, he was a Visiting Scientist with the Massachusetts Institute of Technology (MIT), Cambridge, MA, USA, under an MIT-MISTI grant, with short-term stays focusing on monitoring Earth's ecosystem health using next-generation remote sensing satellites. Between 2015 and 2020, he served as an Associate Professor with the Faculty of Bioscience Engineering and the Earth and Life Institute, UCLouvain. In 2021, he joined the Faculty of Sciences at the University of Liège, Liège, Belgium, as an Associate Professor, where he leads the Earth Observation and Ecosystem Modelling Laboratory. Additionally, since 2020, he has been a Guest Professor with UCLouvain. His research interests include remote sensing for Earth observation, ecosystem modelling, land-surface hydrology, ecophysiology, and hydrogeophysics. He focuses particularly on the interactions between the water and carbon cycles, with an emphasis on the impact of drought on forest ecosystems and their resilience to climate change.



María Piles (Senior Member, IEEE) received the B.Sc. degree in telecommunication engineering from Universitat Politècnica de València, València, Spain, in 2005, and the Ph.D. degree in signal theory and communications from Universitat Politècnica de Catalunya (UPC), Barcelona, Spain, in 2010.

She was a Research Scientist with Melbourne University, the Institute of Marine Sciences, CSIC, and UPC, and a Research Affiliate with the Massachusetts Institute of Technology. In 2017, she joined the Image and Signal Processing Group (ISP, <https://isp.uv.es/>),

Universitat de València, as a Ramón y Cajal Senior Researcher, and was promoted to an Associate Professor in 2023. She has advised more than 15 master's and seven Ph.D. theses (2 underway). She teaches courses in signal theory, image processing, and remote sensing in several engineering and data science degrees. Her research interests include remote sensing for Earth Observation (EO), estimation of soil moisture and vegetation biogeophysical parameters from microwave satellites, and development of multimodal techniques that exploit EO data and machine learning to model and understand the Earth system with a focus on agriculture, forestry, wildfire prediction, extreme detection, and climate studies.

Dr. Piles is a Fellow of the European Laboratory for Learning and Intelligent Systems. She is the past President of the IEEE Spain Geoscience and Remote Sensing Society Chapter.



Natan Holtzman was born in Raleigh, NC, USA, in 1994. He received the B.S. (Hons.) degree in geological sciences from the University of North Carolina at Chapel Hill, Chapel Hill NC, USA, in 2016, and the Ph.D. degree in Earth system science from Stanford University, Stanford, CA, USA, in 2014.

He is currently a Senior Member of Technical Staff with The Aerospace Corporation, El Segundo, CA, USA, with a specialization in signal and image processing and physics. As part of the Microelectronics Technology Department within the Aerospace

Physical Science Laboratories, he performs computational experiments and data analyses to study electronic devices for use in spacecraft. From 2018 to 2024, he was a Graduate Researcher with the Remote Sensing Ecohydrology Laboratory, Stanford University, where he developed techniques to characterize Earth surface vegetation and soil moisture with microwave radiometry and model-data fusion. From 2017 to 2018, he was a Research Associate with the University of North Carolina Institute for the Environment. Dr. Holtzman was awarded the NASA Future Investigators in Earth and Space Sciences and Technology fellowship to fund his graduate studies. He has authored or coauthored ten journal articles and presented at five conferences including the American Geophysical Union Fall Meeting and the Gordon Research Conference on Multi-Scale Plant Vascular Biology.



Alexandra G. Konings received the M.S. degree in environmental science from Duke University, Durham, NC, USA, in 2011, and the Ph.D. degree in hydrology from the Massachusetts Institute of Technology, Cambridge, MA, USA, in 2015.

In 2015–2016, she was a Postdoctoral Fellow with Columbia University and in 2016 she was a Postdoctoral Fellow with the NASA Jet Propulsion Laboratory. Since 2016, she has been an Assistant Professor with the Department of Earth System Science, Stanford University, Stanford, CA, USA, becoming

an Associate Professor in the same department in 2023. Her research interests include understanding water–carbon interactions at continental to global scale, with a particular interest in microwave remote sensing of vegetation water content and its applications.



Andrew F. Feldman received the B.S. and M.S. degrees in civil engineering from Drexel University, Philadelphia, PA, USA, in 2016, and the S.M. and Ph.D. degrees in civil and environmental engineering from the Massachusetts Institute of Technology, Cambridge, MA, USA, in 2018 and 2021, respectively.

He is currently an Assistant Research Scientist with NASA Goddard Space Flight Center Biospheric Sciences Laboratory and University of Maryland, Greenbelt, MD, USA. He was a NASA Postdoctoral Program Fellow with the NASA Goddard Space Flight Center. His research interests include land-surface hydrology, terrestrial ecosystem science, and microwave and thermal remote sensing.

Dr. Feldman is a member of the American Geophysical Union and Ecological Society of America.



Konstantin Schellenberg received the M.Sc. degree in geoinformatics from Friedrich Schiller University Jena, Jena, Germany, in 2022. He is currently working toward the Ph.D. degree in the International Max Planck Research School (IMPRS), Jena.

He is jointly affiliated with the Max Planck Institute for Biogeochemistry, Jena, the Microwaves and Radar Institute of the German Aerospace Center (DLR) and the Friedrich Schiller University Jena. He currently leads a suite of in situ measurements of tree hydraulic parameters for remote sensing model validation. His research interests include forest microwave remote sensing and hydrological parameter retrieval for tackling research questions in tree mortality and forest resilience.



Martin J. Baur received the B.S. degree in geography from the University of Augsburg, Augsburg, Germany, in 2016, and the M.S. degree in global change ecology from the University of Bayreuth, Bayreuth, Germany, in 2019. He is currently working toward the Ph.D. degree in leveraging remote sensing-based information on plant water status for ecosystem models from the University of Cambridge, Cambridge, U.K.

Since 2017, he has been collaborating with the Microwaves and Radar Institute, German Aerospace Center (DLR), Wessling, Germany. Since 2018, he

has been a Visiting Researcher with the Department of Civil and Environmental Engineering, MIT, Cambridge, MA, USA, and more recently with the Department of Environmental Systems Science, ETH Zurich, Zurich, Switzerland. His research interests include multifrequency retrievals of soil moisture and vegetation water content based on the SMAP and AMSR2 missions. He is also conducting research on soil and plant water dynamics as well as the effects of disturbances on ecohydrology.



Harald Kunstmann received the Diploma degree in physics from the University of Heidelberg, Heidelberg, Germany, in 1994, and the Ph.D. degree in environmental natural sciences from ETH Zürich, Zürich, Switzerland, in 1998.

He is currently a Professor and the Chair for regional climate and hydrology with the University of Augsburg in joint appointment with Karlsruhe Institute of Technology, Campus Alpin, Garmisch-Partenkirchen, Germany. His research interests include hydrological climate change impacts, high-resolution coupled atmosphere-hydrology modeling, geostatistical merging of hydrometeorological variables, and quantification of spatio-temporal precipitation variability by attenuation analysis of commercial microwave links.



Susan C. Steele-Dunne (Member, IEEE) received the S.M. and Ph.D. degrees in hydrology from the Massachusetts Institute of Technology, Cambridge, MA, USA, in 2002 and 2006, respectively.

She has been with the Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands, since 2008. She leads the M-WAVE group, who perform research from field to global scales, combining in situ and spaceborne sensors to improve our understanding of microwave interactions with vegetation. Her research interests

include the use of data assimilation, modeling and machine learning to exploit spaceborne radar instruments for applications in ecosystem and agricultural monitoring.