



The scientific value of tractography: accuracy vs usefulness

Francois Rheault¹ · Helen Mayberg² · Michel Thiebaut de Schotten^{3,4} · Alard Roebroeck⁵ · Stephanie J Forkel^{4,6,7}

Received: 27 March 2025 / Accepted: 19 April 2025

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2025

Abstract

Tractography has emerged as a central tool for mapping the cerebral white matter architecture. However, its scientific value continues to be a subject of debate, given its inherent limitations in anatomical accuracy. This concise communication showcases key points of a debate held at the 2024 Tract-Anat Retreat, addressing the trade-offs between the accuracy and utility of tractography. While tractography remains constrained by limitations related to resolution, sensitivity, and validation, its usefulness and utility in areas such as surgical planning, disorder prediction, and the elucidation of brain development are emphasized. These perspectives highlight the necessity of context-specific interpretation, anatomically informed algorithms, and the continuous refinement of tractography workflows to achieve an optimal balance between accuracy and utility.

“Le mieux est le mortel ennemi du bien”.
- Montesquieu's *Pensées* (1726).

This quote, often misattributed to Voltaire, highlights how the pursuit of perfection can sometimes undermine what is already good.

Opening statement

Tractography offers a glimpse into the complex network of connections that enable human cognition and behavior. Despite its promise, tractography faces a formidable challenge: the brain's incredible complexity over all scales. Limitations of diffusion magnetic resonance imaging (dMRI) tractography are inherent to the fact that dMRI is an indirect measurement of the white matter architecture and will always be limited by spatial resolution. However, does it mean that tractography cannot be useful or valuable in specific situations or at a particular scale? At the International Society of Tractography, the debate titled “Neuroanatomy is too complicated to be solved by tractography” led to discussions regarding the strengths and limitations of dMRI tractography and its role in modern neuroscience. This commentary examines the dichotomy between the accuracy and usefulness of tractography, highlighting the importance of anatomically informed tractography reconstruction (Table 1).

Proof tractography is useful

- **Neurosurgical planning:** Tractography helps surgeons identify critical white matter pathways near tumors, enabling them to plan surgical approaches that minimize

✉ Francois Rheault
francois.m.rheault@usherbrooke.ca

¹ Sherbrooke Connectivity Imaging Lab (SCIL), Computer Science Department, Université de Sherbrooke, Sherbrooke, Canada

² Nash Family Center for Advanced Circuit Therapeutics, Icahn School of Medicine at Mount Sinai, Mount Sinai, NY, USA

³ Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives-UMR 5293, CNRS, CEA University of Bordeaux, Bordeaux, France

⁴ Brain Connectivity and Behaviour Laboratory, Paris, France

⁵ Department of Cognitive Neuroscience, Faculty of Psychology & Neuroscience, Maastricht University, Maastricht, The Netherlands

⁶ Donders Institute for Brain Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

⁷ Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands

Table 1 Summary of tractography applications and limitations across Spatial scales. This table highlights how the goals and accuracy of tractography vary across macro-, meso-, and microscale levels of brain analysis. At the **macroscale**, tractography is highly useful for clinical applications such as neurosurgical planning and targeting in deep brain stimulation (DBS) or focused ultrasound (FUS), with reliable identification of major white matter pathways. The **mesoscale** enables functional network modelling and translational insights into neuropsychiatric disorders, with moderate anatomical correspondence to invasive tracer studies. At the **microscale**, tractography contributes to developmental and structural modelling, though its accuracy is limited by the resolution constraints of diffusion MRI and indirect Estimation of axonal pathways

Goal Scale	Useful	Accurate
Macroscale	Assists in neurosurgical planning to avoid critical pathways and guides DBS and FUS for precise targeting.	Identifies major white matter pathways reliably and reproducibly.
Mesoscale	Supports predictions of network functions underlying neuropsychiatric disorders and provides a foundation for personalized therapies.	Provides reasonable correlation with tracer studies in non-human primates for connectivity patterns.
Microscale	Contributes to understanding axonal growth, brain geometry, and developmental processes.	Limited by the resolution of dMRI; indirect approximations of axonal pathways;

damage to essential brain regions (Kamagata et al. 2024; Krieg et al. 2017; Radwan et al. 2024).

- **Deep brain stimulation (DBS):** Tractography aids in understanding the relationship between DBS electrode placement and treatment outcomes, facilitating more precise targeting of brain regions involved in movement disorders and other conditions (Alagapan et al. 2023; Calabrese 2016).
- **Focused Ultrasound (FUS):** This was recently approved for the treatment of refractory essential tremor and similarly to DBS, more precise targeting can be achieved using tractography (Feltrin et al. 2022; Krishna et al. 2019).
- **Cognition and disorder predictions:** Tractography, when combined with complementary datasets, offers a data-driven approach to predicting neurological deficits, such as those seen in stroke or neurodegenerative conditions. This predictive capability supports the development of personalized interventions and enables targeted therapies by linking brain disconnection patterns to cognitive or clinical outcomes (Forkel et al. 2022; Gonzalez-Aguines et al. 2019; Talozzi et al. 2023; Thiebaut de Schotten & Forkel, 2022).
- **Understanding brain geometry and development:** Tractography data can contribute to a better understanding of how axons grow and connect, how brain geometry is influenced by development, and vice versa (Chenot et al. 2019; David et al. 2019; Hau et al. 2016; Maffei et al. 2019; Petit et al. 2022).
- **Correlation with tracer studies:** While some studies noted that correspondence between tractography and tracer is inherently limited (Thomas et al. 2014), several studies have shown a modest to strong correlation between tractography results and findings from invasive tracer studies in non-human primates. The divergence in reports may stem from different studies using different tracer techniques or databases, different dMRI data, and different tractography algorithms. While imperfect, this suggests that tractography can reasonably represent connectivity patterns (Azadbakht et al. 2015; Knösche et al. 2015; Roebroek et al. 2019; Yendiki et al. 2022). However, no complete connectome in non-human primates has been achieved.
- **Correlation with human histology studies:** Several studies have shown a modest to strong correlation between findings from histology techniques with dMRI orientation estimates (Budde and Annesse 2013; Khan and Hashmi 2015; Mollink et al. 2017; Seehaus et al. 2015) and the tractography based on them. Although inferred sensitivity and specificity of tractography are imperfect (Reveley et al. 2015; Roebroek et al. 2008; Thomas et al. 2014) this suggests that tractography can reconstruct macro- and mesoscale connectivity patterns when its validated strengths and limitations are taken into account (Roebroek et al. 2019; Yendiki et al. 2022), as discussed below. Furthermore, its accuracy could be further improved when jointly modeled with microscopy in ex vivo situations (Howard et al. 2019).

Proof tractography is accurate

- **Capable of finding major fiber tracts:** Tractography has consistently demonstrated its ability to identify white matter pathways in the brain at the macroscale. While it may not perfectly replicate the full extent of these tracts, it offers valuable insights into their overall

Tractography is not perfect, but we know its limitations

While different tractography algorithms may yield varying results, the overall tractography process is largely reproducible. This means that repeated scans of the same individual are likely to produce consistent findings, demonstrating a degree of reliability.

- **Sensitivity/Specificity Trade-off:** High sensitivity (detecting all true connections) often comes at the cost of reduced specificity (avoiding false connections) and vice versa. This means that researchers and clinicians must carefully consider the trade-offs and choose tractography parameters that align with their specific objectives (Bastiani et al. 2012; Maier-Hein et al. 2017; Schilling et al. 2019; Seehaus et al. 2013; Thomas et al. 2014).
- **Limited resolution and indirect nature:** The resolution of diffusion MRI data is significantly lower (millimeter scale, typically ranging from 1.5 mm to 2.5 mm) than the microscopic scale of individual axons (micrometer scale. The average axon diameter, excluding the myelin sheath, is typically 1 to 10 micrometers). On top of limited resolution, dMRI only offers indirect measurements for local structures, since its signal emerges from diffusion of water molecules. Consequently, local models, upon which tractography is based, can deviate significantly from accurately representing local axonal orientation. This, coupled with the inferential nature of tractography algorithms, means that the technique may only provide an indirect and potentially incomplete picture of true fiber pathways (Rheault et al. 2020; Schilling et al. 2016, 2018).
- **Over-reliance on local orientation information:** Traditional tractography algorithms primarily rely on local orientation information derived from diffusion MRI. This limited information, without the integration of broader anatomical knowledge or other modalities, contributes to the inaccuracies observed in reconstructions (Bastiani et al. 2012; Daducci et al. 2016; Rheault et al. 2019; Schilling et al. 2022).
- **Limited validation in humans:** While validation studies in non-human primates and phantoms provide valuable insights, the direct validation of tractography in living human brains remains a significant challenge. Complementary post-mortem approaches such as Klingler post-mortem dissections (Akeret et al. 2022; Catani et al. 2012; Thiebaut de Schotten et al. 2011), myelin staining (Seehaus et al. 2015; Thiebaut de Schotten et al. 2011) can help decipher true connections from spurious reconstructions, but the absence of a definitive gold standard for human brain connectivity makes it difficult to

assess the accuracy of tractography findings fully. Additionally, connectivity presents with a significant degree of variability between individuals, even at the macrostructural level (Catani et al. 2007; Croxson et al. 2018; Forkel et al. 2014; 2022) presents an additional level of complexity in the quest for the identification of the “true connectome”. These approaches tackle structural connectivity across regions, an extension of the local tissue structures. For local structure, multiple post-mortem imaging techniques, such as polarized light imaging and polarized optical coherence tomography (OCT), offer high-resolution measurements to validate local models of fiber orientation.

Closing statement

No method is perfect, and the quest to unravel the mysteries of the human connectome is an ongoing endeavor. Despite its limitations, tractography continues to be an invaluable resource. While this technique has undoubtedly advanced our understanding of brain connectivity, the sources underscore the critical need for objective criticism, cautious interpretation, and continuous refinement. As we have seen, the inherent limitations of tractography, stemming from its inferential nature and the complex nature of the multiscale brain organization, can lead to inaccuracies and inconsistencies in reconstructions. Moving forward, a balanced anatomically-validated approach is crucial – one that recognizes the scale of interest and the strengths and weaknesses of tractography to answer neuroanatomical questions at the appropriate scale and ensure clinical use is reliable despite the limited availability of anatomical ground truth.

Author contributions All authors participated equally to the ideas/content, writing and proofreading.

Funding This work was supported by the Donders Mohrmann Fellowship on ‘*Neurovariability*’ No. 2401515 (SJF) and the Dutch Research Council NWO Aspasia Grant ‘*Human individuality: phenotypes, cognition, and brain disorders*’ (SJF).

MTdS was supported by HORIZON-INFRA-2022 SERV (Grant No. 101147319) ‘*EBRAINS 2.0: A Research Infrastructure to Advance Neuroscience and Brain Health*’, by the European Union’s Horizon 2020 research and innovation program under the European Research Council (ERC) Consolidator grant agreement No. 818521 (DISCONNECTOME), the University of Bordeaux’s IdEx ‘*Investments for the Future*’ program RRI ‘*IMPACT*’, and the IHU ‘*Precision & Global Vascular Brain Health Institute – VBHI*’ funded by the France 2030 initiative (ANR-23-IAHU-0001).

HM was supported through NIH funding (UH3NS103550), Wellcome Leap MCPsych, Hope for Depression Research Foundation.

Data availability No datasets were generated or analysed during the current study.

Declarations

Competing interests The authors declare no competing interests.

References

- Akeret K, Forkel SJ, Buzzi RM, Vasella F, Amrein I, Colacicco G, Serra C, Krayenbühl N (2022) Multimodal anatomy of the human fornical commissure. *Commun Biology* 5(1):742
- Alagapan S, Choi KS, Heisig S, Riva-Posse P, Crowell A, Tiruvadi V, Obatusin M, Veerakumar A, Waters AC, Gross RE (2023) & others. Cingulate dynamics track depression recovery with deep brain stimulation. *Nature*, 622(7981), 130–138
- Azadbakht H, Parkes LM, Haroon HA, Augath M, Logothetis NK, de Crespigny A, D'Arceuil HE, Parker GJ (2015) Validation of high-resolution tractography against in vivo tracing in the macaque visual cortex. *Cereb Cortex* 25(11):4299–4309
- Bastiani M, Shah NJ, Goebel R, Roebroeck A (2012) Human cortical connectome reconstruction from diffusion weighted MRI: the effect of tractography algorithm. *NeuroImage* 62(3):1732–1749
- Budde MD, Annesse J (2013) Quantification of anisotropy and fiber orientation in human brain histological sections. *Front Integr Neurosci* 7:3
- Calabrese E (2016) Diffusion tractography in deep brain stimulation surgery: A review. *Front Neuroanat* 10:45
- Catani M, Allin MP, Husain M, Pugliese L, Mesulam MM, Murray RM, Jones DK (2007) Symmetries in human brain language pathways correlate with verbal recall. *Proceedings of the National Academy of Sciences*, 104(43), 17163–17168
- Catani M, Dell'Acqua F, Vergani F, Malik F, Hodge H, Roy P, Valabregue R, De Schotten MT (2012) Short frontal lobe connections of the human brain. *Cortex* 48(2):273–291
- Chenot Q, Tzourio-Mazoyer N, Rheault F, Descoteaux M, Crivello F, Zago L, Mellet E, Jobard G, Joliot M, Mazoyer B & others. (2019) A population-based atlas of the human pyramidal tract in 410 healthy participants. *Brain Struct Funct* 224(2):599–612
- Croxson PL, Forkel SJ, Cerliani L, de Thiebaut M (2018) Structural variability across the primate brain: A cross-species comparison. *Cereb Cortex* 28(11):3829–3841
- Daducci A, Dal Palú A, Descoteaux M, Thiran J-P (2016) Microstructure informed tractography: pitfalls and open challenges. *Front Neurosci* 10:247
- David S, Heemskerk AM, Corrivetti F, Thiebaut de Schotten M, Sarubbo S, Corsini F, De Benedictis A, Petit L, Viergever MA, Jones DK (2019) & others. The superoanterior fasciculus (SAF): A novel white matter pathway in the human brain? *Frontiers in Neuroanatomy*, 13, 24
- de Thiebaut M, Forkel SJ (2022) The emergent properties of the connected brain. *Science* 378(6619):505–510
- Feltrin FS, Chopra R, Pouratian N, Elkurd M, El-Nazer R, Lanford L, Dauer W, Shah BR (2022) Focused ultrasound using a novel targeting method four-tract tractography for magnetic resonance-guided high-intensity focused ultrasound targeting. *Brain Commun* 4(6):fcac273
- Forkel SJ, de Schotten MT, Kawadler JM, Dell'Acqua F, Danek A, Catani M (2014) The anatomy of fronto-occipital connections from early blunt dissections to contemporary tractography. *Cortex* 56:73–84
- Forkel SJ, Friedrich P, Thiebaut de Schotten M, Howells H (2022) White matter variability, cognition, and disorders: A systematic review. *Brain Struct Funct*, 1–16
- Gonzalez-Aguines A, Moreno-Andrade T, Gongora-Rivera F, Cordero-Perez AC, Ortiz-Jiménez X, Cavazos-Luna O, Garza-Villareal E, Campos-Coy M, Elizondo-Riojas G (2019) The role of tractography in ischemic stroke: A review of the literature. *Med Univ* 20:161–165
- Hau J, Sarubbo S, Perchey G, Crivello F, Zago L, Mellet E, Jobard G, Joliot M, Mazoyer BM, Tzourio-Mazoyer N (2016) & others. Cortical terminations of the inferior fronto-occipital and uncinate fasciculi: Anatomical stem-based virtual dissection. *Frontiers in Neuroanatomy*, 10, 58
- Howard AF, Mollink J, Kleinnijenhuis M, Pallegage-Gamarallage M, Bastiani M, Cottaar M, Miller KL, Jbabdi S (2019) Joint modeling of diffusion MRI and microscopy. *NeuroImage* 201:116014
- Kamagata K, Andica C, Uchida W, Takabayashi K, Saito Y, Lukies M, Hagiwara A, Fujita S, Akashi T, Wada A, others (2024) Advancements in diffusion MRI tractography for neurosurgery. *Invest Radiol* 59(1):13–25
- Khan S, Hashmi GS (2015) Histology and functions of connective tissues: A review Article. *Univ J Dent Sci* 1:28–34
- Knösche TR, Anwander A, Liptrot M, Dyrby TB (2015) Validation of tractography: comparison with manganese tracing. *Hum Brain Mapp* 36(10):4116–4134
- Krieg SM, Lioumis P, Mäkelä JP, Wilenius J, Karhu J, Hannula H, Savolainen P, Lucas CW, Seidel K, Laakso A (2017) & others. Protocol for motor and language mapping by navigated TMS in patients and healthy volunteers; workshop report. *Acta Neurochirurgica*, 159, 1187–1195
- Krishna V, Sammartino F, Agrawal P, Changizi BK, Bourekas E, Knopp MV, Rezaei A (2019) Prospective tractography-based targeting for improved safety of focused ultrasound thalamotomy. *Neurosurgery* 84(1):160–168
- Maffei C, Sarubbo S, Jovicich J (2019) Diffusion-based tractography atlas of the human acoustic radiation. *Sci Rep* 9(1):4046
- Maier-Hein KH, Neher PF, Houde J-C, Côté M-A, Garyfallidis E, Zhong J, Chamberland M, Yeh F-C, Lin Y-C, Ji Q (2017) & others. The challenge of mapping the human connectome based on diffusion tractography. *Nature Communications*, 8(1), 1349
- Mollink J, Kleinnijenhuis M, van Walsum A-M, van Sotiropoulos C, Cottaar SN, Mirfin M, Heinrich C, Jenkinson MP, Pallegage-Gamarallage M, Ansoorge M (2017) O., & others. Evaluating fibre orientation dispersion in white matter: Comparison of diffusion MRI, histology and polarized light imaging. *Neuroimage*, 157, 561–574
- Petit L, Ali KM, Rheault F, Boré A, Corsini F, De Benedictis A, Descoteaux M, Sarubbo S (2022) *The Structural Connectivity of The Human Angular Gyrus as Revealed by Microdissection and Diffusion Tractography*
- Radwan AM, Emsell L, Vansteelandt K, Cleeren E, Peeters R, De Vleeschouwer S, Theys T, Dupont P, Sunaert S (2024) Comparative validation of automated presurgical tractography based on constrained spherical Deconvolution and diffusion tensor imaging with direct electrical stimulation. *Hum Brain Mapp* 45(6):e26662
- Reveley C, Seth AK, Pierpaoli C, Silva AC, Yu D, Saunders RC, Leopold DA, Frank QY (2015) Superficial white matter fiber systems impede detection of long-range cortical connections in diffusion MR tractography. *Proc Natl Acad Sci* 112(21):E2820–E2828
- Rheault F, St-Onge E, Sidhu J, Maier-Hein K, Tzourio-Mazoyer N, Petit L, Descoteaux M (2019) Bundle-specific tractography with incorporated anatomical and orientational priors. *NeuroImage* 186:382–398
- Rheault F, Poulin P, Caron AV, St-Onge E, Descoteaux M (2020) Common misconceptions, hidden biases and modern challenges of dMRI tractography. *J Neural Eng* 17(1):011001
- Roebroeck A, Galuske R, Formisano E, Chiry O, Bratzke H, Ronen I, Kim D, Goebel R (2008) High-resolution diffusion tensor imaging and tractography of the human optic Chiasm at 9.4 T. *NeuroImage* 39(1):157–168

- Roebroek A, Miller KL, Aggarwal M (2019) Ex vivo diffusion MRI of the human brain: technical challenges and recent advances. *NMR Biomed*, 32(4), e3941
- Schilling K, Janve V, Gao Y, Stepniewska I, Landman BA, Anderson AW (2016) Comparison of 3D orientation distribution functions measured with confocal microscopy and diffusion MRI. *NeuroImage* 129:185–197
- Schilling K, Gao Y, Janve V, Stepniewska I, Landman BA, Anderson AW (2018) Confirmation of a gyral bias in diffusion MRI fiber tractography. *Hum Brain Mapp* 39(3):1449–1466
- Schilling KG, Nath V, Hansen C, Parvathaneni P, Blaber J, Gao Y, Neher P, Aydogan DB, Shi Y, Ocampo-Pineda M (2019) & others. Limits to anatomical accuracy of diffusion tractography using modern approaches. *Neuroimage*, 185, 1–11
- Schilling KG, Petit L, Rheault F, Remedios S, Pierpaoli C, Anderson AW, Landman BA, Descoteaux M (2020) Brain connections derived from diffusion MRI tractography can be highly anatomically accurate—If we know where white matter pathways start, where they end, and where they do not go. *Brain Struct Funct* 225:2387–2402
- Schilling KG, Tax CM, Rheault F, Landman BA, Anderson AW, Descoteaux M, Petit L (2022) Prevalence of white matter pathways coming into a single white matter voxel orientation: the bottleneck issue in tractography. *Hum Brain Mapp* 43(4):1196–1213
- Seehaus AK, Roebroek A, Chiry O, Kim D-S, Ronen I, Bratzke H, Goebel R, Galuske RA (2013) Histological validation of DW-MRI tractography in human postmortem tissue. *Cereb Cortex* 23(2):442–450
- Seehaus A, Roebroek A, Bastiani M, Fonseca L, Bratzke H, Lori N, Vilanova A, Goebel R, Galuske R (2015) Histological validation of high-resolution DTI in human post mortem tissue. *Front Neuroanat* 9:98
- Talozzi L, Forkel SJ, Pacella V, Nozais V, Allart E, Piscicelli C, Pérenou D, Tranel D, Boes A, Corbetta M, others (2023) Latent disconnectome prediction of long-term cognitive-behavioural symptoms in stroke. *Brain* 146(5):1963–1978
- Thiebaut de Schotten M, Dell'Acqua F, Forkel S, Simmons A, Vergani F, Murphy DG, Catani M (2011a) A lateralized brain network for visuo-spatial attention. *Nat Precedings*, 1–1
- Thiebaut de Schotten M, Ffytche D, Bizzi A, Dell'Acqua F, Allin M, Walshe M, Murray R, Williams S, Murphy D, Catani M (2011b) Atlasing location, asymmetry and inter-subject variability of white matter tracts in the human brain with MR diffusion tractography. *NeuroImage* 54:49e59
- Thomas C, Ye FQ, Irfanoglu MO, Modi P, Saleem KS, Leopold DA, Pierpaoli C (2014) Anatomical accuracy of brain connections derived from diffusion MRI tractography is inherently limited. *Proc Natl Acad Sci* 111(46):16574–16579
- Yeh F-C, Vettel JM, Singh A, Poczos B, Grafton ST, Erickson KI, Tseng W-YI, Verstynen TD (2016) Quantifying differences and similarities in whole-brain white matter architecture using local connectome fingerprints. *PLoS Comput Biol* 12(11):e1005203
- Yendiki A, Aggarwal M, Axer M, Howard AF, van Walsum A-M C., Haber SN (2022) Post mortem mapping of connectonal anatomy for the validation of diffusion MRI. *NeuroImage* 256:119146

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.