

Initial-state preparation with dynamically generated system-environment correlations

C. H. Fleming,¹ Albert Roura,² and B. L. Hu¹¹*Joint Quantum Institute and Department of Physics, University of Maryland, College Park, Maryland 20742, USA*²*Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, D-14476 Golm, Germany*

(Received 26 May 2011; published 4 August 2011)

The dependence of the dynamics of open quantum systems upon initial correlations between the system and environment is an utterly important yet poorly understood subject. For technical convenience most prior studies assume factorizable initial states where the system and its environments are uncorrelated, but these conditions are not very realistic and give rise to peculiar behaviors. One distinct feature is the rapid buildup or a sudden jolt of physical quantities immediately after the system is brought in contact with its environments. The ultimate cause of this is an initial imbalance between system-environment correlations and coupling. In this paper we demonstrate explicitly how to avoid these unphysical behaviors by proper adjustments of correlations and/or the coupling, for setups of both theoretical and experimental interest. We provide simple analytical results in terms of quantities that appear in linear (as opposed to affine) master equations derived for factorized initial states.

DOI: [10.1103/PhysRevE.84.021106](https://doi.org/10.1103/PhysRevE.84.021106)

PACS number(s): 05.30.-d, 03.65.Yz, 05.10.Gg, 05.70.Ln

I. OPEN-SYSTEM INITIAL CORRELATIONS

An open quantum system is a quantum system S that interacts with some environment E whose degrees of freedom have been coarse grained (colloquially, traced out or integrated over). The unitary evolution of the combined system plus environment, C (for combined or closed), is generated by the Hamiltonian

$$\mathbf{H}_C \equiv \mathbf{H}_S + \mathbf{H}_E + \mathbf{H}_I, \quad (1)$$

where \mathbf{H}_I denotes the system-environment interaction. Specifying the initial state of the combined system is necessary to determine the open-system dynamics. The most common choice is to assume factorized initial states for the system and environment

$$\rho_C(0) = \rho_S(0) \otimes \rho_E(0), \quad (2)$$

$$\rho_E(0) = \frac{1}{Z_E(\beta)} e^{-\beta \mathbf{H}_E}, \quad (3)$$

where $Z_E(\beta)$ denotes the partition function of the free (noninteracting) environment and $T = 1/\beta$ is the temperature of the environment, which acts here as a thermal reservoir.

When considering environments with a large number of high-frequency modes and characterized by an UV frequency cutoff Λ , such a factorized initial state (chosen for mathematical simplicity) unfortunately engenders unphysical behavior such as a sudden jolt in physical quantities near the initial time (this was analyzed in some detail in Ref. [1]) or spurious cutoff sensitivity of certain system correlators (see Appendix D in Ref. [2]) [3]. This kind of initial condition assumes that an uncorrelated system and environment are instantaneously coupled with nonvanishing strength. The pathological behavior arises because the factorized initial state contains a number of highly excited energy states of the full Hamiltonian (including \mathbf{H}_I), even when the initial reduced states of the system and environment are not highly excited in the free theory, and it is a reflection of the high-frequency modes of the environment quickly becoming correlated with the system within a time of order $1/\Lambda$.

The next most common choice of initial state (see Ref. [5] and references therein) has been to consider system deforma-

tions or measurements of the global equilibrium state of the combined system C , with density matrix

$$\rho_C(0) = \sum_n \mathbf{O}'_n \frac{1}{Z_C(\beta)} e^{-\beta \mathbf{H}_C} \mathbf{O}_n, \quad (4)$$

where the operators \mathbf{O} and \mathbf{O}' are restricted to act on the system. However, this method still gives rise to jolts for sufficiently general deformations or measurements [6], which can be understood as a consequence of altering the state of the system instantaneously [7].

To cure or tame these drastic effects, especially in the context of linear systems, the following procedure has been suggested: (a) force the system by a constant amount, (b) wait for it to relax into the displaced equilibrium state, and then (c) release the force [5]. Alternatively, in order to generate interesting coherent superposition states for the system, one can start with the equilibrium state of the combined system and act on the system, but for a nonvanishing time [7]. Essentially we view the problem as an imbalance between initial correlations and initial coupling strength; the imbalance can be countered on either side. We also believe that the most natural resolution should be a dynamical preparation that relies upon equilibration [5,8] followed by an additional preparation of the system for a finite time [7]. Our key contribution is showing that this can be achieved while still taking advantage of the simpler analytical results obtained when deriving the master equation for a factorized initial state, without the need to introduce inhomogeneous terms and an affine master equation [8].

In the following section we will briefly discuss the perturbative open-system master equation that we will use to approach these issues. Then in Secs. III and IV we will provide resolutions based, respectively, on balancing the coupling and the correlations.

II. OPEN-SYSTEM DYNAMICS

In the time-local representation (also called the convolutionless or Markovian representation), the equation governing

the dynamics of the reduced density matrix of the system ρ can be expressed as a quantum Liouville equation

$$\frac{d}{dt}\rho(t) = \mathcal{L}(t)\rho(t) \quad (5)$$

for any factorized initial condition. As a perturbative approximation, $\mathcal{L}(t)$ is expanded in powers of the system-environment interaction $\mathbf{H}_I(t)$ and truncated to some order. (In this section we momentarily consider time-dependent Hamiltonians, even in the Schrödinger picture, since the more general relations will be necessary for our techniques.) Such perturbative master equations can be derived in a variety of ways [9–11] and find application in many branches of physics and chemistry [12–15]. The expansion of $\mathcal{L}(t)$ will then take the form

$$\mathcal{L}(t) = \sum_{k=0}^{\infty} \mathcal{L}_{2k}(t), \quad (6)$$

$$\mathcal{L}_0\rho = [-i\mathbf{H}_S(t), \rho], \quad (7)$$

where $\mathcal{L}_{2k} = O(\mathbf{H}_I^{2k})$ and to zeroth order the system is driven in a unitary manner by its Hamiltonian $\mathbf{H}_S(t)$. We take the expansion to be even as we only consider Gaussian noise with vanishing mean [16]. A Gaussian noise distribution is necessary for higher-order perturbation theory to be nonsecular in the late-time limit [17], though at second order one is effectively truncating the noise cumulants in a manner consistent with Gaussian noise.

The linear master equation is derived under the general assumptions of a factorized initial state and an expansion of the interaction as a sum of separable operators:

$$\mathbf{H}_I(t) = \sum_n \mathbf{L}_n(t) \otimes \mathbf{I}_n(t), \quad (8)$$

where $\mathbf{L}_n(t)$ and $\mathbf{I}_n(t)$ are (possibly time-dependent) system and environment operators, respectively, in the Schrödinger picture. The environment coupling operators $\mathbf{I}_n(t)$ will typically be collective observables of the environment, dependent upon many modes.

Using the notation of Ref. [17], the second-order master equation can be expressed as

$$\mathcal{L}_2\{\rho\} \equiv \sum_{n,m} [\mathbf{L}_n, \rho(\mathbf{A}_{nm} \diamond \mathbf{L}_m)^\dagger - (\mathbf{A}_{nm} \diamond \mathbf{L}_m)\rho], \quad (9)$$

where the operators \mathbf{A} and diamond product define the second-order operator

$$(\mathbf{A}_{nm} \diamond \mathbf{L}_m)(t) \equiv \int_0^t d\tau \alpha_{nm}(t, \tau) \{\mathcal{G}_0(t, \tau) \mathbf{L}_m(\tau)\}. \quad (10)$$

Here $\mathcal{G}_0(t, \tau): \rho(\tau) \rightarrow \rho(t)$ is the free-system propagator, which for a constant Hamiltonian \mathbf{H} is given by

$$\mathcal{G}_0(t, \tau)\rho = e^{-i(t-\tau)\mathbf{H}}\rho e^{+i(t-\tau)\mathbf{H}}, \quad (11)$$

while $\alpha_{nm}(t, \tau)$ are the environment correlation functions defined by

$$\alpha_{nm}(t, \tau) = \langle \mathbf{I}_n(t) \mathbf{I}_m(\tau) \rangle_E, \quad (12)$$

where $\mathbf{I}_n(t)$ represents the time-evolving \mathbf{I}_n in the interaction (Dirac) picture. In general the correlation function is Hermitian and positive definite. For constant coupling to any stationary environment, the correlation function will also

be stationary $\alpha(t, \tau) = \alpha(t - \tau)$. Furthermore, for a thermal environment the correlation function will satisfy the Kubo-Martin-Schwinger relation [18,19]

$$\tilde{\alpha}(\omega) = \tilde{\alpha}^*(-\omega)e^{-\beta\omega}, \quad (13)$$

where $\tilde{\alpha}(\omega) = \int_{-\infty}^{+\infty} dt \alpha(t) e^{-i\omega t}$ denotes the Fourier transform.

From this perspective, the mathematical cause of the initial jolt becomes clear. For constant Hamiltonians and an initially stationary environment, the second-order operator obeys the relation

$$\frac{d}{dt}(\mathbf{A}_{nm} \diamond \mathbf{L}_m)(t) = \alpha_{nm}(t) \{\mathcal{G}_0(t) \mathbf{L}_m\}, \quad (14)$$

which can be extremely large near the initial time when considering an environment with a sufficient amount of high-frequency modes (such as low-temperature ohmic and supraohmic environments) since $\alpha(t)$ is typically a very localized distribution in those cases. E.g., for a finite but large cutoff Λ , the environment correlation function for a low-temperature ohmic environment becomes of order Λ for a time of order $1/\Lambda$.

III. COUPLING SWITCH-ON

One method for balancing the initial coupling between the system and environment with their initial lack of correlation is to turn on the coupling slowly with a time-dependent interaction such as

$$\mathbf{H}_I = \theta_s(t) \sum_n \mathbf{L}_n \otimes \mathbf{I}_n, \quad (15)$$

where $\theta_s(t): [0, \infty) \rightarrow [0, 1)$ is a smooth switch-on function with a characteristic time scale τ_s , which vanishes at the initial time and becomes (effectively) one for times longer than τ_s . To some extent, this was considered for linear systems in Ref. [4].

Such a time-dependent interaction is equivalent to employing the second-order operator

$$(\mathbf{A}_{nm} \diamond \mathbf{L}_m)(t) = \theta_s(t) \int_0^t d\tau \theta_s(t - \tau) \alpha_{nm}(\tau) \{\mathcal{G}_0(\tau) \mathbf{L}_m\} \quad (16)$$

for otherwise constant couplings and Hamiltonians. Therefore, any initial jolt due to the localized nature of $\alpha(t)$ will be suppressed by $\theta_s(t)$ as long as $\tau_s \gg 1/\Lambda$.

As can be seen in Fig. 1, the cutoff-frequency jolts are essentially replaced by jolts of frequency $\min[\Lambda, 1/\tau_s]$ and amplitude proportional to the same value. This approach provides a useful way of generating initial system-environment correlations when τ_s is much larger than $1/\Lambda$ but smaller than any other relevant time scales (such as the system frequencies). Furthermore, even if a mild jolt is still present, the important point is that it is cutoff insensitive (for fixed τ_s and sufficiently large Λ).

IV. DYNAMICALLY PREPARED INITIAL STATES

Alternatively, in order to balance the initial correlations with an initially nonvanishing interaction strength, we will consider here initial states with suitable correlations to the




FIG. 1. (Color online) Zero-temperature ohmic decay rate for the instantaneously coupled (red thick line) and gradually coupled (blue thin line) initial states of a two-level system with exponential cutoff frequency $\Lambda = 100\Omega$. In this case the switch-on function is exponential $\theta_s(t) = 1 - e^{-t/\tau_s}$ and the switch-on times τ_s are chosen to take the values $1/\Lambda$, $2/\Lambda$, $4/\Lambda$, $8/\Lambda$, and $16/\Lambda$.

environment. Such states will be obtained via an auxiliary construction that involves evolving an initially uncorrelated state for a sufficiently long time (a similar procedure was used in Refs. [20,21] within the context of semiclassical gravity). The system-environment correlations are then dynamically generated through the environment interaction itself. Our first examples of equilibrium preparation will be the simplest mathematically, while the final examples of nonequilibrium preparation will be closer to actual laboratory experiments.

In all cases we will take the system and environment to be uncorrelated not at $t = 0$ but in the infinite past,

$$\rho_C(-\infty) = \rho_S(-\infty) \otimes \rho_E(-\infty), \quad (17)$$

for some (possibly unimportant) system state $\rho_S(-\infty)$ and thermal $\rho_E(-\infty)$. We then define the system Hamiltonian piecewise in time

$$\mathbf{H}_S(t) = \begin{cases} \mathbf{H}_+(t), & 0 < t \\ \mathbf{H}_-, & t < 0 \end{cases} \quad (18)$$

such that in the past the system is allowed to equilibrate with the environment for an infinite time, which determines the correlated initial state at $t = 0$. The second-order master equation is then determined by

$$(\mathbf{A}_{nm} \diamond \mathbf{L}_m)(t) = \int_{-\infty}^t d\tau \alpha_{nm}(t, \tau) \{\mathcal{G}_S(t, \tau) \mathbf{L}_m(\tau)\}. \quad (19)$$

To analyze the coefficients associated with the initially correlated state we will reduce them to a sum of coefficients for the auxiliary initially uncorrelated state involving various time ranges. First, we split the integration into two parts

$$\int_{-\infty}^t d\tau = \int_0^t d\tau + \int_{-\infty}^0 d\tau, \quad (20)$$

with the first integral depending only upon $\mathcal{G}_+(t, \tau)$ and corresponding to the uncorrelated coefficients. Inserting the

product $\mathcal{G}_-(0, t)\mathcal{G}_-(t, 0)$, which equals the identity, the second integral can be written as

$$\mathcal{M}(t) \int_{-\infty}^0 d\tau \alpha_{nm}(t, \tau) \{\mathcal{G}_-(t, \tau) \mathbf{L}_m(\tau)\}, \quad (21)$$

given the operator

$$\mathcal{M}(t) \equiv \mathcal{G}_+(t, 0)\mathcal{G}_-(0, t). \quad (22)$$

The integral in Eq. (21) is then broken up into two parts

$$\int_{-\infty}^0 d\tau = \int_{-\infty}^t d\tau - \int_0^t d\tau, \quad (23)$$

corresponding to the asymptotic and finite-time coefficients for an initially uncorrelated system driven by the time-independent preparation Hamiltonian \mathbf{H}_- . Finally, our correlated coefficients can be expressed in terms of the uncorrelated coefficients as

$$\underbrace{(\mathbf{A}_{nm} \diamond \mathbf{L}_m)(t)}_{\text{correlated}} = \underbrace{(\mathbf{A}_{nm} \diamond \mathbf{L}_m)_+(t)}_{\text{uncorrelated}} - \underbrace{\mathcal{M}(t) \{ (\mathbf{A}_{nm} \diamond \mathbf{L}_m)_-(t) - (\mathbf{A}_{nm} \diamond \mathbf{L}_m)_-(\infty) \}}_{\substack{\text{jolt suppression} \\ \text{preparation eraser}}}, \quad (24)$$

where the subscripted $(\mathbf{A} \diamond \mathbf{L})_{\pm}$ coefficients are defined as

$$(\mathbf{A}_{nm} \diamond \mathbf{L}_m)_{\pm}(t) \equiv \int_0^t d\tau \alpha_{nm}(t, \tau) \{\mathcal{G}_{\pm}(t, \tau) \mathbf{L}_m(\tau)\}. \quad (25)$$

If the system frequencies are always small compared to the cutoff, we can inspect the early-time behavior (and jolts) by taking $\mathcal{G}_{\pm}(t) \approx \mathbf{1}$. Then one can see that the first two terms of Eq. (24) will precisely cancel in the early-time regime. Therefore, the correlated initial states are jolt-free given sufficiently small system frequencies compared to the cutoff: $\Omega \ll \Lambda$. The final term in turn is such that in the late-time limit it precisely cancels the second term and erases all memory of \mathbf{H}_- . Finally note that, quite trivially, if we choose $\mathbf{H}_+(t) = \mathbf{H}_-$, then the first two terms cancel and we recover the equilibrium coefficients at any finite time.

A. Equilibrium preparation

To prepare an initial state in this approach, we choose the past Hamiltonian \mathbf{H}_- such that its dynamics along with the environment interaction relaxes our system to the desired initial state:

$$\lim_{t \rightarrow \infty} e^{t\mathcal{L}_-(\infty)} \rho_0 = \rho_0, \quad (26)$$

$$\mathcal{L}_-(\infty) \rho_0 = \mathbf{0}, \quad (27)$$

where $\mathcal{L}_-(\infty)$ is the stationary limit of the Liouvillian for a system with the past Hamiltonian as well as the coupling to the environment. Our target state ρ_0 will only be specified to zeroth order in the system-environment interaction. This is because for sufficiently long times (and in particular for the asymptotic equilibrium state) the diagonal elements of the reduced density matrix in the energy basis cannot be determined beyond zeroth order anyway when using the second-order perturbative master equation [22]. Due to unavoidable degeneracy present in all open-system dynamics, one actually requires components of the fourth-order master equation to calculate the full second-order solutions. The second-order master equation provides

for all second-order dynamical quantities, such as frequency shifts, dissipation, diffusion, and decoherence rates. We are concerned here with the induced jolts, which are dynamical quantities, and so this subtle point does not raise any additional problems for us.

1. Preparation by decoherence

For \mathbf{L}_n all commuting with each other, one can force a general environment into ℓ -state preparation via decoherence. If the past Hamiltonian is deactivated or, more generally, taken to commute with \mathbf{L}_n , then since all system operators commute with each other, the master equation and its solutions will trivially result in a system that decoheres in the ℓ basis associated with the \mathbf{L}_n . Thus coefficients prepared in this manner are consistent with any initial state that is a completely incoherent mixture of ℓ states. [Note that if $\rho_S(-\infty)$ corresponds to a pure eigenstate of the set $\{\mathbf{L}_n\}$, this procedure simply adjusts the state of the environment, while the system and environment remain unentangled.]

2. Preparation by equilibration

A finite-temperature environment allows mixed-state preparation by equilibration. Essentially one chooses the past Hamiltonian so that its thermal state (or some other steady state) is the desired initial state. For a positive-temperature environment, at zeroth order one can prepare a (sufficiently) mixed state ρ_0 with the past Hamiltonian $\mathbf{H}_- = -T \ln(\rho_0)$. However, one must be careful that past system frequencies are small compared to the high-frequency jolts; otherwise, this preparation will fail to remedy jolting. One can work out that the adiabatic preparation regime is given by

$$\frac{p_{\max}}{p_{\min}} \ll e^{\beta\Lambda}, \quad (28)$$

where Λ is the jolt frequency and p are the initial-state probabilities of preparation energy levels connected by \mathbf{L}_n . (Clearly, for this method to work there can only be a finite number of such energy levels.)

3. Preparation by freezing

To prepare an initially pure state via equilibration at the order that we are working, one requires a zero-temperature environment for preparation by freezing (see Fig. 2). Then one can choose any \mathbf{H}_- with ground state ρ_0 . It is important to emphasize that the reduced density matrix of the system corresponding to the ground state of the combined system will not be a pure state in general due to the entanglement between the system and the environment: The free ground state of the system is a pure state, but the reduced density matrix of the open system is in general a mixed state beyond zeroth order in the system-environment coupling. However, this point becomes irrelevant at the order that we are working since, as explained above, when using the second-order perturbative master equation to prepare the initial state by equilibration, one cannot meaningfully specify ρ_0 beyond zeroth order.




FIG. 2. (Color online) Zero-temperature, ohmic decay rate for the unprepared (red thick line) and prepared (blue thin line) initial states of a two-level system with exponential cutoff frequency $\Lambda = 100\Omega$. In this case preparation by freezing was used to create an initially excited state.

B. Nonequilibrium preparation

In order to consider situations closer to actual laboratory experiments, here we will first allow the system to equilibrate with the environment (as described in Sec. IV A) and then choose some preparation Hamiltonian $\mathbf{H}_P(t)$, which would (in the absence of coupling to the environment) generate the desired initial state in some finite time τ_P . One simply applies the master-equation coefficients in Eq. (24) with the future Hamiltonian

$$\mathbf{H}_+(t) = \begin{cases} \mathbf{H}_0(t), & \tau_P < t \\ \mathbf{H}_P(t), & t < \tau_P, \end{cases} \quad (29)$$

where $\mathbf{H}_0(t)$ is the desired postpreparation Hamiltonian. All jolts will be avoided if $1/\tau_P \ll \Lambda$: The introduction of a nonvanishing preparation time serves to tame the jolts and eliminate their high-cutoff sensitivity.

1. State flipping

A possible preparation Hamiltonian, which could model as a particular case Rabi oscillations induced by an appropriate laser field acting on a two-level system, is the following:

$$\mathbf{H}_P = \frac{\pi}{2\tau_P} (|\psi_0\rangle\langle 0| + |0\rangle\langle\psi_0|). \quad (30)$$

Assuming that one has a zero-temperature environment and that the system is already equilibrated, driving the system with this Hamiltonian for a time τ_P provides a relatively easy way of preparing an initial pure state $|\psi_0\rangle$. As discussed above, the reduced density matrix of the system at quadratic order and beyond would actually be a mixed state in general because of the system-environment entanglement of the equilibrium state as well as the interaction to the environment while evolving the combined system during this additional finite preparation time. In fact, the preparation time τ_P cannot be too long if we want the state of the system to be more or less close to $|\psi_0\rangle$.

2. State swapping

Let us consider a system that is initially equilibrated without making any assumption as to the temperature of the environment. We couple the system to an ancillary and analog system (equivalent Hilbert spaces) that is already prepared in the desired initial state. The system of interest and ancilla are temporarily coupled in such a way that they swap states, for instance, by means of the following block-matrix preparation Hamiltonian:

$$\mathbf{H}_P = \frac{\pi}{2\tau_P} \begin{bmatrix} \mathbf{0} & \mathbf{1} \\ \mathbf{1} & \mathbf{0} \end{bmatrix}. \quad (31)$$

In the absence of coupling to the environment this would exactly swap the system and ancilla states in a time τ_P . The same remarks as for state flipping concerning the purity and accuracy of the prepared state when taking into account the coupling to the environment also apply in this case.

3. Other possibilities

Within the second-order perturbative approach, the generation of equilibrium correlations in a laboratory setting can always be calculated using Eq. (24). One only needs to make sure that any additional state preparation does not rely upon large system energies as compared to the bath cutoff. For instance, one can consider the preparation of Ref. [7], which relies on ancillary degrees of freedom to drive the equilibrium state into a coherent superposition. In fact, one could simply apply their own time-dependent Hamiltonian to our formulas as $\mathbf{H}_+(t)$ and obtain results consistent with theirs.

V. DISCUSSION

In this paper we have provided a technically and computationally simple method for generating properly correlated initial states of open quantum systems, which is an important issue when the environment contains a large number of high-frequency degrees of freedom. The need for considering appropriately correlated states is because factorized initial states (with uncorrelated system and environment) typically lead to unphysical pathological results at very short times whose effect on certain properties of the system can persist at much longer times. They are a consequence of the high-frequency modes of the environment suddenly becoming correlated with the system due to the interaction Hamiltonian \mathbf{H}_I . (In fact, while being well behaved with respect to the free Hamiltonian, factorized initial states are highly excited states of the full Hamiltonian.)

In previous studies dealing with this issue an appropriately correlated state was obtained by considering the equilibrium state for the combined system (including the system-environment interaction), which is usually done employing Euclidean path integrals. Other states with a more interesting dynamics have been generated by starting with such an equilibrium state and acting on the system S to prepare nontrivial states (including those involving interesting coherent superpositions). Carrying out this preparation procedure instantaneously [5] will still lead in general to a pathological behavior [6], which can be avoided by performing

the preparation in a finite time [7]. Unfortunately in these cases (as well as in approaches based on affine master equations [8]) the formalism is rather involved, especially compared to the much simpler one for factorized initial states. The salient merit of our method is that the results can be written in terms of the same kind of objects that appear in the study of factorized initial states; see Eq. (24).

The key element of our approach is the use of an auxiliary construction that involves evolving the combined system for an infinite time so that it reaches a stationary (equilibrium) state with the system and the environment properly correlated. This global equilibrium state is then taken as the initial state and is further evolved with the full system plus environment Hamiltonian. However, the system Hamiltonian $\mathbf{H}_S(t)$ used for this subsequent evolution can differ from that employed in the generation of the initial state and even be time dependent, so that nontrivial states can be produced in a second preparation stage. With our formulation for the perturbative master equation it is easy to show that the result can be entirely written as a combination of the different objects that appear for factorized initial states, associated with the auxiliary Hamiltonian (for both finite and infinite times) on the one hand and with the Hamiltonian during the actual evolution (for a finite time) on the other. Furthermore, there is a cancellation between the contributions corresponding to the auxiliary and actual Hamiltonians involving finite-time evolutions, which separately exhibit the jolts and unphysical features characteristic of factorized initial states.

It should be emphasized that the crucial role in the suppression of jolts and related pathologies is played by the interaction Hamiltonian, which establishes the appropriate correlations between the fast environment degrees of freedom (high-frequency modes) and the system. In contrast, this is fairly insensitive to the details of the system Hamiltonian as long as its characteristic frequency scales are much smaller than those of the environment high-frequency modes. We have provided a number of explicit examples illustrating how to prepare nontrivial states for the system making use of suitable time-dependent preparation Hamiltonians for the system that last for a finite time. Other similar preparation schemes, such as that of Ref. [7], can be straightforwardly implemented within our approach. In this manner, our results might be applied toward specific experiments or other theoretical formalisms, such as stochastic preparation schemes in quantum process tomography [23,24]. Initial correlations with an environment provide a natural means of generating maps that are not completely positive [25–27] and, as we have shown, such correlations are a general feature of non-Markovian dissipative quantum mechanics when physically relevant initial states are considered.

ACKNOWLEDGMENTS

This work was supported partially by National Science Foundation Grant Nos. PHY-0426696 and PHY-0801368, Defense Advanced Research Projects Agency Grant No. DARPAHR0011-09-1-0008, and the Laboratory of Physical Sciences.

- [1] B. L. Hu, J. P. Paz, and Y. Zhang, *Phys. Rev. D* **45**, 2843 (1992).
- [2] B. L. Hu, A. Roura, and E. Verdaguier, *Phys. Rev. D* **70**, 044002 (2004).
- [3] In addition, even after resolving issues of renormalization, the system will typically be displaced by a finite amount of the order of the induced damping within a time of order $1/\Lambda$. This by itself could be easily dealt with (see Appendix C in Ref. [4]).
- [4] C. H. Fleming, A. Roura, and B. L. Hu, *Ann. Phys. (NY)* **326**, 1207 (2011).
- [5] H. Grabert, P. Schramm, and G. L. Ingold, *Phys. Rep.* **168**, 115 (1988).
- [6] L. D. Romero and J. P. Paz, *Phys. Rev. A* **55**, 4070 (1997).
- [7] J. R. Anglin, J. P. Paz, and W. H. Zurek, *Phys. Rev. A* **55**, 4041 (1997).
- [8] H. P. Breuer, B. Kappler, and F. Petruccione, *Ann. Phys. (NY)* **291**, 36 (2001).
- [9] N. Kampen and I. Oppenheim, *J. Stat. Phys.* **87**, 1325 (1997).
- [10] H. P. Breuer, A. Ma, and F. Petruccione, in *Quantum Computing and Quantum Bits in Mesoscopic Systems*, edited by A. J. Leggett, B. Ruggiero, and P. Silvestrini (Kluwer, Dordrecht, 2003).
- [11] W. T. Strunz and T. Yu, *Phys. Rev. A* **69**, 052115 (2004).
- [12] W. T. Pollard, A. K. Felts, and R. A. Friesner, *Adv. Chem. Phys.* **93**, 77 (1997).
- [13] H. J. Carmichael, *Statistical Methods in Quantum Optics I* (Springer, New York, 1999).
- [14] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, New York, 2002).
- [15] N. G. van Kampen, *Adv. Chem. Phys.* **34**, 245 (2007).
- [16] The case with $\langle \bar{\mathbf{I}}_n(t) \rangle_E \neq 0$ [see Eqs. (8) and (12) for the definitions] can be included in our treatment by shifting the operator $\mathbf{I}_n(t) \rightarrow \mathbf{I}_n(t) - \langle \bar{\mathbf{I}}_n(t) \rangle_E$ and including the term $\sum_n \mathbf{L}_n(t) \langle \bar{\mathbf{I}}_n(t) \rangle_E$, which depends on the environment initial state, in the system Hamiltonian.
- [17] C. H. Fleming and B. L. Hu (unpublished).
- [18] R. Kubo, *J. Phys. Soc. Jpn.* **12**, 570 (1957).
- [19] P. C. Martin and J. Schwinger, *Phys. Rev.* **115**, 1342 (1959).
- [20] G. Pérez-Nadal, A. Roura, and E. Verdaguier, *Phys. Rev. D* **77**, 124033 (2008).
- [21] G. Pérez-Nadal, A. Roura, and E. Verdaguier, *Classical Quantum Gravity* **25**, 154013 (2008).
- [22] C. H. Fleming and N. I. Cummings, *Phys. Rev. E* **83**, 031117 (2011).
- [23] A. M. Kuah, K. Modi, C. A. Rodríguez-Rosario, and E. C. G. Sudarshan, *Phys. Rev. A* **76**, 042113 (2007).
- [24] K. Modi and E. C. G. Sudarshan, *Phys. Rev. A* **81**, 052119 (2010).
- [25] T. F. Jordan, A. Shaji, and E. C. G. Sudarshan, *Phys. Rev. A* **70**, 052110 (2004).
- [26] M. Ziman, e-print [arXiv:quant-ph/0603166](https://arxiv.org/abs/quant-ph/0603166).
- [27] H. A. Carteret, D. R. Terno, and K. Życzkowski, *Phys. Rev. A* **77**, 042113 (2008).