

Supplemental Figure 1:

Effect of pseudocanthatidins on several alternative exons. HEK293 cells were treated with 1 mM of the substances indicated for 24 hrs and effects in alternative splicing were assayed by RT-PCR. The sequences of the alternative exons were as follows:

CAMKK2	CTGCACCCCTGGGTCA CGAGGCATGGGGCGGAGCCGTTGCCGTCGGAGGATGAGAA CTGCACGCTGGTCAAGTGA CTGAAGAGGAGGTCGAGAACTCAGTCAAACACATTC CCAGCTTGGCAACCGTG
IDH3B	GTGCGGACTCGAGACATGGGCGGCTACAGCACCAACCGACTTCATCAAGTCTGT CATCGGTACACTGCAGACTAAAGGGAGCTAGAGCCTTATTCTTCCAACCTTGCA AGGACCACACTCCCCATACCCCTCAGTGCAGTGTACCAGGGAAGAGACCTTGTGCCT CTAACGAGTGGACCATGGTCACCTGCTGGTAGAGCCTAGGTTGTCCCTGGGCCGG CTTCCTTAGGGGACAGACTGTTGGGTGGTATGGGGATTGTTAGGATGGAGCCCAG GCCACATGGATGATGATTCTCCCCACAG
SMNDC1	gtaagtgtaaaaaaaactctaactataacatgaataataaaaatacaatggtaagaagtttattcagtttaactaccctatttaatcttgaa cttcatttaggcagttatgattatagccaaatttgcatttgcattaaataca
TIMM8B	Gccacgtgcccgtagaaaagatactcatccactgtgggttgggtcgccgtccccactgcctactggattgtgaggatcatatgcga caatgtatttggaaacgactagaacattatcgaggaggaagggtgactctgaagttagtgcgttagactatggatgtagaacaagggttggag ccctcggacatggtctaaacgcggcctgacttgcggctacatgacccctggactacataatcacgcctttaatggggagggtgatgacag ctatccttgaggaccccttagagagaactgatttgcatttgcacaaata
ZNF207	ATGGGGACACCTGTCA CAAGCTCAAGTACAGCTTCATCCAATT CAGAAAGTCTGTCT GCATCTTCTAAAGCTCTGTTCTAGCACAGCACAA

Supplemental data

